

# Sussex Research

## Computational model of a modulatory cell type in the feeding network of the snail, *Lymnaea stagnalis*

Dimitris V. Vavoulis, Eugeny S. Nikitin, Jianfeng Feng, Paul Benjamin, George Kemenes

### Publication date

06-07-2007

### Licence

This work is made available under the **Copyright not evaluated** licence and should only be used in accordance with that licence. For more information on the specific terms, consult the repository record for this item.

### Citation for this work (American Psychological Association 7th edition)

Vavoulis, D. V., Nikitin, E. S., Feng, J., Benjamin, P., & Kemenes, G. (2007). *Computational model of a modulatory cell type in the feeding network of the snail, Lymnaea stagnalis* (Version 1). University of Sussex. <https://hdl.handle.net/10779/uos.23312603.v1>

### Published in

BMC Neuroscience

### Link to external publisher version

<https://doi.org/10.1186/1471-2202-8-S2-P113>

### Copyright and reuse:

This work was downloaded from Sussex Research Open (SRO). This document is made available in line with publisher policy and may differ from the published version. Please cite the published version where possible. Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners unless otherwise stated. For more information on this work, SRO or to report an issue, you can contact the repository administrators at [sro@sussex.ac.uk](mailto:sro@sussex.ac.uk). Discover more of the University's research at <https://sussex.figshare.com/>

# The ionic basis of action potentials in an identified modulatory cell type in the feeding network of the snail, *Lymnaea stagnalis*: a computational study

Dimitris V. Vavoulis<sup>1,2</sup>, Eugeny S. Nikitin<sup>2</sup>, Jianfeng Feng<sup>1</sup>, Paul R. Benjamin<sup>2</sup>, György Kemenes<sup>2</sup>

<sup>1</sup>*Ctr. for Scientific Computing, Uni. Of Warwick, Coventry, United Kingdom*

<sup>2</sup>*Sussex Ctr. for Neuroscience, Uni. of Sussex, Brighton, United Kingdom*

email: [Dimitris.Vavoulis@dcs.warwick.ac.uk](mailto:Dimitris.Vavoulis@dcs.warwick.ac.uk)

## Introduction

Realistic mathematical models of single neurons are significant in assessing the contribution of specific ionic conductances to neuronal excitability. This study presents a detailed computational model of the Cerebral Giant Cells (CGCs), a pair of serotonergic neurons in the feeding network of *Lymnaea stagnalis*, which are critical for the expression of motor behaviour (feeding) and the formation of long-term memory.

## Methods

First, we fitted a single-compartment, Hodgkin-Huxley model of the CGCs to two-electrode voltage- and current-clamp data [1] using a combination of linear and non-linear least-square fitting techniques. Then, we selectively blocked each ionic current to assess its role in the model, thus mimicking the application of pharmacological agents in the biological neuron.

## Results

The model replicates accurately the shape of the action potentials and the tonic firing ( $\sim 0.74$  Hz) of the biological neuron (Fig. 1A). A persistent sodium current  $I_{NaP}$  and a transient low-threshold calcium current  $I_{LVA}$  keep the neuron spontaneously active (Fig. 1Bi,ii). A transient potassium current  $I_A$  regulates the interspike interval, while a transient high-threshold calcium current  $I_{HVA}$  increases the duration of each spike (Fig. 1Biii,iv). Transient sodium and delayed rectifier potassium currents are responsible for the depolarizing and repolarizing phases of the action potential, as in the classical Hodgkin-Huxley model. The available experimental data [1] are in agreement with these conclusions.

## Conclusions


The model we have developed here provides an accurate description of the CGCs at the biophysical level and it is a useful tool for studying the electrical properties of these important modulatory neurons.

## Acknowledgments

This research was supported by EPSRC and BBSRC, United Kingdom

## References

1. K. Staras, J. Gyori & G. Kemenes: **Voltage-gated ionic currents in an identified modulatory cell type controlling molluscan feeding.** *Eur J Neurosci* 2002, **15**: 109-119



**Fig. 1:** Overview of the CGCs model and the contribution of specific currents to neuronal excitability. In A, the model has been slightly (2msec) shifted to the right for illustration purposes