

Sussex Research

Flavor violation and warped geometry

Stephan Huber

Publication date

19-07-2003

Licence

This work is made available under the **Copyright not evaluated** licence and should only be used in accordance with that licence. For more information on the specific terms, consult the repository record for this item.

Citation for this work (American Psychological Association 7th edition)

Huber, S. (2003). *Flavor violation and warped geometry* (Version 1). University of Sussex.
<https://hdl.handle.net/10779/uos.23312687.v1>

Published in

Nuclear Physics B

Link to external publisher version

[https://doi.org/10.1016/S0550-3213\(03\)00502-9](https://doi.org/10.1016/S0550-3213(03)00502-9)

Copyright and reuse:

This work was downloaded from Sussex Research Open (SRO). This document is made available in line with publisher policy and may differ from the published version. Please cite the published version where possible. Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners unless otherwise stated. For more information on this work, SRO or to report an issue, you can contact the repository administrators at sro@sussex.ac.uk. Discover more of the University's research at <https://sussex.figshare.com/>

Huber, S.J.

Flavor violation and warped geometry

Article (Unspecified)

Original citation:

Huber, S.J. (2003) Flavor violation and warped geometry. Nuclear Physics B, 666 (1-2). pp. 269-288. ISSN 0550-3213

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/1637/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository URL above for details on accessing the published version

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Flavor violation and warped geometry

Stephan J. Huber¹

Deutsches Elektronen-Synchrotron DESY, Hamburg, Germany

Abstract

Extra dimensions have interesting consequences for flavor physics. We consider a setup where the standard model fermions and gauge fields reside in the bulk of a warped extra dimension. Fermion masses and mixings are explained by flavor dependent fermion locations, without relying on hierarchical Yukawa couplings. We discuss various flavor violating processes induced by (Kaluza-Klein) gauge boson exchange and non-renormalizable operators. Experimental constraints are satisfied with a Kaluza-Klein scale of about 10 TeV. Some processes, such as muon-electron conversion, are within reach of next generation experiments.

¹stephan.huber@desy.de

1 Introduction

Models with extra dimensions have attracted great attention in recent years as they offer new perspectives on challenging problems in modern physics. It was demonstrated by Randall and Sundrum that a small but warped extra dimension provides an elegant solution to the gauge hierarchy problem [1] (see also [2]). The fifth dimension is an S_1/Z_2 orbifold with an AdS_5 geometry of curvature k , bordered by two 3-branes with opposite tensions and separated by distance R . The red shift induced by the AdS warp factor $\Omega = e^{-\pi k R}$ generates an exponential hierarchy between the energy scales on the two branes. If the brane separation is $kR \simeq 11$, the scale on the negative tension brane is of TeV-size, while the scale on the other brane is of order M_{Pl} . The AdS curvature k and the 5D Planck mass M_5 are assumed to be of order M_{Pl} . Gravity is weak at the TeV-brane because the zero mode corresponding to the 4D graviton is localized at the positive tension brane (Planck-brane).

In contrast to the original proposal [1] we take the standard model (SM) fermions and gauge bosons as bulk fields. In the non-supersymmetric framework we are studying, the Higgs field has to be confined to the TeV-brane in order to preserve the solution to the gauge hierarchy problem [3, 4]. Our motivation is twofold. With the SM fermions residing in the 5-dimensional bulk, the hierarchy of quark and lepton masses can be related to a higher dimensional geography [5, 6]. Different fermion flavors are localized at different positions in the extra dimension or, more precisely, have different wave functions. The fermion masses are in direct proportion to the overlap of their wave functions with the Higgs field [7]. Also the CKM mixing can be explained along these lines. A second point is that at the TeV-brane non-renormalizable operators, now only TeV-scale suppressed, are known to induce rapid proton decay, large neutrino masses and flavor violating interactions. Because of the warp factor, the effective cut-off scale varies along the extra dimension. If the quarks and leptons are localized towards the Planck-brane in the extra dimension, the suppression scales of dangerous operators can be significantly enhanced [5, 6]. Small Majorana neutrino masses can then arise from dimension-five interactions, without introducing new degrees of freedom [8]. The atmospheric and solar neutrino anomalies can be satisfactorily resolved. Alternatively, Dirac neutrino masses can be generated by a coupling to right-handed neutrinos in the bulk [9, 10].

Fields living in the 5D bulk can be expanded as a tower of Kaluza-Klein (KK) states. Electroweak observables, in particular the weak mixing angle and gauge boson masses, require the KK excitations of SM particles to be heavier than about 10 TeV [4, 11, 12]. A small hierarchy separates the weak and KK scales. In the presence of brane-localized kinetic terms the bound on the KK scale may be somewhat relaxed [13], but we are not considering this possibility in the following. For fermions confined to the TeV-brane, the electroweak constraints have been analyzed in ref. [14].

In this paper we address the issue of flavor violation in the warped SM which is an

immediate consequence of our approach to the fermion mass problem. Some aspects of this topic have already been discussed in the literature [5, 6, 12, 15–18]. The crucial connection to the issue of fermion masses, however, has not been thoroughly investigated so far. This connection allows us to obtain more reliable predictions for the flavor violating rates. Moreover, we are including lepton flavor violating processes which have not yet been considered in context of the warped SM. Some results have already been presented in ref. [19].

In the next section we summarize some results on bulk fermions and gauge bosons in a warped background. In section 3 we are using the observed fermion masses and mixings to determine the locations of the various fermion flavors in the extra dimension. The fermion mass pattern can be accommodated without relying on hierarchical Yukawa couplings. In section 4 we discuss how the mixing between different KK levels leads to a non-unitary CKM matrix. Flavor violation by (KK) gauge boson exchange, being a natural consequence of our setup, is studied in section 5. We find that with a KK scale of 10 TeV the model is consistent with the experimental constraints, while some processes, such as muon-electron conversion, are within reach of next generation experiments. In section 6 we show that contributions from non-renormalizable operators to flavor violating processes are naturally within experimental bounds, while proton decay cannot be adequately suppressed. Finally, in section 7 we conclude.

2 Bulk fields in a warped background

We assume that the standard model gauge bosons and fermions reside in the bulk of the warped 5D space-time [1]

$$ds^2 = e^{-2\sigma(y)} \eta_{\mu\nu} dx^\mu dx^\nu + dy^2, \quad (2.1)$$

where $\sigma(y) = k|y|$. The 4-dimensional metric is $\eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1)$ and y denotes the fifth coordinate. The AdS curvature k is related to the bulk cosmological constant and the brane tensions. To set the notation let us briefly review some properties of gauge and fermion fields in a slice of AdS_5 .

A gauge field propagating in a curved background with metric G^{MN} is described by the equation of motion

$$\frac{1}{\sqrt{-G}} \partial_M (\sqrt{-G} G^{MN} G^{RS} F_{NS}) - M_A^2 G^{RS} A_S = 0. \quad (2.2)$$

The mass term M_A^2 arises from spontaneous symmetry breaking and is present only for the weak gauge bosons. Since the Higgs field is localized at the TeV-brane, we have

$$M_A^2 = \frac{1}{2} (g^{(5)})^2 v_0^2 \delta(y - \pi R), \quad (2.3)$$

where $g^{(5)}$ is the 5D gauge coupling. The Higgs vev v_0 is expected to be of order M_{Pl} . Imposing the gauge $A_5 = 0$, we decompose the 5D fields as

$$A_\mu(x, y) = \frac{1}{\sqrt{2\pi R}} \sum_{n=0}^{\infty} A_\mu^{(n)}(x) f_n^A(y). \quad (2.4)$$

Inserting the metric (2.1) and the decomposition (2.4) into the equation of motion (2.2), the wave functions f_n^A have to satisfy

$$(\partial_y^2 - 2\sigma' \partial_y - M_A^2 + e^{2\sigma} m_n^2) f_n^A = 0, \quad (2.5)$$

where $\sigma' = d\sigma/dy$. This equation is solved by [20]

$$f_n^A(y) = \frac{e^\sigma}{N_n} \left[J_1\left(\frac{m_n}{k} e^\sigma\right) + b^A(m_n) Y_1\left(\frac{m_n}{k} e^\sigma\right) \right]. \quad (2.6)$$

The spectrum of KK masses m_n and the coefficients b^A are determined by the boundary conditions of the wave functions at the branes. The localized mass term M_A^2 only affects the boundary condition at the TeV-brane [4, 11]. The normalization constants are fixed by

$$\frac{1}{2\pi R} \int_{-\pi R}^{\pi R} dy f_m^A(y) f_n^A(y) = \delta_{mn}. \quad (2.7)$$

In the case $M_A^2 = 0$ eq. (2.2) admits a constant zero mode solution [20]. The excited states are localized towards the TeV-brane and have TeV-scale KK masses $m_n \sim n\pi k e^{-\pi k R}$. In the presence of the mass term M_A^2 , the zero mode acquires a mass

$$m_0^2 = (g^{(5)})^2 v_0^2 e^{-2\pi k R} / (2\pi R) + \mathcal{O}((g^{(5)})^4 v_0^4 e^{-4\pi k R} / M_{KK}^2), \quad (2.8)$$

where $M_{KK} = m_1$ denotes the KK scale. The order $1/M_{KK}^2$ corrections are related to a dip in the zero mode wave function caused by the boundary mass term [4]. Compared to the 4D standard model they induce a tree-level shift of the W and Z boson mass ratio relative to its 4D standard model value. In refs. [4, 11, 12] it was shown that the electroweak precision data thus implies the constraint $M_{KK} \gtrsim 10$ TeV. If the boundary mass term was not included in the KK reduction, but evaluated with the $M_A^2 = 0$ wave functions, the order $1/M_{KK}^2$ effects would show up in the 4D effective action as mixings between the KK states.

The equation of motion of a fermion in curved space-time reads

$$E_a^M \gamma^a (\partial_M + \omega_M) \Psi + m_\Psi \Psi = 0, \quad (2.9)$$

where E_a^M is the fünfbein, $\gamma^a = (\gamma^\mu, \gamma^5)$ are the Dirac matrices in flat space, and

$$\omega_M = \left(\frac{1}{2} e^{-\sigma} \sigma' \gamma_5 \gamma_\mu, 0 \right) \quad (2.10)$$

is the spin connection induced by the metric (2.1). Fermions have two possible transformation properties under the Z_2 orbifold symmetry, $\Psi(-y)_\pm = \pm \gamma_5 \Psi(y)_\pm$. Thus, $\bar{\Psi}_\pm \Psi_\pm$ is odd under Z_2 , and the Dirac mass term, which is odd as well, can be parametrized as $m_\Psi = c\sigma'$. The Dirac mass should therefore originate from the coupling to a Z_2 odd scalar field which acquires a vev. On the other hand, $\bar{\Psi}_\pm \Psi_\mp$ is even. Using the metric (2.1) one obtains for the left- and right-handed components of the Dirac spinor [5, 9]

$$[e^{2\sigma} \partial_\mu \partial^\mu + \partial_5^2 - \sigma' \partial_5 - M_f^2] e^{-2\sigma} \Psi_{L,R} = 0, \quad (2.11)$$

where $M_f^2 = c(c \pm 1)k^2 \mp c\sigma''$ and $\Psi_{L,R} = \pm \gamma_5 \Psi_{L,R}$.

Decomposing the 5D fields as

$$\Psi(x^\mu, y) = \frac{1}{\sqrt{2\pi R}} \sum_{n=0}^{\infty} \Psi^{(n)}(x^\mu) e^{2\sigma} f_n(y), \quad (2.12)$$

one ends up with a zero mode wave function [5, 9]

$$f_0(y) = \frac{e^{-c\sigma}}{N_0}, \quad (2.13)$$

and a tower of KK excited states

$$f_n(y) = \frac{e^{\sigma/2}}{N_n} \left[J_\alpha\left(\frac{m_n}{k} e^\sigma\right) + b_\alpha(m_n) Y_\alpha\left(\frac{m_n}{k} e^\sigma\right) \right]. \quad (2.14)$$

The order of the Bessel functions is $\alpha = |c \pm 1/2|$ for $\Psi_{L,R}$. The spectrum of KK masses m_n and the coefficients b_α are determined by the boundary conditions at the branes [5, 9]. The normalization constants follow from

$$\frac{1}{2\pi R} \int_{-\pi R}^{\pi R} dy e^\sigma f_m(y) f_n(y) = \delta_{mn}. \quad (2.15)$$

Because of the orbifold symmetry, the zero mode of Ψ_+ (Ψ_-) is a left-handed (right-handed) Weyl spinor. For $c > 1/2$ ($c < 1/2$) the fermion is localized near the boundary at $y = 0$ ($y = \pi R$), i.e. at the Planck- (TeV-) brane (see also fig. 1).

The zero modes of leptons and quarks acquire masses from their coupling to the Higgs field

$$\int d^4x \int dy \sqrt{-G} \lambda_{ij}^{(5)} H \bar{\Psi}_{i+} \Psi_{j-} \equiv \int d^4x m_{ij} \bar{\Psi}_{iL}^{(0)} \Psi_{jR}^{(0)} + \dots, \quad (2.16)$$

where $\lambda_{ij}^{(5)}$ are the 5D Yukawa couplings. The 4D Dirac masses are given by

$$m_{ij} = \int_{-\pi R}^{\pi R} \frac{dy}{2\pi R} \lambda_{ij}^{(5)} H(y) f_{0iL}(y) f_{0jR}(y) = \frac{l_{ij} v_0}{\pi k R} f_{0iL}(\pi R) f_{0jR}(\pi R). \quad (2.17)$$

In the second step we have used $H(y) = v_0 \delta(y - \pi R)/k$ and introduced the dimensionless couplings $l_{ij} = \lambda_{ij}^{(5)} \sqrt{k}$.

The gauge interaction between bulk gauge bosons and fermions, $g^{(5)} \bar{\Psi} i E_a^M \gamma^a A_M \Psi$, induces the effective 4D couplings [5]

$$g_{ijn} = \frac{g^{(5)}}{(2\pi R)^{3/2}} \int_{-\pi R}^{\pi R} e^\sigma f_i(y) f_j(y) f_n^A(y) dy \quad (2.18)$$

between the different KK levels. In the case of a massless gauge boson ($f_0^A(y) \equiv 1$) this integral reduces to the normalization condition of fermions (2.15), and one finds $g_{ij0} = \delta_{ij} g^{(5)} / \sqrt{2\pi R}$. For a massive gauge field the gauge coupling of the zero mode to TeV-brane fermions is somewhat reduced due to the dip in its wave function [4]. In the SM this effect leads, for instance, to smaller gauge couplings of W and Z bosons relative to that of the photon. In order to keep these corrections within experimental bounds we derived in ref. [11] the constraint $M_{KK} \gtrsim 60$ TeV if the fermions reside on the TeV-brane. For fermions localized towards the Planck-brane this constraint becomes weaker than the 10 TeV bound from the gauge boson masses discussed before.

Combining eqs. (2.8) and (2.18), we can determine the Higgs vev required to provide the measured gauge boson masses. To keep the electroweak corrections small enough, we take $M_{KK} = 10$ TeV. Assuming $k = M_{\text{Pl}}$ we find $kR = 10.83$ and $v_0 = 0.043 M_{\text{Pl}}$. Thus, there is still a small hierarchy of about twenty between the fundamental scale and the Higgs vev.

3 Quark and lepton masses

The fermion masses crucially depend on the overlap of the fermion wave functions and the Higgs profile. They are functions of the 5D mass parameters of the left- and right-handed fermions, c_L and c_R respectively, which enter eq. (2.17). As the 5D Dirac mass, i.e. c parameter increases, the fermion gets localized closer towards the Planck-brane. Its overlap with the Higgs profile at the TeV-brane is reduced, which is reflected in a smaller 4D fermion mass from electroweak symmetry breaking. Since the fermionic zero modes depend exponentially on the 5D mass parameters, the large hierarchy of charged fermion masses can be generated from c parameters of order unity [5, 6]. If right-handed neutrinos are introduced in the bulk, sub-eV neutrino masses can be explained in the same manner [9, 10].

Building up the fermion mass matrices from eq. (2.17) requires also the specification of the 5D Yukawa couplings. Thus there are considerably more independent parameters in the model than there are observable fermion masses and mixings. Relating the measured fermion properties to their locations in the extra dimension needs therefore additional assumptions. In ref. [6] it was assumed that the 5D Yukawa couplings are of order unity, and the fermion mass pattern is solely due to the different locations. Additionally, the fermions were localized as closely as

possible towards the Planck-brane in order to maximally suppress the impact of non-renormalizable operators on rare processes.

In the following we are using a somewhat different approach. While still assuming the 5D Yukawa couplings to be of order unity, we are looking for a set of c parameters which “most naturally” accounts for the observed fermion masses and mixings. More precisely, we are taking random 5D Yukawa couplings and require the averaged fermion properties to fit the experimental data. Similar methods have been used to study models where the fermion mass pattern is due to (approximately) conserved charges [21]. There the role of the fermion locations is taken by a set of Higgs fields.

Let us first focus on the quark sector. At the scale of 10 TeV, where we are matching our model to the observational data, the (running) quark masses are

$$\begin{aligned} m_u &= 0.7 - 2.3 \text{ MeV}, & m_c &= 420 - 540 \text{ MeV}, & m_t &= 140 - 148 \text{ GeV} \\ m_d &= 1.4 - 4.2 \text{ MeV}, & m_s &= 28 - 80 \text{ MeV}, & m_b &= 2.1 - 2.3 \text{ GeV}. \end{aligned} \quad (3.19)$$

We have used one loop renormalization group equations [22] to run the quark masses given in ref. [23] from the scale M_Z to 10 TeV. The running reduces the quark masses by about 20 percent. The moduli of the CKM matrix are given by [24]

$$|V_{\text{CKM}}| = \begin{pmatrix} 0.9741 - 0.9756 & 0.219 - 0.226 & 0.0025 - 0.0048 \\ 0.219 - 0.226 & 0.9732 - 0.9748 & 0.038 - 0.044 \\ 0.004 - 0.014 & 0.037 - 0.044 & 0.9990 - 0.9993 \end{pmatrix}. \quad (3.20)$$

A convention independent measure of CP violation is the Jarlskog invariant which experimentally is found to be [24]

$$J = (3 \pm 0.3) \times 10^{-5}. \quad (3.21)$$

Assuming non-hierarchical 5D Yukawa couplings, the fermion mass matrix (2.17) leads to a product-like structure

$$M \sim \begin{pmatrix} a_1 b_1 & a_1 b_2 & a_1 b_3 \\ a_2 b_1 & a_2 b_2 & a_2 b_3 \\ a_3 b_1 & a_3 b_2 & a_3 b_3 \end{pmatrix} \quad (3.22)$$

where a_i and b_i are given by the fermion wave functions $f_{0iL,R}(\pi R)$. As a function of the fermion location there is a slow increase $f_0(\pi R) \propto \sqrt{1/2 - c}$ for $c < 1/2$, and an exponential suppression $f_0(\pi R) \propto \exp(-c\pi k R)$ for $c > 1/2$. Note that some non-equal Yukawa couplings are needed to render the mass matrix (3.22) non-singular. If the mass matrix is diagonalized by $U_L M U_R^\dagger$, the left- and right-handed mixings are typically of order $U_{L,ij} \sim a_i/a_j$ and $U_{R,ij} \sim b_i/b_j$, respectively. Fermions which have similar positions (c parameters) have large mixings. The mass matrix (3.22) predicts the approximate relation $U_{13} \sim U_{12}U_{23}$ between the mixing angles, which for the observed CKM matrix (3.20) is satisfied up to a factor of about two.




Figure 1: The wave functions of the left-handed quark zero modes $q_i^{(0)}$ and the first KK state, $q_1^{(1)}$, of Q_1 for the parameters of eq. (3.24).

In our numerical analysis we generate $N = 25000$ random sets of up and down-type Yukawa couplings and diagonalize the emerging mass matrices (2.17). We require the averaged fermion masses and mixings to fit the experimental data, using the logarithmic average of a quantity X

$$\langle X \rangle = \exp \left(\sum_i^N \frac{\ln(X_i)}{N} \right). \quad (3.23)$$

Taking $2/3 < |l_{ij}| < 4/3$ and random phases from 0 to 2π , we find the “most natural” locations

$$\begin{aligned} c_{Q1} &= 0.643, & c_{D1} &= 0.643, & c_{U1} &= 0.671, \\ c_{Q2} &= 0.583, & c_{D2} &= 0.601, & c_{U2} &= 0.528, \\ c_{Q3} &= 0.317, & c_{D3} &= 0.601, & c_{U3} &= -0.460. \end{aligned} \quad (3.24)$$

The wave functions of the left-handed quark zero modes and the first excited state of Q_1 are shown in fig. 1. We fix the relative positions of Q_1 and Q_2 by fitting $|V_{us}|$, while $|V_{cb}|$ determines the relative positions of Q_2 and Q_3 . The Jarlskog invariant $J = \text{Im}(V_{cs}V_{us}^*V_{ud}V_{cd}^*)$ [25] and $|V_{ub}|$ are then fixed as well. Note that the CKM mixings are determined by the locations of the left-handed quarks. We then use the locations of the right-handed quarks to fit the quark masses. Taking the quark locations (3.24), we find for the averages

$$\begin{aligned} m_u &= 2.0 \text{ MeV}, & m_c &= 506 \text{ MeV}, & m_t &= 144 \text{ GeV}, \\ m_d &= 4.0 \text{ MeV}, & m_s &= 58 \text{ MeV}, & m_b &= 2.2 \text{ GeV}, \\ |V_{us}| &= 0.222, & |V_{cb}| &= 0.040, & |V_{ub}| &= 0.0088, \\ J &= 3.1 \times 10^{-5}. \end{aligned} \quad (3.25)$$

While the prediction of J agrees well with the experimental value (3.21), $|V_{ub}|$ is found to be about two times too large (3.20). As discussed above this discrepancy is a consequence of the mass pattern (3.22).

The quark locations of eq. (3.24) are not a unique solution. Moving all the left-handed quarks in such a way that the related factors a_i in (3.22) are changed by a common factor, can be compensated by an appropriate change in the locations of the right-handed quarks. So in fact there is a one parameter family of solutions, labeled, for instance, by δc_{Q1} . However, if some of the light quarks are localized towards the TeV-brane ($c < 1/2$), large deviations from electroweak precision variables drive the KK scale far above 10 TeV [11]. In section 5 we will see that in this case flavor violation is greatly enhanced. Since c_{U2} is already close to $1/2$, we cannot move the right-handed quarks much closer to the TeV-brane. This would also require to localize the right-handed top quark extremely close towards the TeV-brane by $c_{U3} \lesssim -2$. On the other hand, the Q_i cannot be localized much closer towards the TeV-brane without getting too large modifications of the left-handed bottom couplings. Thus we end up with the quite constrained range of $-0.01 \lesssim \delta c_{Q1} \lesssim 0.02$. The quarks can be localized closer towards the Planck-brane if the 5D Yukawa couplings are increased by a common factor. Increasing them by a factor of 10, for instance, corresponds to $\delta c_{Q1} = 0.037$. However, very large Yukawa couplings $l \gg 1$ introduce a new and unexplained hierarchy in the model. Taking Yukawa couplings $l \ll 1$ would mean to move the quarks closer towards the TeV-brane which worsens the electroweak fit. The quark locations also depend on k/M_{Pl} which we take to be one.

The experimental errors (3.19) and (3.20) translate into uncertainties in the fermion locations. For instance, from $|V_{cb}|$ we can fix the position of Q_3 relative to Q_2 only up to an error of $-0.02 \lesssim \Delta c \lesssim 0.03$. The position of Q_1 relative to Q_2 is determined quite accurately up to about $-6 \times 10^{-4} \lesssim \Delta c \lesssim 5 \times 10^{-4}$ by $|V_{us}|$. For the right-handed quarks the errors induced by the quark masses range from $\Delta c \approx \pm 8 \times 10^{-4}$ for D_3 to $\Delta c \approx \pm 0.03$ for U_3 . More precisely, these are errors on the relative positions of left- and right-handed quarks, for instance, $c_{Q3} - c_{U3}$ in the case of the top. Note that positions $c < 1/2$ have larger errors since in this range the fermion mass does not depend exponentially but as $\sqrt{1/2 - c}$ on the fermion location.

The statistical approach provides us with distributions of fermion properties rather than with their precise values. This adds further uncertainties to the fermion locations. In fig. 2a we present the distribution of $\ln(m_u)$ which shows a half width of about 1.1. Requiring that, with a different location, the average value of $\ln(m_u)$ still lies within this half width range translates into an uncertainty of $-0.017 \lesssim \Delta c_{U1} \lesssim 0.019$. In table 1 we have summarized the statistical and experimental uncertainties in the quark locations. Fig. 2b shows the distribution of $\ln(|V_{ub}|)$ which overlaps considerably with the experimentally favored range. Despite the factor of two deviation in the average $\langle |V_{ub}| \rangle$, there is an acceptably good fit to the data once the statistical variations are taken into account.




Figure 2: Logarithmic distributions of m_u (a) and $|V_{ub}|$ (b) for Yukawa coupling distributions $2/3 < |l_{ij}| < 4/3$ (solid lines), $0 < |l_{ij}| < 2$ (dashed dotted lines) and $|l_{ij}| = 1$ (solid lines). The vertical dashed lines indicate the experimental uncertainties.

Our results depend only weakly on the range taken for the moduli of the 5D couplings l_{ij} . In fig. 2 we present the distributions of m_u and $|V_{ub}|$ additionally for the case $0 < |l_{ij}| < 2$ and for pure phase Yukawa couplings $|l_{ij}| = 1$. In the context of flat extra dimensions pure phase Yukawa couplings have recently been investigated in ref. [26]. We find that a wider range of $|l_{ij}|$ leads only to a slight broadening in the distributions of fermion properties. The induced shift in the average values is always much smaller than the width of the distributions. For instance, we find an average $m_u = 2.3$ (2.0) MeV in the case $0 < |l_{ij}| < 2$ ($|l_{ij}| = 1$). This behavior stresses that the fermion mass pattern is indeed determined by the fermion locations, not by accidental properties of the 5D Yukawa matrices. Note that the robustness of our results concerning different ranges of $|l_{ij}|$ is due to the complex phases in the 5D Yukawa couplings. For real values of l_{ij} there is a high probability of cancellations in the mass matrices. This leads to multiple peaks in the distributions of fermion masses and mixings for a narrow range of $|l_{ij}|$.

Let us close this section with a brief discussion of the lepton sector. To determine the lepton locations we have to take into account neutrino masses and

	Q_1	Q_2	Q_3	D_1	D_2	D_3	U_1	U_2	U_3
c	0.643	0.583	0.317	0.643	0.601	0.601	0.671	0.528	-0.460
exp.	+0.0004 -0.0006	+0.002 -0.003	+0.03 -0.02	+0.017 -0.008	+0.012 -0.006	+0.0008 -0.0008	+0.017 -0.002	+0.004 -0.002	+0.03 -0.03
stat.	+0.022 -0.027	+0.029 -0.024	+0.13 -0.21	+0.016 -0.023	+0.006 -0.017	+0.004 -0.006	+0.019 -0.017	+0.006 -0.023	+0.50 -0.76

Table 1: Experimental and statistical errors on the fermion locations of eq. (3.24).

mixings. Neutrino masses can arise from different sources in the context of warped geometry. In one scenario sub-eV Dirac masses are generated by a coupling to right-handed neutrinos in the bulk [9, 10]. The necessary tiny Yukawa couplings can naturally be generated if the right-handed neutrinos are localized closely towards the Planck-brane by taking $1.2 \lesssim c \lesssim 1.5$. Neutrino and charged fermion masses are then treated on the same footing. Alternatively, the dimension five interaction $(1/Q)HHLL$ can induce small Majorana neutrino masses [8]. The suppression scale Q of the non-renormalizable interaction depends on the position of the left-handed leptons L in the extra dimension. This mechanism is minimal in the sense that it does not require to extend the standard model particle content.

As in the case of quarks, the locations of the left-handed states determine the observed fermion mixings. Large neutrino mixings require the neutrinos and thus the $SU(2)$ lepton doublets to have similar positions c_{Li}

$$\begin{aligned} c_{L1} &= 0.565, & c_{L2} &= 0.565, & c_{L3} &= 0.565, \\ c_{E1} &= 0.764, & c_{E2} &= 0.609, & c_{E3} &= 0.509. \end{aligned} \quad (3.26)$$

The right-handed positions c_{Ei} we fixed by requiring that with random Yukawa couplings the average charged lepton masses fit their observed values.

To suppress the matrix element U_{e3} in the neutrino mixing matrix it is favorable to separate the electron doublet somewhat from the muon and tau doublets [8]

$$\begin{aligned} c_{L1} &= 0.631, & c_{L2} &= 0.565, & c_{L3} &= 0.565, \\ c_{E1} &= 0.725, & c_{E2} &= 0.594, & c_{E3} &= 0.497. \end{aligned} \quad (3.27)$$

Like in the case of quarks there is a one parameter family of degenerate locations if the left- and right-handed states are shifted against each other. The widths of the distributions allow to fix the locations only up to $\Delta c \approx \pm 0.01$. Taking the 5D Yukawa couplings larger than one, we could shift the leptons closer towards the Planck-brane, as was done in ref. [8].

4 Mixings of KK states and the unitarity of the CKM matrix

So far we have only dealt with the zero modes of quarks and leptons. But the Yukawa interaction (2.16) also induces mixings between the zero modes and the vector-like excited states. In the following we show that this effect hardly modifies the conclusions we have reached above.

From the KK reduction of an $SU(2)$ doublet quark Q_L we obtain a left-handed zero mode $q_L^{(0)}$, corresponding to the SM quark, and an infinite tower of left- and right-handed KK excited states $q_L^{(n)}$ and $q_R^{(n)}$, where we omit flavor indices. The $SU(2)$ singlet up quarks decompose into the right-handed zero mode $u_R^{c(0)}$ and the

KK excited states $u_L^{c(n)}$ and $u_R^{c(n)}$. After electroweak symmetry breaking the up quark mass matrix takes the form

$$M_U = (\bar{u}_L^{(0)}, \bar{u}_L^{(1)}, \bar{u}_L^{c(1)}, \dots) \begin{pmatrix} m^{(0,0)} & 0 & m^{(0,1)} & \dots \\ m^{(1,0)} & m_{Q,1} & m^{(1,1)} & \dots \\ 0 & 0 & m_{U,1} & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} \begin{pmatrix} u_R^{c(0)} \\ u_R^{(1)} \\ u_R^{c(1)} \\ \vdots \end{pmatrix} \quad (4.28)$$

where we again suppress flavor indices, i.e. every entry represents a 3×3 matrix in flavor space. The masses $m^{(m,n)}$ arise from electroweak symmetry breaking. They are obtained by inserting the relevant wave functions into eq. (2.17). The $m_{Q,m}$ and $m_{U,m}$ denote the KK masses of the excited quark states. The zeros in (4.28) follow from the Z_2 orbifold properties of the wave functions. Analogous mass matrices arise for the down quarks and charged leptons. The wave functions of the the KK fermions are localized at the TeV-brane (see fig. 1). Their overlap with the Higgs is large and almost independent of the location of the zero mode. We obtain

$$\begin{aligned} m^{(0,n)} &= \frac{lv_0}{\pi k R} f_{0L}(\pi R) f_{nR}(\pi R) \\ m^{(m,n)} &= \frac{lv_0}{\pi k R} f_{mL}(\pi R) f_{nR}(\pi R) \approx (-1)^{m+n} \cdot l \cdot 349 \text{ GeV}, \quad m, n \geq 1. \end{aligned} \quad (4.29)$$

These masses are large compared to the corresponding zero mode masses.

The mass matrix (4.28) induces a mixing between SU(2) singlet and doublet states. This effect diminishes the weak charge of the left-handed quarks [16]. In the context of neutrinos this behavior was discussed in refs. [9, 10]. Ignoring flavor mixing for a moment, the singlet admixture in the left-handed zero mode is

$$\sum_{n=1}^{\infty} \left(\frac{m^{(0,n)}}{m_{U,n}} \right)^2 \approx \left(\frac{m^{(0,1)}}{m_{U,1}} \right)^2 \sum_{n=1}^{\infty} \frac{1}{n^2} = \left(\frac{m^{(0,1)}}{m_{U,1}} \right)^2 \frac{\pi^2}{6}.$$

In the second step we approximated the KK spectrum by $m_n \approx nm_1$. Taking the quark positions of eq. (3.24) we obtain for the top quark a singlet admixture of 1.5×10^{-4} . For the other quark flavors the admixture is smaller than 10^{-6} . The mixing between the SU(2) doublet zero mode and its KK excitations is even more suppressed and on the order of $(m^{(0,1)} m^{(1,1)})^2 / m_{Q,1}^4$. Our neglect of the KK states in the discussion of quark and lepton masses in the previous section is therefore well justified.

The gauge couplings of fermions (2.18) are modified by KK mixing in the weak gauge boson sector [4, 11]. The W and Z bosons couple somewhat weaker to fermions at the TeV-brane, while their KK states couple stronger [5, 20], as we show in fig. 3. For $c > 1/2$ the gauge coupling is almost independent of the precise location, and the coupling of the KK states is small, $g_1/g_{SM} \approx 0.19$.




Figure 3: Gauge couplings of the Z boson (a) and its first KK state relative to the SM value as a function of the fermion location.

The charged current interaction of the KK fermions is described by an infinitely dimensional version of the CKM matrix

$$\mathcal{V} = \mathcal{U}_{L,U} \mathcal{G} \mathcal{U}_{L,D}^\dagger \quad (4.30)$$

where \mathcal{U}_L and \mathcal{U}_R diagonalize the full fermion mass matrices (4.28). The flavor diagonal matrix \mathcal{G} contains the gauge couplings of the various KK states to the W boson. We normalize \mathcal{G} to the weak coupling of the muon as obtained from eq. (2.18) with the muon location of eq. (3.26). Since the weak gauge bosons are massive, they induce couplings between different KK levels. These couplings are suppressed by M_Z/M_{KK} and numerically small. Note that in general even the full CKM matrix \mathcal{V} is not unitary. Its truncation to the zero mode sector, V , governs the low energy charged current interaction. Three sources contribute to the non-unitarity of V : the mixing of SU(2) doublet and singlet states, the truncation to the zero mode sector and the modification of the fermion gauge couplings. The last contribution is dominant, except for the third generation quarks. Defining $\Delta V_i = \sum_{j=1}^3 |V_{ij}|^2 - 1$, we find with the quark locations of eq. (3.24) $\Delta V_1 \approx 1 \cdot 10^{-5}$, $\Delta V_2 \approx 2 \cdot 10^{-7}$ and $\Delta V_3 \approx -3 \cdot 10^{-3}$. Again we have averaged over random sets of Yukawa couplings. The deviation from unitarity in the first generation is two orders of magnitude below the current experimental sensitivity [24]. Also the expectation of ΔV_3 is small compared to the few percent precision in the weak charge of the top quark to be reached at the LHC [27].

Charged current interactions are also mediated by the KK states of the W boson. These corrections are suppressed by $(g_1^2/\delta g_0^2)(M_W^2/M_{KK}^2) \lesssim 0.1$ and therefore sub-leading. The inequality becomes satisfied for the top quark which is localized closest towards the TeV-brane. Only for fermions localized very closely to the TeV-brane these corrections become important.

5 Flavor violation

The mechanisms we have discussed in the previous section in the context of charged current sector modify the neutral current interactions in a similar way. In analogy to eq. (4.30) we define the neutral current gauge couplings in the basis of mass eigenstates as

$$\mathcal{X}_{L,R}^{\psi(n)} = \mathcal{U}_{L,R}^{\psi} \mathcal{G}_{L,R}^{\psi(n)} \mathcal{U}_{L,R}^{\psi\dagger} \quad (5.31)$$

where the unitary matrices $\mathcal{U}_{L,R}^{\psi}$ diagonalize the full mass matrices, including the KK states of the fermion species $\psi = q_L, u_R, l_L$, etc. In ref. [19] we had not yet included KK mixing in the fermion sector. With $\mathcal{G}_{L,R}^{\psi n}$ we denote the coupling to the n th KK state of the Z boson as obtained from eq. (2.18). If the fermion families are localized at different positions in the extra dimension, the gauge couplings $\mathcal{G}_{L,R}^{\psi n}$ are non-universal in flavor space. These deviations in the fermion gauge couplings are too small to directly show up in flavor conserving processes at present accelerator experiments if the KK scale is about 10 TeV [11]. However, the transformation to the mass eigenstates (5.31) induces flavor violating couplings which lead to interesting new phenomena. These flavor violating interactions are an immediate consequence of our approach to the problem of fermion masses. The phenomenology is similar to what happens in models with family non-universal Z' bosons, so we can adopt the formalism described, for instance, in ref. [28] with the complication of the KK mixing of fermions. In the gauge boson sector we work with mass eigenstates from the very beginning. There is no mixing of the Z boson and its KK states, instead the zero mode has flavor violating couplings as well.

Flavor violation as induced by the couplings (5.31) is driven by the non-universality δg_{ij} in the couplings of different flavor states which are mixed by an angle θ_{ij}

$$\mathcal{X}_{ij} \approx \delta g_{ij} \sin \theta_{ij}. \quad (5.32)$$

Very different fermion locations on the one hand increase non-universality, on the other hand they lead to small mixing angles. Non-universality is larger for fermions localized towards the TeV-brane, $c < 1/2$, as is shown in fig. 3, and for the Z zero mode depends on the KK scale as $1/M_{KK}^2$. Note that, different from the standard model, the right-handed mixings become physically relevant. The left-handed mixings of quarks and leptons are naturally comparable to the observed CKM and neutrino mixing angles, respectively. This requirement fixed the left-handed fermion locations in section 3. The locations of right-handed fermions we have used to fit the fermion masses. For the locations (3.24) and (3.27) the right-handed mixings are then predicted as

$$\begin{aligned} \langle |\mathcal{U}_{R,1,2}^u| \rangle &= 0.017, & \langle |\mathcal{U}_{R,2,3}^u| \rangle &= 0.073, & \langle |\mathcal{U}_{R,1,3}^u| \rangle &= 0.0012, \\ \langle |\mathcal{U}_{R,1,2}^d| \rangle &= 0.23, & \langle |\mathcal{U}_{R,2,3}^d| \rangle &= 0.60, & \langle |\mathcal{U}_{R,1,3}^d| \rangle &= 0.23, \\ \langle |\mathcal{U}_{R,1,2}^e| \rangle &= 0.016, & \langle |\mathcal{U}_{R,2,3}^e| \rangle &= 0.057, & \langle |\mathcal{U}_{R,1,3}^e| \rangle &= 0.0016. \end{aligned} \quad (5.33)$$

Again we have averaged over random sets of Yukawa couplings. The right-handed down quarks mix strongly with each other, while the mixings of up quarks and leptons are found to be small. For the alternative other set of lepton locations (3.26) the right-handed mixings are similarly small.

In the following we are using a KK scale of 10 TeV which might be somewhat lowered by introducing brane-kinetic terms [13]. Note that the rates of flavor violating processes which we present below cannot be translated to that case by simple scaling with the appropriate powers of M_{KK} . Brane-kinetic terms modify the KK wave functions and thus possible non-universalities even if the KK scale is not changed.

5.1 Lepton flavor violation

Let us start the discussion of flavor violation with the lepton sector. In addition to the lepton locations (A) (3.26) and (B) (3.27) we are considering two modified settings

$$\begin{aligned} (A') : \quad c_{L1} &= 0.585, & c_{L2} &= 0.565, & c_{L3} &= 0.545, \\ (B') : \quad c_{L1} &= 0.520, & c_{L2} &= 0.200, & c_{L3} &= 0.150 \end{aligned} \quad (5.34)$$

to explore the dependence on the fermion locations. The right-handed positions are taken as in (A) and (B), respectively. In the set (A') we have shifted the left-handed leptons by a small amount to induce some non-universality in the left-handed sector. The changes in the lepton masses and mixings remain within the widths of the statistical distributions. In (B') we have reduced the Yukawa couplings by a common factor of twenty, which is compensated by moving the left-handed leptons closer towards the TeV-brane where non-universality is especially large. We also separated somewhat the left-handed muon and tau states. Using these locations we find the following lepton flavor violation Z couplings

	(A)	(A')	(B)	(B')	
$\langle \mathcal{X}_{L,1,2}^{e(0)} \rangle$	2.3×10^{-7}	6.5×10^{-7}	3.7×10^{-7}	8.4×10^{-5}	
$\langle \mathcal{X}_{L,2,3}^{e(0)} \rangle$	5.1×10^{-7}	1.2×10^{-6}	3.5×10^{-7}	3.2×10^{-5}	
$\langle \mathcal{X}_{L,1,3}^{e(0)} \rangle$	2.3×10^{-7}	8.5×10^{-7}	2.2×10^{-7}	4.9×10^{-5}	(5.35)
$\langle \mathcal{X}_{R,1,2}^{e(0)} \rangle$	7.1×10^{-9}	9.7×10^{-9}	1.0×10^{-8}	7.7×10^{-9}	
$\langle \mathcal{X}_{R,2,3}^{e(0)} \rangle$	7.6×10^{-7}	1.1×10^{-6}	2.3×10^{-6}	1.7×10^{-6}	
$\langle \mathcal{X}_{R,1,3}^{e(0)} \rangle$	7.3×10^{-9}	8.5×10^{-9}	6.5×10^{-8}	4.6×10^{-8}	

where we have included the zero modes and the first KK states of the leptons. Note that in the case (B') there is a large increase in the left-handed couplings compared to the cases with $c \gtrsim 1/2$. For (A) the flavor violating left-handed couplings completely

stem from fermion KK mixing since there is no non-universality among the left-handed states. In the other cases fermion KK mixing is a subleading effect.

The couplings (5.35) induce lepton flavor violating decays of the Z boson with a branching ratio $\text{Br}(Z \rightarrow l_i \bar{l}_j) \approx 0.29 \cdot (|\mathcal{X}_{L,i,j}^{e(0)}|^2 + |\mathcal{X}_{R,i,j}^{e(0)}|^2)$. The largest rate occurs in scenario (B') with $(Z \rightarrow e \bar{\mu}) \approx 2 \times 10^{-9}$ which is still three orders of magnitude below the experimental bound of 1.7×10^{-6} [24]. The other decay modes are even stronger suppressed. Thus there are no constraints on the model from lepton flavor violating Z decays.

The off-diagonal elements of \mathcal{X} also lead to flavor violating decays of charged leptons. Tree-level exchange of a Z boson and its KK states induces the processes $l_i \rightarrow 3l_j$ at a rate

	(A)	(A')	(B)	(B')	
$\text{Br}(\mu \rightarrow eee) :$	5.2×10^{-14}	4.7×10^{-13}	1.0×10^{-13}	5.3×10^{-9}	
$\text{Br}(\tau \rightarrow \mu\mu\mu) :$	1.1×10^{-13}	2.1×10^{-12}	7.3×10^{-13}	1.7×10^{-10}	(5.36)
$\text{Br}(\tau \rightarrow eee) :$	7.5×10^{-15}	3.2×10^{-13}	7.8×10^{-15}	3.9×10^{-10}	

Again we have averaged over random lepton Yukawa couplings in the four different scenarios. In the result (5.36) we have only included the contribution of the Z boson exchange. The amplitude contributed by its first KK state is already suppressed by a factor of about twenty. We have included the first KK level of lepton states which is important in scenario (A) where the left-handed lepton positions are degenerate. The branching ratios depend on the KK scale as $1/M_{KK}^4$. The experimental bound $\text{Br}(\mu \rightarrow eee) < 1.0 \times 10^{-12}$ [29] is satisfied for leptons localized towards the Planck-brane and the rates can come close to the experimental sensitivity. It would be very interesting if a slow muon facility could test this process at the precision of 10^{-16} [30]. The scenario (B'), where the leptons are moved towards the TeV-brane, leads to strong flavor violation. The rate could only be brought down to the experimental bound if the KK scale is increased by an order of magnitude to about 100 TeV. The expected rates for tau decays are in all cases well below the latest Belle results $\text{Br}(\tau \rightarrow \mu\mu\mu) < 3.8 \times 10^{-7}$ and $\text{Br}(\tau \rightarrow eee) < 2.7 \times 10^{-7}$ [31].

Severe experimental bounds have been put on μe -conversion in muonic atoms. The best exclusion limit $\text{Br}(\mu N \rightarrow eN) \equiv \Gamma(\mu N \rightarrow eN)/\Gamma(\mu N \rightarrow \nu_\mu N') < 6.1 \times 10^{-13}$ comes from the Sindrum-II collaboration [32]. We expect a branching ratio of

	(A)	(A')	(B)	(B')	
$\text{Br}(\mu N \rightarrow eN) :$	5.0×10^{-16}	5.2×10^{-15}	1.0×10^{-15}	5.8×10^{-11}	(5.37)

While the scenario (B') is again excluded for a KK scale of 10 TeV, leptons close towards the Planck-brane easily avoid the experimental limit. The forthcoming MECO experiment (E940 at BNL) plans to increase the sensitivity to about 5×10^{-17} [33]. It would completely cover the predicted interaction rates, even for leptons close to the Planck-brane.

At the one-loop level radiative decays that violate lepton flavor are induced at a rate

$$\begin{array}{cccc}
& (A) & (A') & (B) & (B') \\
\text{Br}(\mu \rightarrow e\gamma) : & 2.1 \times 10^{-16} & 1.9 \times 10^{-15} & 4.1 \times 10^{-16} & 2.1 \times 10^{-11} \\
\text{Br}(\tau \rightarrow \mu\gamma) : & 6.7 \times 10^{-16} & 9.5 \times 10^{-15} & 4.2 \times 10^{-15} & 6.6 \times 10^{-13} \\
\text{Br}(\tau \rightarrow e\gamma) : & 2.8 \times 10^{-17} & 1.3 \times 10^{-15} & 3.3 \times 10^{-17} & 1.5 \times 10^{-12}.
\end{array} \tag{5.38}$$

Except for the scenario (B') the expected branching ratios are far below the experimental bounds $\text{Br}(\mu \rightarrow e\gamma) < 1.2 \times 10^{-11}$, $\text{Br}(\tau \rightarrow \mu\gamma) < 1.1 \times 10^{-6}$ and $\text{Br}(\tau \rightarrow e\gamma) < 2.7 \times 10^{-6}$. The MEG collaboration at PSI plans to increase the sensitivity with respect to $\mu \rightarrow e\gamma$ to about $10^{-13} - 10^{-14}$ within the next years [34]. This will not be sufficient to test the model for leptons localized towards the Planck-brane. Note that our model leads to predictions quite different from supersymmetric models, where radiative decays occur at much larger rates than the decays of eqs. (5.36) and (5.37) [35]. In the warped SM radiative decays occur only at the loop level, while the latter are tree-level processes. In the scenario of Dirac neutrino masses radiative decays are also mediated by the KK states of the sterile neutrinos. If the SM neutrinos are confined to the TeV-brane, a too large branching ratio for $\mu \rightarrow e\gamma$ pushes the KK scale up to 25 TeV and thus imposes a stringent constraint on the model [15]. However, the rate for $\mu \rightarrow e\gamma$ is sensitive to the mixing between light and heavy neutrino states. With bulk neutrinos the mixing with heavy states is considerably reduced. In ref. [10] a branching ratio $\text{Br}(\mu \rightarrow e\gamma) \approx 10^{-15}$ was found for leptons localized towards the Planck-brane ($c > 1/2$). While this value is still well below the experimental sensitivity, it is larger than the contribution from gauge boson exchange (5.38). Other lepton flavor violating processes, such as muonium-antimuonium oscillations [36], do not lead to new constraints.

5.2 Meson mass splittings and CP violation

The flavor violating gauge couplings (5.31) also contribute to the mass splittings in neutral pseudo-scalar meson systems. The mass splitting between the flavor eigenstates of a meson, P^0 and \bar{P}^0 , is given by

$$\Delta m_P = \frac{\text{Re}\langle P^0 | -\mathcal{L}_{\text{FCNC}} | \bar{P}^0 \rangle}{m_K} \tag{5.39}$$

where $\mathcal{L}_{\text{FCNC}}$ contains the flavor violating couplings (5.31). Phases in the flavor violation couplings are constrained by the indirect CP-violation in the kaon system

$$\epsilon_K = \frac{\text{Im}\langle P^0 | -\mathcal{L}_{\text{FCNC}} | \bar{P}^0 \rangle}{2\sqrt{2}m_K\Delta m_H}. \tag{5.40}$$

These expressions can be evaluated using the vacuum insertion approximation (see for instance [37]). The largest contributions to (5.39) and (5.40) come from the

exchange of KK gluons [38]. Excited gluons have flavor non-universal couplings to fermions like the KK states of the weak gauge bosons and induce flavor violating couplings analogous to those of eq. (5.31). The gluon zero modes couple universally and therefore do not mediate flavor changing interactions.

For the quark locations of eq. (3.24) we obtain the following contributions of first level of KK gluons and the Z boson zero mode

	gluon	Z	exp.
$\Delta m_K :$	1.5×10^{-14}	1.2×10^{-17}	3.5×10^{-12}
$\Delta m_B :$	5.1×10^{-11}	3.0×10^{-14}	3.2×10^{-10}
$\Delta m_D :$	3.8×10^{-13}	5.2×10^{-15}	4.6×10^{-11}
$\epsilon_K :$	1.1×10^{-3}	1.1×10^{-6}	2.3×10^{-3}

(5.41)

The meson mass splittings are given in units of MeV. In eq. (5.41) we give results averaged over random sets of Yukawa couplings $2/3 < |l_{ij}| < 4/3$. As in the case of flat extra dimensions [38] the dominating contributions come from the exchange of KK gluons. KK states of the Z boson give only an about ten percent correction to the zero mode exchange. In the computation of the fermion mass eigenstates we have included the first level of KK states. Compared to using only fermion zero modes the gluon contribution to ϵ_K is enhanced by a factor of two. The meson mass splittings are less sensitive to whether or not KK fermions are included. The neglect of higher KK levels amounts to an uncertainty of order unity in our results.

The contributions to meson mass splittings we find are much smaller than the experimental values [24] listed in the last column of eq. (5.41). The KK gluon contribution to ϵ_K comes rather close to the observed value. About one third of the random sets of Yukawa couplings we tested gave $\epsilon_K > 2.3 \times 10^{-3}$, which is still acceptable. In ref. [17] KK contributions to $b \rightarrow s\gamma$ were considered. For the quark locations of eq. (3.24) a bound on the KK scale of about 5 TeV was found. From the the process $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ the limit on the Z coupling $|\mathcal{X}_{L,1,2}^{d(0)}| < 5.1 \times 10^{-6}$ was obtained in ref. [18]. We find a considerably lower value of $\langle |\mathcal{X}_{L,1,2}^{d(0)}| \rangle = 2.2 \times 10^{-7}$. Our approach to fermion masses and mixings is therefore nicely compatible with experimental constraints on flavor violation.

We stress that our results are very different from models where the fermions mass pattern is explained by flavor-dependent fermion locations in one or more universal *flat* extra dimensions. There flavor violation, in particular kaon mixing, leads to very restrictive bounds on the KK scale, $M_{KK} \gtrsim 10^3$ TeV [38], disfavoring these models as a solution to the the gauge hierarchy problem. Recently it was shown that models constructed from intersecting D-branes suffer from a similar problem [39]. The crucial difference lies in the wave functions of the KK gauge bosons. With a warped extra dimension the gauge boson wave functions are almost constant away from the TeV-brane. Therefore the gauge couplings of KK gauge bosons are nearly universal for fermions localized somewhat towards the Planck-brane ($c > 1/2$) as

we show in fig. 3. Flavor violation is thus automatically suppressed for the light fermion species which are expected to reside closely towards the Planck-brane to explain their small masses.

6 Non-renormalizable operators and rare processes

In models where the weak scale is identified with the fundamental scale of gravity, the low cut-off scale dramatically amplifies the impact of non-renormalizable operators in weak scale interactions. As a consequence, large rates for rare processes, such as flavor violation and proton decay, are a challenge for model building. With bulk fermions localized towards the Planck-brane the corresponding suppression scales can be significantly enhanced without relying on ad-hoc symmetries [5, 6]. However, there are limits because the SM fermions need to have sufficient overlap with the Higgs field at the TeV-brane to acquire their observed masses, as discussed in section 3. We consider the following generic four-fermion operators which are relevant for flavor violation as well as for proton decay

$$\int d^4x \int dy \sqrt{-g} \frac{1}{M_5^3} \bar{\Psi}_i \Psi_j \bar{\Psi}_k \Psi_l \equiv \int d^4x \frac{1}{Q^2} \bar{\Psi}_i^{(0)} \Psi_j^{(0)} \bar{\Psi}_k^{(0)} \Psi_l^{(0)}. \quad (6.42)$$

Integrating over the extra dimension, the effective 4D suppression scales Q associated with these operators depend on where the relevant fermion states are localized in the extra dimension.

Let us focus on some examples. The lepton flavor violating decay $\mu \rightarrow eee$ is induced by the operator μeee at a rate $\Gamma \sim m_\mu^5/Q^4$. The experimental constraint on the corresponding branching ratio translates into $Q > 5 \times 10^5$ GeV. For the lepton locations considered in the previous section we obtain the suppression scales

$$\begin{array}{cccc} \text{(A)} & \text{(A')} & \text{(B)} & \text{(B')} \\ Q(\mu eee)[\text{GeV}] : & 4.8 \times 10^6 & 1.1 \times 10^7 & 7.9 \times 10^7 & 2.4 \times 10^5, \end{array} \quad (6.43)$$

where we have taken all the leptons to be left-handed. Except for the case (B') where the leptons are localized towards the TeV-brane, the operator μeee is safely suppressed. The same holds for similar lepton flavor violating operators such as $\tau \mu \mu \mu$ and τeee . The operator $\mu e q_1 q_1$ contributing to muon electron conversion is constrained by $Q > 1 \times 10^5$ GeV while we are finding

$$\begin{array}{cccc} \text{(A)} & \text{(A')} & \text{(B)} & \text{(B')} \\ Q(\mu e q_1 q_1)[\text{GeV}] : & 4.4 \times 10^7 & 3.5 \times 10^7 & 1.1 \times 10^8 & 6.6 \times 10^6. \end{array} \quad (6.44)$$

These suppression scales even exceed $Q > 1 \times 10^6$ GeV, a bound which could be set by the upcoming MECO experiment. Thus non-renormalizable operators are not expected to induce lepton flavor violation at an observable rate, unless the leptons would be localized closely towards the TeV-brane.

Constraints on the $K - \bar{K}$ mass splitting require the dimension-six operator $(ds)^2$ to be suppressed by $Q > 5 \times 10^6$ GeV. If it contributes to CP violation, an even stronger suppression of $Q > 5 \times 10^7$ GeV is required. Using the quark locations of eq. (3.24), we obtain $Q > 7.2 \times 10^7$ GeV for left-handed and $Q > 1.2 \times 10^8$ GeV for right-handed states. For the operator $(db)^2$ we find $Q > 3.3 \times 10^6$ GeV, which is above the corresponding experimental bound of 2×10^6 GeV. Other flavor violating operators like $(cu)^2$ are also within their experimental bounds. The quark locations of eq. (3.24) which we obtained from the fermion mass pattern automatically lead to the required suppression of flavor violating non-renormalizable operators. This conforms the conclusions reached in refs. [5, 6].

Dimension-six operators contributing to proton decay are highly constrained by experimental searches. For instance the operator $q_1 q_1 q_2 l_3$ has to be suppressed by $Q > 10^{15}$ GeV [40]. Taking again the quark locations of eq. (3.24) and the lepton locations of eq. (3.26) we obtain $Q > 5.6 \times 10^7$ GeV. So some small coupling of order 10^{-14} should multiply the non-renormalizable operator to be consistent with experimental limits on proton decay. This might point to an additional symmetry, such as baryon or lepton number. A tiny coupling might also originate from non-perturbative effects of gravity, especially if there is an extra dimension somewhat larger than M_{PL} [41]. If the proton is stabilized by imposing lepton number (or lepton parity in the case of Majorana neutrino masses), baryon number violating processes like neutron-antineutron oscillations could still occur [42]. With the quark locations of eq. (3.24) the rate is, however, far below the experimental bound.

In ref. [6] somewhat different quark locations were used in order to maximally suppress proton decay by non-renormalizable interactions. Similar conclusions were reached, even though the suppression scale Q could be enhanced by an order of magnitude. Non-renormalizable interactions can be further suppressed if the fermions are shifted closer towards the Planck-brane by allowing for larger 5D Yukawa couplings. A smaller AdS curvature compared to the fundamental Planck mass has the same effect. However, using these means to stabilize the proton would necessarily introduce new hierarchies in the model parameters.

7 Conclusions

In this paper we have studied aspects of flavor physics in the warped SM. Gauge bosons and fermions are bulk fields, while the Higgs field is confined to the TeV-brane. We have shown in detail how the fermion mass hierarchies and mixings can naturally be explained in a geometrical way, without relying on hierarchical Yukawa couplings. The observed fermion masses and mixings fix the relative positions of the fermion fields in the extra dimension. Large mixings are attributed to similar locations, and masses become small for fermions localized towards the Planck-brane. Small neutrino masses can arise from a coupling to right-handed neutrinos in the bulk or from dimension-five interactions.

Mixings between different KK levels induces deviations from the SM. Electroweak fits require the KK scale to be at least 10 TeV. KK mixings in the quark and weak gauge boson sectors lead to a non-unitary CKM matrix. However, deviations from unitarity are safely within the experimental bounds.

Flavor violation by (KK) gauge boson exchange is an immediate consequence of our approach to the fermion mass problem. After transformation to fermion mass eigenstates, non-universal gauge couplings generate flavor changing neutral currents. Fixing the fermion locations by the fermion masses and mixings, we can predict the rates of flavor violations processes. Since the light fermion flavors are localized towards the Planck-brane, where non-universality is small, flavor violation is within experimental bounds even for a KK scale of 10 TeV. Some processes, such as muon-electron conversion, are in the reach of next generation experiments and can provide valuable hints to the higher dimensional theory. This result is quite different from models with flat extra dimensions, where constraints on kaon mixing require the KK scale to be in the 10^3 TeV range. Flavor violation in radiative decays is suppressed which is an important distinction to supersymmetric models.

Bulk fermions also help to reduce the impact of non-renormalizable operators. We have demonstrated that their contribution to flavor violating processes is naturally below the experimental sensitivity, even for order unity couplings. Dimension-six operators leading to proton decay, however, cannot be sufficiently suppressed. The required tiny couplings might point to some additional symmetry.

Thus, the warped SM allows to generate the fermion mass pattern from flavor-dependent locations without giving up the solution to the gauge hierarchy problem or inducing unacceptable rates for flavor violating processes.

Acknowledgements

The author thanks David E. Costa and Qaisar Shafi for valuable discussions.

References

- [1] L. Randall and R. Sundrum, *Phys. Rev. Lett.* **83** (1999) 3370.
- [2] M. Gogberashvili, hep-ph/9812296.
- [3] S. Chang, J. Hisano, H. Nakano, N. Okada and Yamaguchi, *Phys. Rev.* **D62** (2000) 084025 [hep-ph/9912498].
- [4] S.J. Huber and Q. Shafi, *Phys. Rev.* **D63** (2001) 045010 [hep-ph/0005286].
- [5] T. Gherghetta and A. Pomarol, *Nucl. Phys.* **B586** (2000) 141 [hep-ph/0003129].
- [6] S.J. Huber and Q. Shafi, *Phys. Lett.* **B498** (2001) 256 [hep-ph/0010196].

- [7] N. Arkani-Hamed and M. Schmaltz, *Phys. Rev. D* **61** (2000) 033005 [hep-ph/9903417]; G.R. Dvali and M.A. Shifman, *Phys. Lett.* **B475** (2000) 295 [hep-ph/0001072].
- [8] S.J. Huber and Q. Shafi *Phys. Lett.* **B544** (2002) 295 [hep-ph/0205327].
- [9] Y. Grossman and M. Neubert, *Phys. Lett.* **B474** (2000) 361 [hep-ph/9912408].
- [10] S.J. Huber and Q. Shafi, *Phys. Lett.* **B512** (2001) 365 [hep-ph/0104293].
- [11] S.J. Huber, C.-A. Lee and Q. Shafi, *Phys. Lett.* **B531** (2002) 112 [hep-ph/0111465].
- [12] J.L. Hewett, F.J. Petriello and T.G. Rizzo, *JHEP* **0209** (2002) 030 [hep-ph/0203091].
- [13] H. Davoudiasl, J.L. Hewett and T.G. Rizzo, hep-ph/0212279; M. Carena, E. Ponton, T.M.P. Tait and C.E.M. Wagner, hep-ph/0212307.
- [14] C. Csaki, J. Erlich and J. Terning *Phys. Rev.* **D66** (2002) 064021 [hep-ph/0203034].
- [15] R. Kitano, *Phys. Lett.* **B481** (2000) 39 [hep-ph/0002279].
- [16] F. del Aguila and J. Santiago, *Phys. Lett.* **B493** (2000) 175 [hep-ph/0008143].
- [17] C.S. Kim, J.D. Kim and J. Song, *Phys. Rev.* **D67** (2003) 015001 [hep-ph/0204002].
- [18] G. Burdman, *Phys. Rev.* **D66** (2002) 076003 [hep-ph/0205329].
- [19] S.J. Huber, to appear in the Proceedings of SUSY 2002, hep-ph/0211056.
- [20] H. Davoudiasl, J.L. Hewett and T.G. Rizzo, *Phys. Lett.* **B473** (2000) 43; A. Pomarol, *Phys. Lett.* **B486** (2000) 153.
- [21] H.B. Nielsen and Y. Takanishi, *Phys. Lett.* **B543** (2002) 249 [hep-ph/0205180]; C.D. Froggatt, H.B. Nielsen and Y. Takanishi, *Nucl. Phys.* **B631** (2002) 285 [hep-ph/0201152].
- [22] H. Fusaoka and Y. Koide, *Phys. Rev.* **D57** (1998) 3986-4001 [hep-ph/9712201].
- [23] H. Fritzsch and Z. Xing, *Prog. Part. Nucl. Phys.* **45** (2000) 1 [hep-ph/9912358].
- [24] K. Hagiwara *et al.* [Particle Data Group], *Phys. Rev.* **D66** (2002) 010001.
- [25] C. Jarlskog, *Phys. Rev. Lett.* **55** (1985) 1039.
- [26] P.Q. Hung and M. Seco, hep-ph/0111013.

- [27] M. Beneke *et al.*, hep-ph/0003033.
- [28] P. Langacker and M. Plümacher, *Phys. Rev.* **D62** (2000) 013006 [hep-ph/0001204].
- [29] U. Bellgardt *et al.* [SINDRUM Collaboration] *Nucl. Phys.* **B299** (1988) 1.
- [30] J. Aysto *et al.*, hep-ph/0109217.
- [31] Y. Yusa *et al.*, [BELLE Collaboration], hep-ex/0211017.
- [32] P. Wintz, in *Proceedings of the first International Symposium on Lepton and Baryon Number Violation*, p. 534 (1998).
- [33] M. Bachmann *et al.*, [MECO Collaboration], Research Proposal E940 for an experiment at BNL (1997).
- [34] L.M. Barkov, *et al.*, [MEGA Collaboration], Research Proposal for an experiment at PSI (1999).
- [35] J. Hisano and D. Nomura, *Phys. Rev.* **D59** (1999) 116005 [hep-ph/9810479]; J.A. Casas and A. Ibarra, *Nucl. Phys.* **B618** (2001) 171-204 [hep-ph/0103065].
- [36] L. Willmann *et al.*, *Phys. Rev. Lett.* **82** (1999) 49 [hep-ex/9807011].
- [37] F. Gabbiani, E. Gabrielli, A. Masiero and L. Silvestrini *Nucl. Phys.* **B477** (1996) 321 [hep-ph/9604387]; T. Moroi, *JHEP* **0003** (2000) 019 [hep-ph/0002208].
- [38] A. Delgado, A. Pomarol and M. Quiros *JHEP* **0001** (2000) 030 [hep-ph/9911252].
- [39] S. Abel, M. Masip and J. Santiago, hep-ph/0303087.
- [40] K. Benakli and S. Davidson, *Phys. Rev.* **D60** (1999) 025004 [hep-ph/9810280].
- [41] A.B. Kobakhidze, *Phys. Lett.* **B514** (2001) 131-138 [hep-ph/0102323]; R. Kallosh, A.D. Linde, D.A. Linde and L. Susskind, *Phys. Rev.* **D52** (1995) 912.
- [42] C.E. Carlson and C.D. Carone, *Phys. Lett.* **B512** (2001) 121 [hep-ph/0103180].