

Sussex Research

The moment index of minima (II)

D. J. Daley, Charles Goldie

Publication date

01-01-2006

Licence

This work is made available under the **Copyright not evaluated** licence and should only be used in accordance with that licence. For more information on the specific terms, consult the repository record for this item.

Citation for this work (American Psychological Association 7th edition)

Daley, D. J., & Goldie, C. (2006). *The moment index of minima (II)* (Version 1). University of Sussex.
<https://hdl.handle.net/10779/uos.23312708.v1>

Published in

Statistics and Probability Letters

Link to external publisher version

<https://doi.org/10.1016/j.spl.2005.10.013>

Copyright and reuse:

This work was downloaded from Sussex Research Open (SRO). This document is made available in line with publisher policy and may differ from the published version. Please cite the published version where possible. Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners unless otherwise stated. For more information on this work, SRO or to report an issue, you can contact the repository administrators at sro@sussex.ac.uk. Discover more of the University's research at <https://sussex.figshare.com/>

The moment index of minima (II)

Article (Unspecified)

Citation:

Daley, D.J. and Goldie, Charles M. (2006) The moment index of minima (II). Statistics and Probability Letters, 76. pp. 831-837. ISSN 0167-7152

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/1647/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

THE MOMENT INDEX OF MINIMA (II)

D.J. Daley*

and

Charles M. Goldie

The Australian National University

The University of Sussex

Abstract

The moment index $\kappa(X) = \sup\{k: E(X^k) < \infty\}$ of a nonnegative random variable X has the property that $\kappa(\min(X, Y)) \geq \kappa(X) + \kappa(Y)$ for independent r.v.s X and Y . We characterize conditions under which equality holds for a given r.v. X and every independent nonnegative r.v. Y , and discuss extensions to related r.v.s and their distributions.

Key words and phrases: exponential index, moment index, regular variation.

1. Introduction

In Daley (2001) to which this note is a sequel, the *moment index* $\kappa(X)$ of a nonnegative random variable (r.v.) X is defined by

$$\kappa(X) = \sup\{k \geq 0 : E(X^k) < \infty\}. \quad (1)$$

It was shown that for independent nonnegative r.v.s X and Y each with a finite moment index,

$$\kappa(\min(X, Y)) \geq \kappa(X) + \kappa(Y), \quad (2)$$

that equality holds when the tail of the d.f. of either X or Y is regularly varying, and an example in which X and Y have discrete supports that are ‘increasingly sparse’ and ‘well interspersed’ demonstrated that the inequality at (2) can be strict.

The main purpose of this paper is to prove the theorem below; it characterizes independent nonnegative r.v.s X and Y for which equality holds in (2). We precede its proof in Section 2 with further discussion. In Section 3 we note companion results for the exponential index of a r.v., and Section 4 looks at questions surrounding the finiteness or otherwise of $E(X^{\kappa(X)})$.

*Centre for Mathematics and its Applications (SMS), Australian National University, Canberra, ACT 0200, Australia. Email: daryl@maths.anu.edu.au

Definition 1. (a) \mathcal{M}^α denotes the class of all nonnegative r.v.s X with moment index $\kappa(X) = \alpha$ for which, for every nonnegative r.v. Y independent of X ,

$$\kappa(\min(X, Y)) = \kappa(X) + \kappa(Y).$$

(b) For $\alpha < \infty$, $\mathcal{M}^{\alpha+} \subseteq \mathcal{M}^\alpha$ consists of those r.v.s $X \in \mathcal{M}^\alpha$ for which $E(X^\alpha) < \infty$, and $\mathcal{M}^{\alpha-} = \mathcal{M}^\alpha \setminus \mathcal{M}^{\alpha+}$.

Note that $\mathcal{M}^{0+} = \mathcal{M}^0$. For intermediate values $0 < \alpha < \infty$ the inclusion (b) is proper.

Theorem 2. $X \in \mathcal{M}^\alpha$ if and only if the tail \bar{F} of its d.f. F satisfies

$$\lim_{x \rightarrow \infty} [-\log \bar{F}(x)] / \log x = \alpha = \kappa(X). \quad (3)$$

We remark that the class \mathcal{M}^α of d.f.s for $\alpha < \infty$, which by Daley (2001) includes d.f.s with regularly varying tails of index α , is indeed larger than the latter family. This follows essentially as in Proposition 2.2.8 of Bingham, Goldie and Teugels (1989) (hereafter, [BGT]), where there is an example of a monotone function whose lower and upper orders coincide (see equations (5)–(6) below) but for which the representation theorem [BGT 2.2.7] is not of the form of the corresponding theorem [BGT Theorem 1.3.1] for regularly varying functions.

We owe to a referee the idea that Theorem 2 might have implications for statistical estimation of the moment index. Estimation of tail properties of a random variable is a ‘difficult’ statistical question, being the estimation of a rare event. The definition of moment index does not suggest any estimating procedure: can one maybe estimate α by (3) for the subclass \mathcal{M}^α ? Assuming the problem is cast in the setting of a sequence of i.i.d. r.v.s, one immediately runs up against the obstacle that the Strong Law of Large Numbers holds only when the first moment is finite. Now considering the class of Paréto distributions on $(1, \infty)$, namely those with $\Pr\{X > x\} = x^{-\alpha}$ for $x \geq 1$, we have

$$\Pr\{X > x\} = \Pr\{X > y\} \Pr\{X > x/y\} \quad (1 < y < x),$$

and this suggests that observations should be aggregated from indicators of tail events such as $\{X \geq x\}$ rather than using the values of individual realizations x_i , because we can then plot

$$\frac{\sum_{i=1}^n \mathbf{1}\{x_i > x\}}{n \ln x}$$

for a range of values $x > 1$. This leads us to something like the Hill estimator for a tail index. In other words, we are led to the further subclass of distributions with regularly varying tails, for which there is extensive experience (and knowledge of the problems) of estimating α , as a tail index. A recent contribution to such estimation problems is Goncalves and Riedi (2005).

2. Discussion and proof of Theorem 2

The identification of α in (3) with $\kappa(X)$ is a matter of definition. Also, it is known (though perhaps not well known; see Baltrūnas, Daley and Klüppelberg, 2004) that

$$\liminf_{x \rightarrow \infty} \frac{-\log \bar{F}(x)}{\log x} = \kappa(X), \quad (4)$$

so we give it as Lemma BDK below and indicate its proof; note that for a positive function f , its *lower order* $\mu(f)$ is just

$$\mu(f) = \liminf_{x \rightarrow \infty} [\log f(x)] / \log x, \quad (5)$$

while the companion *upper order* $\nu(f)$ say is

$$\nu(f) = \limsup_{x \rightarrow \infty} [\log f(x)] / \log x \quad (6)$$

[BGT Section 2.2.2]. Consequently, what is new in Theorem 2 is the identification of \mathcal{M}^α with tails of d.f.s \bar{F} for which $\mu(1/\bar{F}) = \nu(1/\bar{F})$.

Now the tail of the d.f. of $\min(X, Y)$ is just the function $\bar{F}\bar{G}$, where G is the d.f. of Y , and for any real-valued functions f and g with finite limits infima, $\liminf_{t \rightarrow \infty} [f(t) + g(t)] \geq \liminf_{t \rightarrow \infty} f(t) + \liminf_{t \rightarrow \infty} g(t)$, where equality holds for any given f and all g if and only if

$\lim_{t \rightarrow \infty} f(t)$ exists. In exploiting this property to demonstrate that inequality at (2) may hold for given \bar{F} for which $\nu(1/\bar{F}) > \mu(1/\bar{F}) = \alpha$, we need to ensure that the function \bar{G} we construct with given lower moment order smaller than its upper order is indeed a distribution function.

Nevertheless, there are pairs of independent r.v.s X and Y for which the limits infima and suprema are different but for which equality holds at (2). One such pair is as in Lemma 3 (the proof is at the end of this section), which shows that for strict inequality to hold in (2), the regions where the ratios at (4) are close to their limits infima must not overlap but rather be well interspersed as in the example in Daley (2001). Indeed, in an unpublished student essay at the ANU, Tu Anh Nguyen has given an example for which the limits infima are finite (and hence so too is the right-hand side of (2)) but the left-hand side of (2) is infinite because the limits suprema are infinite, the ratios in (4) being ‘large’ in different regions.

Lemma 3. *Let the nonnegative r.v. X have finite first moment, and let the nonnegative r.v. Y , independent of X , have probability density function proportional to the tail $\bar{F}(\cdot)$ of the d.f. of X . Then $\kappa(Y) = \kappa(X) - 1$ and $\kappa(\min(X, Y)) = 2\kappa(X) - 1$.*

Lemma BDK. *For a nonnegative r.v. X , the lower order of the reciprocal $1/\bar{F}(x)$ of the tail of its d.f. equals its moment index, i.e.*

$$\mu(1/\bar{F}) \equiv \liminf_{x \rightarrow \infty} \frac{-\log \bar{F}(x)}{\log x} = \kappa(X) \equiv \sup\{k \geq 0 : E(X^k) < \infty\}. \quad (7)$$

PROOF. We first show that $\kappa(X) \geq \mu(1/\bar{F})$. If $\mu(1/\bar{F}) = 0$ there is nothing to prove. When $0 < \mu(1/\bar{F}) < \infty$ let $\lambda = \mu(1/\bar{F})$ and observe that for arbitrary $\varepsilon > 0$, $-\log \bar{F}(x) \geq (\lambda - \varepsilon) \log x = \log x^{\lambda - \varepsilon}$ for all $x \geq$ some $x_0 = x_0(\varepsilon)$. Then for such x , $\bar{F}(x) \leq 1/x^{\lambda - \varepsilon}$ and therefore

$$\int_{x_0}^{\infty} x^{\lambda - 1 - 2\varepsilon} \bar{F}(x) dx \leq \int_{x_0}^{\infty} \frac{dx}{x^{1 + \varepsilon}} < \infty. \quad (8)$$

Since

$$E(X^k) = \int_0^{\infty} kx^{k-1} \bar{F}(x) dx \quad \text{when } k > 0,$$

(8) implies that $\kappa(X) \geq \lambda - 2\varepsilon$, and as $\varepsilon > 0$ is otherwise arbitrary, we conclude $\kappa(X) \geq \mu(1/\bar{F})$.

In the case $\mu(1/\bar{F}) = \infty$ the same argument works for all $\lambda > 0$, so we conclude $\kappa(X) = \infty$ as wanted.

For the converse assertion, that $\mu(1/\bar{F}) \geq \kappa(X)$, the argument is similarly structured. If $\kappa(X) = 0$ there is nothing to prove. If $0 < \kappa(X) < \infty$ we write $\lambda = \kappa(X)$ and note that for $0 < \varepsilon < \lambda$,

$$\infty > \int_0^\infty x^{\lambda-\varepsilon} dF(x) = E(X^{\lambda-\varepsilon}) \geq x^{\lambda-\varepsilon} \bar{F}(x),$$

so $(\lambda - \varepsilon) \log x + \log \bar{F}(x) \leq \log E(X^{\lambda-\varepsilon})$ and

$$\frac{-\log \bar{F}(x)}{\log x} \geq \lambda - \varepsilon - \frac{\log E(X^{\lambda-\varepsilon})}{\log x} \rightarrow \lambda - \varepsilon \quad (x \rightarrow \infty), \quad (9)$$

and hence $\mu(1/\bar{F}) \geq \lambda = \kappa(X)$. If $\kappa(X) = \infty$ this argument works for all $\lambda > 0$, hence $\mu(1/\bar{F}) = \infty$ as wanted. ■

Proof of Theorem 2. The proof of (1.2) in Daley (2001) is probabilistic. Here we apply Lemma BDK to the r.v. $\min(X, Y)$, whose d.f. has the tail $\bar{F} \bar{G}$, in writing

$$\kappa(\min(X, Y)) = \liminf_{x \rightarrow \infty} \frac{-\log \bar{F}(x) - \log \bar{G}(x)}{\log x} \quad (10)$$

$$\geq \liminf_{x \rightarrow \infty} \frac{-\log \bar{F}(x)}{\log x} + \liminf_{x \rightarrow \infty} \frac{-\log \bar{G}(x)}{\log x} = \kappa(X) + \kappa(Y), \quad (11)$$

with equality holding when (3) holds.

For the converse, suppose X has moment index $\alpha < \infty$ but that $1/\bar{F}$ has upper order exceeding α , i.e.

$$\alpha = \liminf_{x \rightarrow \infty} \frac{-\log \bar{F}(x)}{\log x} < \limsup_{x \rightarrow \infty} \frac{-\log \bar{F}(x)}{\log x} \leq \infty.$$

Then there is a sequence $x_n \rightarrow \infty$ and constant $\varepsilon > 0$ such that

$$-\log \bar{F}(x_n) \geq (\alpha + 2\varepsilon) \log x_n \quad \text{for all } n, \quad (12)$$

and moreover we may choose the initial member x_0 of the sequence so large that

$$-\log \bar{F}(x) \geq (\alpha - \varepsilon) \log x \quad \text{for all } x \geq x_0. \quad (13)$$

Further, we can assume without loss of generality, just by taking a subsequence of x_n if need be, that

$$\log x_{n+1} > \log x_n'' \equiv \frac{\alpha + 2\varepsilon}{\alpha + \varepsilon} \log x_n. \quad (14)$$

The idea now is to choose $\beta > 2(\alpha + \varepsilon)$ and construct Y , or rather the tail \bar{G} of its d.f., with $\kappa(Y) = \beta$ and the additional properties that

$$-\log \bar{G}(x) \geq (\beta + 2\varepsilon) \log x \quad \text{whenever} \quad -\log \bar{F}(x) \leq (\alpha + \varepsilon) \log x, \quad (15)$$

and that on some sequence $x'_n \rightarrow \infty$,

$$-\log \bar{G}(x'_n) = \beta \log x'_n \quad \text{for all } n. \quad (16)$$

We will ensure that $-\log \bar{G}$ has lower order β , and hence $\kappa(Y) = \beta$, by insisting that

$$-\log \bar{G}(x) \geq \beta \log x \quad \text{for all } x \geq x_0. \quad (17)$$

If we can do all this then

$$-\log \bar{F}(x) - \log \bar{G}(x) \geq (\alpha + \beta + \varepsilon) \log x \quad \text{for all } x \geq x_0, \quad (18)$$

as this is so by (15) and (13) when $-\log \bar{F}(x) \leq (\alpha + \varepsilon) \log x$, and by (17) when $-\log \bar{F}(x) > (\alpha + \varepsilon) \log x$. By (18), (10) exceeds $\alpha + \beta = \kappa(X) + \kappa(Y)$, so $X \notin \mathcal{M}^\alpha$ as is to be proved.

There remains the construction of \bar{G} , i.e. of a non-decreasing function $-\log \bar{G}$ satisfying (15), (16) and (17). To the right of any x_n , because $-\log \bar{F}(x)$ is non-decreasing, we have

$$-\log \bar{F}(x) \geq -\log \bar{F}(x_n) \geq (\alpha + 2\varepsilon) \log x_n > (\alpha + \varepsilon) \log x \quad \text{for all } x \in [x_n, x_n''),$$

where

$$\log x_n'' := \frac{\alpha + 2\varepsilon}{\alpha + \varepsilon} \log x_n.$$

Within this interval, \bar{F} does not satisfy the condition that activates (15). Let us define \bar{G} to have the constant value

$$-\log \bar{G}(x) := (\beta + 2\varepsilon) \log x_n \quad \text{for } x \in [x_n, x_n'], \quad (19)$$

where

$$\log x_n' := \frac{\beta + 2\varepsilon}{\beta} \log x_n.$$

Our having fixed $\beta > 2(\alpha + \varepsilon)$ ensures that

$$\frac{\beta + 2\varepsilon}{\beta} = 1 + \frac{2\varepsilon}{\beta} < \frac{\alpha + 2\varepsilon}{\alpha + \varepsilon},$$

and so $x_n' < x_n''$ and thus (19) does not conflict with (15). Note that (19) implies (16).

For the rest of the definition of \bar{G} , just put

$$-\log \bar{G}(x) := (\beta + 2\varepsilon) \log x \quad \text{for all } x \in (x_0, \infty) \setminus \left(\bigcup_{n=1}^{\infty} (x_n, x_n'] \right). \quad (20)$$

Then $-\log \bar{G}$ is non-decreasing, and (20) and (19) together ensure (17). The construction is complete. ■

Proof of Lemma 3. That $\kappa(Y) = \kappa(X) - 1$ follows from the definition (1) as noted in Daley (2001). For the rest, let $\alpha = \kappa(X)$, and let $\{x_n\}$ be an increasing sequence for which $x_n \rightarrow \infty$ as $n \rightarrow \infty$ and

$$\lim_{n \rightarrow \infty} \frac{-\log \bar{F}(x_n)}{\log x_n} = \alpha. \quad (21)$$

For any fixed $c > 1$,

$$\frac{-\log \bar{F}(x_n/c)]}{\log(x_n/c)} \leq \frac{-\log \bar{F}(x_n)}{\log x_n - \log c} \rightarrow \alpha \quad (n \rightarrow \infty), \quad (22)$$

where we have used monotonicity of \bar{F} and the limit from (21). From the limit infimum property of α , it then follows that the left-hand side of (22) must have α as its limit for $n \rightarrow \infty$.

Next consider

$$\frac{-\log \left(\int_{x_n/c}^{\infty} \bar{F}(u) du / E(X) \right)}{\log(x_n/c)}, \quad (23)$$

whose limit infimum is bounded below by $\alpha - 1$ because this equals the moment index of Y , while the numerator with the factor $E(X)$ omitted (without affecting the limit property) is bounded above by

$$-\log \left[\int_{x_n/c}^{x_n} \bar{F}(u) du \right] \leq -\log [x_n(1 - c^{-1})\bar{F}(x_n)].$$

It is readily checked that this quantity, when divided by $\log x_n$, has limit as $n \rightarrow \infty$ equal to $-1 + \alpha$.

Then (23) has a limit as $n \rightarrow \infty$, and it equals $\alpha - 1$, which result can be combined with the limit of (22) to give

$$\frac{-\log \bar{F}(x_n/c) - \log \int_{x_n/c}^{\infty} \bar{F}(u) du}{\log(x_n/c)} \rightarrow \alpha + (\alpha - 1) = 2\alpha - 1. \quad \blacksquare$$

3. The exponential index of a nonnegative r.v.

We hope that the following discussion will facilitate greater use of moment generating functions as a handy technical device.

Definition 4. The exponential index $\varepsilon(X)$ of a real-valued r.v. X for which $\bar{F}(x) = \Pr\{X \geq x\}$, is defined by

$$\varepsilon(X) = \sup \{t: E(e^{tX}) < \infty\}. \quad (24)$$

We could now develop analogues of Lemma BDK and Theorem 2 for $\varepsilon(\cdot)$. However there is no point in reproducing the earlier construction and arguments because *the moment index of a positive r.v. X is related to the exponential index of the real r.v. $Y = \log X$ by $\varepsilon(Y) = \kappa(X)$* . This follows immediately, given a r.v. Y , from writing $Z = e^Y$ in

$$\varepsilon(Y) = \sup \{t: E(e^{tY}) < \infty\} = \sup \{t: E(Z^t) < \infty\} = \kappa(Z) = \kappa(e^Y). \quad (25)$$

The analogues to which we have alluded can be stated as follows without need of further proof.

Lemma 5.

$$\varepsilon(X) = \liminf_{x \rightarrow \infty} \frac{-\log \bar{F}(x)}{x}. \quad (26)$$

Theorem 6. *For independent real-valued r.v.s X and Y ,*

- (i) $\varepsilon(X + Y) = \min(\varepsilon(X), \varepsilon(Y))$;
- (ii) $\varepsilon(\max(X, Y)) = \min(\varepsilon(X), \varepsilon(Y))$; and
- (iii) $\varepsilon(\min(X, Y)) \geq \varepsilon(X) + \varepsilon(Y)$, where equality holds for all r.v.s Y if and only if

$$\limsup_{x \rightarrow \infty} \frac{-\log \bar{F}(x)}{x} = \varepsilon(X).$$

In this theorem, parts (i) and (ii) are included for the sake of completeness (cf. Daley, 2001), while (iii) follows from Theorem 2.

4. Discussion

For independent r.v.s $X \in \mathcal{M}^{\alpha+}$ and $Y \in \mathcal{M}^{\beta+}$ it follows from $E([\min(X, Y)]^{\alpha+\beta}) \leq E(X^\alpha Y^\beta) = E(X^\alpha)E(Y^\beta) < \infty$ that $\min(X, Y) \in \mathcal{M}^{(\alpha+\beta)+}$. Indeed, we have the following.

Lemma 7. *For independent r.v.s $X_j \in \mathcal{M}^{\alpha_j}$ and for nonnegative α_j ($j = 1, \dots, k$), $Z \equiv \min(X_1, \dots, X_k) \in \mathcal{M}^\alpha$ where $\alpha = \alpha_1 + \dots + \alpha_k$.*

If, for each j , α_j is finite and $X \in \mathcal{M}^{\alpha_j+}$, then $Z \in \mathcal{M}^{\alpha+}$.

On the other hand, there exist r.v.s X and $Y \in \mathcal{M}^{\alpha-}$ but $\min(X, Y) \in \mathcal{M}^{\alpha+}$. For example it is enough that they have d.f.s given by $\bar{F}(x) = \bar{G}(x) = 1/[x(1 + \log x)^{3/5}]$ for $x \geq 1$.

Finiteness of the moment of the order of the moment index is also preserved for given X and Y under addition (without requiring independence: just use the c_r -inequality), and, for independent X and Y , when $\alpha = \kappa(X) < \kappa(Y)$ so that $\kappa(\max(X, Y)) = \min(\kappa(X), \kappa(Y)) = \alpha$, we have $X \in \mathcal{M}^{\alpha+}$ if and only if $\max(X, Y) \in \mathcal{M}^{\alpha+}$.

In work underlying Scheller-Wolf (2003), interest centres on independent nonnegative r.v.s X and Y with finite positive moment indexes α and β for which both

$$E(X^\alpha) = \infty = E(Y^\beta) \quad (27)$$

and

$$E([\min(X, Y)]^{\alpha+\beta}) = \infty \quad (28)$$

hold. Clearly, $E([\min(X, Y)]^{\alpha+\beta-\epsilon}) < \infty$ for arbitrary $0 < \epsilon < \alpha + \beta$, and elementary algebra yields the rest of Lemma 8.

Lemma 8. *Let independent nonnegative r.v.s X and Y satisfy $\kappa(X) = \alpha \in (0, \infty)$, $\kappa(Y) = \beta \in (0, \infty)$, and Condition (27). If also Condition (28) holds for given X and all Y as described, then $\kappa(\min(X, Y)) = \kappa(X) + \kappa(Y)$, and the d.f. F of X satisfies (3).*

A sufficient condition on X for (27) to imply (28) is that

$$\liminf_{x \rightarrow \infty} x^\alpha \bar{F}(x) > 0. \quad (29)$$

It remains to consider conditions under which (27) does *not* imply (28). Let $X \in \mathcal{M}^{\alpha-}$, and suppose that for some $\epsilon > 0$ and positive integer $k \geq 2$, its d.f. F satisfies

$$\limsup_{x \rightarrow \infty} (\text{Log}_{k-1}(x))^\epsilon x^\alpha \bar{F}(x) < \infty, \quad (30)$$

where the positive monotonic nondecreasing function $\text{Log}_k(x)$ is defined for $x > 0$ by

$$\text{Log}_k(x) = \begin{cases} \max(1, x) & (k = 0), \\ \max(1, \log \text{Log}_{k-1}(x)) & (k = 1, 2, \dots) \end{cases}$$

(this function Log_k is similar to but not the same as the functional iterate of \log denoted \log_k and defined in [BGT 1.3.3]). We assert that there exists a r.v. Y satisfying (27) such that $E([\min(X, Y)]^{\alpha+\beta}) < \infty$.

To check this assertion, observe that the function $\text{Log}_k(x)$ has derivative

$$(\text{Log}_k(x))' = \begin{cases} 0 & \text{if } \text{Log}_k(x) = 1, \\ \frac{(\text{Log}_{k-1}(x))'}{\text{Log}_{k-1}(x)} = \frac{1}{\text{Log}_{k-1}(x) \cdots \text{Log}_0(x)} & \text{otherwise.} \end{cases}$$

In the latter case, this function is monotone decreasing to 0 as $x \rightarrow \infty$. Its product with $(\max(1, x))^{-(\beta-1)}$, where $\beta \geq 1$, can therefore be taken as the upper tail \bar{G} say, of the d.f. of a nonnegative r.v. Y_L say for which $E(Y_L^\beta) = \infty$, and $E(Y_L^{\beta-\epsilon}) < \infty$ for every positive $\epsilon \leq \beta$. Thus, $Y_L \in \mathcal{M}^{\beta-}$.

From (30) it follows that for some finite positive A ,

$$\bar{F}(x) \leq Ax^{-\alpha} (\text{Log}_{k-1}(x))^{-\epsilon}.$$

Then $E([\min(X, Y_L)]^{\alpha+\beta}) < \infty$ because the function $\bar{F}(x)\bar{G}(x)$, which equals the tail of the d.f. of the r.v. $\min(X, Y_L)$, has $x^{\alpha+\beta-1}\bar{F}(x)\bar{G}(x)$ bounded above for sufficiently large x , $x \geq x_0$ say, by

$$\frac{A}{(\text{Log}_{k-1}(x))^{1+\epsilon} \text{Log}_{k-2}(x) \cdots \text{Log}_0(x)},$$

and this function, being the derivative of $-A(\text{Log}_{k-1}(x))^{-\epsilon}/\epsilon$, is integrable on (x_0, ∞) .

References

- Baltrūnas, A., Daley, D.J. and Klüppelberg, C. (2004), Tail behaviour of the busy period of a GI/G/1 queue with subexponential service times, *Stoch. Proc. Appl.* **111**, 237–258.
- Bingham, N.H., Goldie, C.M. and Teugels, J.L. (1989), *Regular Variation* (Cambridge U.P., Cambridge, UK, pbk. ed. with additions).
- Daley, D.J. (2001), The moment index of minima. In *Probability, Statistics and Seismology (A Festschrift for David Vere-Jones)*, J. Appl. Probab. **38A**, 33–36.
- Goncalves, P. and Riedi, R.H. (2005). Diverging moments and parameter estimation. *J. Amer. Statist. Assoc.* (to appear).
- Scheller-Wolf, A. (2003), Necessary and sufficient conditions for delay moments in FIFO multiserver queues with an application comparing s slow servers with one fast one, *Operat. Res.* **51**, 748–758.