

Sussex Research

A rapid non-radioactive technique for measurement of repair synthesis in primary human fibroblasts by incorporation of Ethynyl deoxyuridine (EdU)

Siripan Limsirichaikul, Atsuko Niimi, Heather Fawcett, Alan Lehmann, Shunichi Yamashita, Tomoo Ogi

Publication date

01-03-2009

Licence

This work is made available under the CC [BY-NC](https://creativecommons.org/licenses/by-nc/4.0/) 4.0 licence and should only be used in accordance with that licence. For more information on the specific terms, consult the repository record for this item.

Citation for this work (American Psychological Association 7th edition)

Limsirichaikul, S., Niimi, A., Fawcett, H., Lehmann, A., Yamashita, S., & Ogi, T. (2009). A rapid nonradioactive technique for measurement of repair synthesis in primary human fibroblasts by incorporation of Ethynyl deoxyuridine (EdU) (Version 1). University of Sussex. https://hdl.handle.net/10779/uos.23313545.v1

Published in Nucleic Acids Research

Link to external publisher version

<https://doi.org/10.1093/nar/gkp023>

Copyright and reuse:

This work was downloaded from Sussex Research Open (SRO). This document is made available in line with publisher policy and may differ from the published version. Please cite the published version where possible. Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners unless otherwise stated. For more information on this work, SRO or to report an issue, you can contact the repository administrators at [sro@sussex.ac.uk.](mailto:sro@sussex.ac.uk) Discover more of the University's research at <https://sussex.figshare.com/>

Sussex Research Online

A rapid non-radioactive technique for measurement of repair synthesis in primary human fibroblasts by incorporation of Ethynyl deoxyuridine (EdU)

Article (Unspecified)

Limsirichaikul, Siripan, Niimi, Atsuko, Fawcett, Heather, Lehmann, Alan, Yamashita, Shunichi and Ogi, Tomoo (2009) A rapid non-radioactive technique for measurement of repair synthesis in primary human fibroblasts by incorporation of Ethynyl deoxyuridine (EdU). Nucleic Acids Research, 37 (4). ISSN 0305-1048

This version is available from Sussex Research Online: http://sro.sussex.ac.uk/id/eprint/2115/

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

A rapid non-radioactive technique for measurement of repair synthesis in primary human fibroblasts by incorporation of Ethynyl deoxyuridine (EdU)

 $\overline{1}$

A rapid non-radioactive technique for measurement of repair synthesis in primary human fibroblasts by incorporation of Ethynyl deoxyuridine (EdU)

Siripan Limsirichaikul^{1, 2}, Atsuko Niimi^{1, 3}, Heather Fawcett³, Alan Lehmann³⁺, Shunichi Yamashita¹⁺ and Tomoo Ogi¹*

¹Department of Molecular Medicine, Atomic Bomb Disease Institute, Nagasaki University, Nagasaki, 852-8523 Japan

²Department of Biopharmacy, Faculty of Pharmacy, Silpakorn University, Nakhon Pathom, 73000 Thailand.

³Genome Damage and Stability Centre, University of Sussex, Falmer, Brighton, BN1 9RQ, United Kingdom

*corresponding author: togi@nagasaki-u.ac.jp / Tel: $+81-95-819-7103$ / Fax: $+81-95-819-7104$
+These authors contributed equally to this work.

ABSTRACT

Example 12 togi@nagasaki-u.ac.jp / Tel: +81-95-819-7103 / uted equally to this work.

mentosum (XP) is an autosomal recessive genetic

sun-sensitivity and skin cancer predisposition. X

mt nucleotide excision repair (NER Xeroderma pigmentosum (XP) is an autosomal recessive genetic disorder. Afflicted patients show extreme sun-sensitivity and skin cancer predisposition. XP is in most cases associated with deficient nucleotide excision repair (NER), which is the process responsible for removing photolesions from DNA. Measuring NER activity by nucleotide incorporation into repair patches, termed 'unscheduled DNA synthesis (UDS)', is one of the most commonly used assays for XP-diagnosis and NER research. We have established a rapid and accurate procedure for measuring UDS by replacement of thymidine with 5-ethynyl-2'-deoxyuridine (EdU). EdU incorporated into repair patches can be directly conjugated to fluorescent azide derivatives, thereby obviating the need for either radiolabelled thymidine or denaturation and antibody detection of incorporated bromodeoxyuridine (BrdU). We demonstrate that the EdU incorporation assay is compatible with conventional techniques such as immunofluorescent staining and labeling of cells with micro-latex beads. Importantly, we can complete the entire UDS assay within half a day from preparation of the assay coverslips; this technique provides a rapid and convenient non-radioactive UDS assay for XP-diagnosis and NER related research.

INTRODUCTION

 Cells are continuously exposed to various types of DNA damage induced by both endogenous and exogenous sources. Consequently, all living organisms have developed DNA repair mechanisms to ensure that the integrity of the genetic information is maintained (1). Nucleotide excision repair (NER) is one of the most versatile DNA repair systems, which deals with the major UV photoproducts, as well as many DNA adducts arising from exposure to a variety of chemical carcinogens (2,3). NER involves removal of damaged nucleotides from the DNA by dual incisions, and filling and sealing of the residual single strand gaps by DNA polymerases and ligases (1-3).

 Xeroderma pigmentosum (XP) is the prototype NER deficient genetic disorder (2-4), the incidence of which is 1 in 80,000 to 1,000,000 in the general population, depending on the location (5). XP patients suffer from severe photosensitivity, a high incidence of sunlightinduced skin cancer, and in some cases neurological abnormalities. In $\sim80\%$ of XP cases, the patients have a defect in one of the genes responsible for NER. XP-variant patients, who comprise the residual 20% of XP cases, are proficient in NER but have a defect in post-

replication repair (PRR) as they lack the specialised DNA polymerase, pol η, which accurately replicates past cyclobutane pyrimidine dimers (CPDs) (6).

 Cockayne Syndrome (CS) and Trichothiodystrophy (TTD) are also NER-deficient disorders (4,7,8). NER consists of two sub-pathways: global genome repair (GGR) is relatively slow but functions genome-wide (3), whereas transcription coupled repair (TCR) is a faster process that is specialised for the transcribed strand of actively transcribed regions (9,10). With the exception of XPC and XPE, which are used exclusively in GGR (3), all genes defective in patients with XP, the combined features of XP and CS (XP/CS), and TTD are associated with both pathways, whereas in CS only TCR is compromised (7,9-11). NER-deficient patients have been so far assigned to at least eleven complementation groups (XP-A, to G, CS-A, and B, TTD-A, and ERCC1), which correspond to 11 different NER damage recognition and incision proteins (3,4).

ased assay for assessing NEK-deficiency entails n
sociated with DNA repair activity. GGR, which c
eed by damage induced non-S phase gap-filling D
thesis (UDS)' (12), while TCR activity can be me
levels after DNA damaging t The most commonly used assay for assessing NER-deficiency entails measuring nucleotide incorporation levels associated with DNA repair activity. GGR, which contributes to \sim 90% of NER activity, is assessed by damage induced non-S phase gap-filling DNA replication, termed 'unscheduled DNA synthesis (UDS)' (12), while TCR activity can be measured by 'recovery of RNA synthesis (RRS)' levels after DNA damaging treatment (13). Currently, a variety of methods for measuring UDS have been established (8,12,14-17). 254nm UVC-irradiation is commonly used for inducing DNA damage, and cells are then incubated in the presence of either radiolabelled thymidine or a nucleoside analog. The most frequently used methods in XP-diagnostic laboratories are: $\left[1\right]$ ³H-thymidine incorporation, followed by evaluation of grain counting after autoradiography of tissue-culture coverslips: this technique is accurate and compatible with most immunostaining based methods, but the procedures are labor-intensive and time-consuming (15,16). [2] Bromodeoxyuridine (BrdU) incorporation, followed by alkaline denaturation of DNA, and immunofluorescent detection by anti-BrdU antibody: this is a convenient variant of method [1]; replacement of ${}^{3}H$ -thymidine by BrdU diminishes the time needed for the assay, however it has reduced sensitivity \sim 50% reduction of UDS activity is not detectable)(18). [3] Liquid scintillation counting of ³H-thymidine incorporation into repair patches: although this method is rapid, the quantitative capacity is lower than the other methods as this is a batch assay. Furthermore, replicative DNA synthesis has to be abolished by using non-dividing cells and the DNA replication inhibitor, hydroxyurea (HU), during the assay (12).

 In this report, we have employed 5-ethynyl-2'-deoxyuridine (EdU), an alkyne conjugated nucleoside analogue of thymidine, as an alternative to radioactive thymidine or BrdU (19). In contrast to the BrdU based UDS assay, EdU can be directly conjugated to fluorescent azide (20), and thus, the assay does not require denaturing of DNA nor the use of antibodies. We have successfully improved the sensitivity, which is one of the weak points of BrdU based methods. Furthermore, the total time for the assay was dramatically reduced even compared to the BrdU based assay. Using this technique, a standard UDS assay can be completed within half a day from preparation of the assay coverslips.

MATERIALS AND METHODS

Cell strains and cultures

All the cells used in this report are human primary fibroblasts. Normal 1BR (21), 48BR (21), 142BR (22), and 251BR (22), XP-patient-derived XP12BR (XP-D) (this study), XP15BR (XP-A) (22), XP13BR (XP-C) (this study) and XP20BE (XP-G) (23), and CS-patient-derived CS10LO (CS-B) (24) cells were used. Cells were cultured in Dulbecco's Modified Eagle

Medium (DMEM) supplemented with 10% FBS (Gibco-BRL) and penicillin-streptomycin (Wako), and maintained at semiconfluent density during the experiments unless otherwise stated. For labeling with latex-beads, 48BR cells were cultured with a few drops of 0.5 μ m diameter polybead Hydroxylate (Polysciences) in DMEM for 3 days.

UV-induced UDS measurement in normal primary human fibroblasts by Ethynyl deoxyuridine (EdU) incorporation

often contains thymidine, which competes with E
then washed with PBS, followed by fixation and p
rmaldehyde, 0.5% triton X-100, and 300 mM succ
ig with PBS, cells were blocked with 10% FBS ir
detected by fluorescent-azide 5-ethynyl-2'-deoxyuridine (EdU) is available from Invitrogen (19). To optimize UVinduced UDS by EdU incorporation, effects of UV-dose and EdU incubation period were examined. 48BR cells were cultured on coverslips and maintained at confluent density. Cells were washed with PBS, followed by irradiation with different doses $(5{\text -}20 \text{ J/m}^2)$ of UVC (254nm). After UV-irradiation, cells were immediately incubated with serum free DMEM supplemented with 10 μ M EdU for different periods (0.5, 1, 2, and 4 h). Serum-free medium was used since serum often contains thymidine, which competes with EdU for incorporation into DNA. Cells were then washed with PBS, followed by fixation and permeabilisation with PBS containing 2% formaldehyde, 0.5% triton X-100, and 300 mM sucrose for 20 minutes. After extensive washing with PBS, cells were blocked with 10% FBS in PBS for 30 minutes. Incorporated EdU was detected by fluorescent-azide coupling reaction (Click-iT, Invitrogen) (19). Briefly, cells were incubated for 30min with azide-conjugated Alexa Fluor-488 dye in TBS supplemented with 4mM CuSO ⁴. Cells were then washed 3 times with PBS containing 0.05% Tween20 (PBST). Coverslips were finally soaked with PBS, fixed with 3.7% formaldehyde in PBS for 20min, and mounted on glass slides with Aquapolymount (Polysciences). Photographs of the cells were captured with a fluorescent microscope equipped with a CCD camera (BIOREVO9000-KEYENCE), and captured images were processed and analysed with ImageJ software (NIH). At least 50 non-S-phase cells were randomly selected from a single captured field, and the average nuclear fluorescent intensity was calculated. Data points presented in the text are the averages calculated from five different fields. We found that 20J/m² UV-irradiation followed by 2 h EdU incubation was the optimal condition for the UDS assay. This condition was used for all experiments except for Figure 1A.

BrdU incorporation assay

Cells were cultured and UVC irradiated $(20J/m²)$, and incubated in conditions identical to the EdU incorporation assay described above (5 µM BrdU, instead of EdU, was supplemented). Cells were then washed with PBS, followed by fixation and permeabilisation in PBS containing 2% formaldehyde, 0.5% tritonX-100, and 300 mM sucrose for 20 min. After extensive washing with PBS, cells were treated with 4 M HCl in PBS for 15 min for denaturing DNA. Cells were extensively washed with PBS for neutralisation, followed by blocking with 10% FBS in PBS for 30 min. Incorporated BrdU was detected by 1 h incubation with mouse anti-BrdU antibody (BD, diluted 1:150 in PBST). Cells were then washed 3 times with PBST, followed by 1 h incubation with Alexa Fluor-488 conjugated goat anti-mouse IgG (Invitrogen, diluted 1:500 in PBST). Coverslips were finally soaked in PBS, fixed with 3.7% formaldehyde in PBS for 20 min, and mounted on glass slides with Aquapolymount (Polysciences). Image capture and analyses were performed as described above.

$3H$ -thymidine incorporation assay

 Details of the assay have been previously described (12). Cells were cultured in 1% DMEM for three days $(3 \times 10^5 \text{ cells per 5cm-diameter dish})$. They were then incubated for 1h with 10mM hydroxyurea (HU), UV irradiated and incubated for a further 3h with 1% DMEM containing 10μ Ci/ml 3 H-thymidine and 10 mM HU. Incorporated 3 H-thymidine incorporated into acid-insoluble materials was measured by liquid scintillation counting.

58 59 60

Immunostaining

48BR and XP20BE cells were mix-cultured at 1:1 ratio (48BR alone for ki67 staining). UV-irradiation, EdU incubation, and fixing and permeabilisation steps were as described above. For antibody detection, cells were blocked with 10% FBS in PBS for 30 min, followed by 1 h incubation with mouse monoclonal anti-XPG (8H7, Santa Cruz Biotechnology), or rabbit monoclonal anti-ki67 (SP6, Thermo scientific) antibodies (diluted 1: 100 in PBST). Cells were then washed 3 times with PBST, followed by 1 h incubation with secondary antibodies: Alexa Fluor-594 conjugated goat-anti-mouse-, and goat-anti-rabbit-IgG, for XPG and ki67 detection, respectively (Invitrogen, diluted 1:1000 in PBST). To avoid unnecessary nuclear fluorescence, DAPI staining was omitted. After extensive washing with PBST, cells were fixed for 20 min with 3.7% formaldehyde in PBS. Subsequently, EdU detection was carried out as described above.

RESULTS
Effects of UV-dose and EdU-incubation period on measurement of UV-induced UDS in normal human primary fibroblasts

Example 15
For Form Control Set Comparison
Formal Set Constant Constant Constant Constant Constant
Formal Set Constant Summan fibroblasts were cultured on cover

Set Summan fibroblasts were cultured on cover

Inter 48BR normal primary human fibroblasts were cultured on coverslips and subsequently exposed to different doses of UVC, followed by incorporation of EdU for different time periods. UVC treatment elicited UDS activity as shown by an increase in nuclear fluorescence, after conjugation of fluorescent azide to incorporated EdU (Figures 1A, 1B). As reported in previous publications employing ³H-thymidine labeling and autoradiography (15,16), strong signals from scheduled DNA synthesis in S-phase cells can be distinguished from much weaker cell-cycle independent unscheduled EdU incorporation, resulting from repair synthesis (Figure 1B). Additionally, the fluorescent-azide coupling reaction seems specific to incorporated EdU, as we barely detected any non-specific cytoplasmic staining or DNA replication independent nuclear signals (Figure 1B). We observed that the nuclear fluorescent intensities were proportional to both UV dose and EdU incubation time, indicating that this endpoint satisfactorily represents the amount of incorporated EdU, which is directly convertible to semiquantitative UDS measurement (Figure 1A). We were able to detect nuclear fluorescent signals at relatively low UV exposures, and with short time-periods of post UV EdU incubation, demonstrating that this technique is sensitive enough to detect low levels of UDS activity.

Sensitivity and resolution of the EdU based UDS assay

We evaluated the resolution as well as the sensitivity of the EdU based UDS assay, by comparing the results obtained from the EdU technique with that of conventional autoradiography and BrdU based methods. Many publications that have utilized autoradiography for the UDS assay report UDS levels for the most severe XP-cases (i.e. XP-A, or XP-G/CS) as $\leq 5\sim 10\%$ of normal levels (25). These can be distinguished from cell strains from XP patients, typically from XP-C or XP-D groups, that often have UDS levels of 20~40% of residual UDS activity (intermediate UDS level). The intensity distribution plots for the EdU based UDS assay for both normal 48BR (UDS positive, Figure 1C) and XP-deficient XP15BR (UDS negative, Figure 1D) fibroblasts resemble to those of published autoradiography based experiments, though we acknowledge that the background UDS levels were relatively higher than that from autoradiography; this point is discussed later. The EdU assay consistently produced a satisfactory S.D. of 10~15%.

We compared the relative sensitivities of the EdU assay and the anti-BrdU based method. Though BrdU has been commonly used for S-phase labeling, publications using it for UDS

 $\mathbf{1}$

assays are quite limited because of its restricted sensitivity and resolution (18). (Figures 1E, 1F). Although the histograms of normal 48BR and XP-deficient XP15BR fibroblasts could be discriminated by the BrdU based assay (Figures 1E, 1F), the resolution as well as S.D. were both markedly improved in the EdU method, suggesting that EdU is a more sensitive substitute for BrdU in the UDS assay.

UV-induced EdU incorporation in NER-deficient primary fibroblasts

To apply the EdU technique to XP diagnosis, we estimated the levels of UV-induced EdU incorporation in several NER-deficient primary fibroblasts as well as normal controls (Figures 2A-2G for histograms, and 2I for typical heterogenous disorder, both the NER genes affected and the types of mutations determine the magnitude of the defect in UDS. We measured UDS in XP15BR (XP-A), XP20BE (XP-G/CS), XP13BR (XP-C), XP12BR (XP-D), and CS-patient-derived CS10LO (CS-B) cells, and compared the results with those obtained using ³H thymidin irradiation, we observed substantial UDS in normal fibroblasts, 48BR (Figure 2A) and 1BR (Figure 2B), as measured by EdU incorporation. We barely detected any UDS in severe XP patient-derived XP15BR (XP-A, Figure 2C) and XP20BE (XP-G, Figure 2D) fibroblasts, while we detected marginal (~20% of normal) UDS activity in XP-C patient-derived XP13BR (Figure 2E) fibroblasts.

For the UDS in normal fibroblasts, A8BR (IT) and the solution of the same of the SIR (C-20% of normal JUDS activity in XP-C patient-compaction. We barely detected any R (XP-A, Figure 2C) and XP20BE (XP-G, Figure 2C) and XP XP12BR is homozygous for R511Q mutation in the XPD gene (our unpublished data), and, as shown in Figure 2F, we found approx. 40% of normal UDS activity. In Figure 2H, we show the data obtained with the same set of cell strains using the ³H thymidine incorporation. In good agreement with the EdU assay, UDS was barely detectable in XP15BR, but were about \sim 20% and ~40% of normal in XP13BR and XP12BR, respectively. These results show good agreement between the two methods and demonstrate that cells exhibiting intermediate UDS activity can be distinguished from both normal and severe UDS defects by the EdU based assay.

 We then further examined whether CS-deficient fibroblasts were distinguishable from normal cell-lines by the UDS assay. CS is only compromised in the TCR pathway of NER, and this is not precisely diagnosed by UDS assays, since the deficiency is usually subtle. Though Figure 2G indicates a modest $(\sim 20\%)$ reduction of the UDS level compared to the normal fibroblasts, we can't conclude this to be significant and is likely to fall within the normal range.

 Levels of nuclear fluorescence after UVC irradiation in severe NER-deficient fibroblasts (Figures 2C, 2D) were virtually the same as those in unirradiated samples, suggesting that the background fluorescent signals resulting from minor EdU incorporation or non-specific fluorescent conjugation do not affect the analyses. We conclude that the EdU based UDS assay provides a specific measurement of nucleotide incorporation associated with the repair synthesis step of NER.

Compatibilities of the EdU based UDS assay with other techniques

 We anticipate applying the EdU based UDS assay system to clinical diagnoses, in which rigorous intra-assay controls are required. We therefore assessed the compatibility of the UDS assay with other standard techniques that use internal references. We found that fluorescent-azide conjugation to EdU is fully compatible with immunofluorescent staining (Figures 3A, 3B).

 Co-culture of an index fibroblast as well as a target under study on the same coverslip provides a rigorous internal control. In Figure 3A, normal and XPG-deficient fibroblasts were co-cultured, followed by UV-UDS assayed in combination with α -XPG immunostaining. EdU positive cells (excluding S-phase) perfectly correspond to XPG positive cells, indicating that two different cell populations can be distinguished in the UDS assay.

 Similarly, we examined if there was a cell-cycle preference for the EdU assay: coimmunostaining with a proliferation marker, ki-67, demonstrated that UDS can be detected in quiescent fibroblasts (no Ki67 stain) as well as proliferating populations (Figure 3B).

 As discussed above, the level of the defect in UDS activity varies between different XP patients. Ideally, ascertaining whether a patient is NER compromised is evaluated by comparing the residual UDS activity of the patient derived fibroblast with index cells. For this purpose, in a practical UDS based XP-diagnosis, normal fibroblasts have been preloaded with latex-beads in the cytoplasm and co-cultured with patient-derived fibroblasts. The 'beadlabeled-cells' provide an internal standard that can be detected under phase contrast microscopy and eliminates sample-to-sample staining variability (15). We tested if this technique is also compatible with the EdU assay. First we checked that the latex-beads did not affect the nuclear fluorescent intensity. As shown in Figures 4A (top panel) and corresponding histogram 4B, the fluorescent intensity in 48BR cells loaded with beads was practically the same as those in cells without beads, indicating that the beads do not interfere with the EdU based UDS assay. Modest (XP12BR, Figure 4A middle panel and 4C) as well as severe (XP15BR, Figure 4A bottom panel and 4D) UDS-deficient fibroblasts were both easily distinguishable from cocultured 48BR fibroblasts by their nuclear fluorescent levels. The UDS measurements with or without internal controls were nearly the same (compare Figures 2A, 2C, and 2F with 4B, 4D and 4C), suggesting that the prep-to-prep or coverslip-to-coverslip variation in the EdU based assay is small; this may be an advantage when applying the EdU assay to experiments that are incompatible with the use of an internal standard (e.g. fluorescent-based high throughput screening using GE's In-Cell-Analyzer or flow cytometry).

Further optimisation of the EdU assay

Factor and the beads of the Ball Linduce the background in the Ball Linduce of A middle panel and 4C) as well as severe (*UDS*-deficient fibroblasts were both easily distingates by their nuclear fluorescent levels. The UDS The EdU based experiments described above are sufficient for routine XP-screening as well as many types of NER research; however, further precise experiments such as distinguishing between no UDS activity and very low levels, may require higher sensitivity. We have attempted to increase the specific EdU uptake as well as to reduce the non-specific background. The incorporation of a thymidine analogue into the DNA is dependent on its concentration relative to endogenously synthesized thymine nucleotides in the nucleotide pool. Fluorodeoxyuridine (FdU) is an inhibitor of thymidylate synthetase, which renders the cell dependent on exogenously supplied thymidine or its analogs. We therefore used FdU and a longer incubation period (4 h) after UV irradiation. Additionally, we used stringent acid extraction with Bouin's fixative to try and reduce the background. As shown in Figure 5, although not dramatic, both background (compare white and red bars and their corresponding asterisks) and UDS specific EdU uptake (compare black and blue bars) were improved.

DISCUSSION

 Rapid and accurate assessment of UDS levels is indispensable for clinical XP-diagnosis as well as NER research. In this report, we have described a novel UDS assay system that provides a very simple method for measurement of repair replication activity in human cells. Orthodox UDS assay methods have been based on 3 H-thymidine incorporation (12,15,16). Although a variety of alternative reagents have been developed, most XP-diagnosis laboratories still measure UDS by autoradiography or direct scintillation counting of incorporated 3 H-thymidine. The biggest disadvantages of using 3 H-thymidine for UDS assay is its time-consuming procedures and the requirement for laboratories designated for radioactive work.

Page 7 of 16 Nucleic Acids Research

60

 The autoradiographic process provides an accurate measurement of UDS activity as each silver-grain represents an individual event of nucleotide incorporation into a repair patch. However, the experiment is complicated, requiring significant skills, and takes 1-2 weeks to be completed; furthermore, counting of silver-grains under the microscope is laborious. An alternative procedure uses direct scintillation counting of 3 H-thymidine incorporation (12). This method requires abrogation of S-phase DNA replication. To accomplish this, cells are brought into quiescence by serum starvation, and hydroxyurea is needed during the nucleotide incorporation period, to block replicative DNA synthesis. This technique significantly reduced the time required for the UDS assay, but quantitation of UDS is less accurate. In contrast to the above ³H-thymidine based methods, BrdU incorporation assay is faster and a convenient option. Indeed, BrdU has been frequently used for cell cycle analysis applications. On the other hand, regardless of the type of experiments, denaturation of DNA is needed to detect the incorporated BrdU as the anti-BrdU antibody is too big to access the epitope in native DNA. DNA denaturation and subsequent immunofluorescence detection sometimes generate an ambiguous staining pattern (data not shown, see high S.D. in Figure 1E, 1F). This is the biggest disadvantage of the BrdU incorporation assay: nucleotide incorporation into repair patches based on DNA repair activity is quite low and difficult to detect. For this reason, UDS based on BrdU has not been widely used.

naturation and subsequent immunotitionescence des

staining pattern (data not shown, see high S.D. if

tage of the BrdU incorporation assay: nucleotide

in DNA repair activity is quite low and difficult to

as not been wid The EdU incorporation method was originally developed for complementing known problems of S-phase population labeling by ³H-thymidine and BrdU (19). In EdU, the methyl group on the 5- position of thymidine is replaced by an alkyne group, and this analog is incorporated into DNA during replication. Subsequent fluorescent-azide coupling, termed 'click reaction' is rapid, sensitive, and specific (19,20): because of this property, the sensitivity and background fluorescent signal were both dramatically improved, and hence, EdU can be used for the UDS assay.

We have presented data demonstrating the usefulness of EdU for the UDS assay. Although the sensitivity of the EdU based method is not as high as the autoradiographic technique, we easily detected the defect in an XP-D cell-line with ~40% residual UDS activity and based on our data we conservatively estimate that we can detect a level of 10-20% of normal UDS.

 The EdU based UDS assay may have an advantage in high-throughput XP-diagnosis using flow cytometry. An attempt to use BrdU labeling and flow cytometry for UDS has been reported (14). Because of the improved sensitivity of EdU over BrdU, further introduction of multicolor labeling (i.e. normal internal control as well as specimen cells, with or without UV damage, all labeled with different fluorescent dyes) may provide a rapid and precise determination of the relative UDS levels.

 In this report, we have demonstrated that: [1] the accuracy and resolution of the UDS assay based on EdU incorporation was comparable to the conventional autoradiographic method: we could distinguish between a normal and an XP-deficient primary fibroblast when the cells were co-cultured on a single coverslip. [2] The assay was compatible with standard immunostaining and latex-bead-labeling, which provide proper internal controls. [3] The time required for the assay was dramatically reduced: the entire UDS assay can be completed within half a day from the preparation of specimen coverslips. The UDS assay based on EdU could potentially become a standard technique in NER research as well as XP-diagnosis.

FUNDING

 This work was supported by KAKENHI from Japan Society for the Promotion of Science [grant number 20810021] and a Butterfield Award from the Great Britain Sasakawa $\mathbf{1}$

Foundation to T.O.

ACKNOWLEDGEMENTS

 We are grateful to Dr. Nicolaas Jaspers for helpful comments on the manuscript. We also thank Dr. Motohiro Yamauchi for critical advice on the EdU incorporation assay.

FIGURE LEGENDS

associated with UDS activity, was analysed using
processing software. For each data point, at least
rror-bars indicate means of discrete nuclei fluores
tively. (**B**) Typical photos of the EdU assay are sl
 r^2 , +UV) or m Figure 1. UV-induced UDS assay in normal human primary fibroblasts. (A) Normal 48BR cells were cultured on coverslips, UVC-irradiated at the indicated doses, followed by incubation with 10 µM EdU for different time periods, fixation and conjugation of fluorescentdye to the incorporated EdU as described in Materials and Methods. The intensity of nuclear fluorescence, which is associated with UDS activity, was analysed using a fluorescence microscope and image processing software. For each data point, at least 250 nuclei were analysed. Points and error-bars indicate means of discrete nuclei fluorescent intensity and standard errors, respectively. (B) Typical photos of the EdU assay are shown. 48BR cells were UVC irradiated $(20J/m^2, +UV)$ or mock treated (-UV), followed by 2h incubation with EdU. EdU, coupled to Alexa fluor 488-azide; PH, phase contrast. (C-F) UDS performed by incorporation of EdU (C, D) or BrdU (E, F) was compared. XP15BR cells are primary fibroblasts from an XP-A patient. Cells were UVC irradiated (20J/m²), followed by incubation with 10 μ M EdU or 5 μ M BrdU for 2 h. Bars represent frequencies of the fluorescence levels in the indicated classes, with (black) or without (white) UVC irradiation. Asterisks indicate the mean values of nuclear fluorescent intensities, which correspond to the UDS levels. [∆] represents UDS difference between irradiated and unirradiated samples.

Figure 2. UV-induced EdU incorporation in non-S phase cells is NER repair replication specific. (A-G) Levels of UV-UDS in NER-proficient and -deficient cells were examined. Primary fibroblasts were cultured on coverslips, UVC irradiated $(20J/m²)$, followed by 2h incubation with 10 µM EdU. Levels of EdU incorporation were analysed as in Figure 1A and presented as in Figure 1. (H) UDS assay performed by ³H thymidine incorporation. Quiescent cells were UVC irradiated at the indicated dose and ³H thymidine incorporation was measured for 3h after UVC irradiation in the media containing 10mM hydroxyurea. UDS levels were normalised and expressed as percentages of the UDS in the normal cells at 10 J/m². Mean Normal represents the average UDS levels of the 4 different normal cell lines (1BR, 48BR, 142BR, and 251BR). (I) Typical photos of the EdU incorporation examined in Figure 2A-G are shown (2h after 20 J/m² UVC irradiation). Arrows indicate non-S phase cells. 48BR, normal; 1BR, normal; XP15BR, XP-A; XP20BE, XP-G; XP13BR, XP-C; XP12BR, XP-D; CS10LO, CS-B.

Figure 3. EdU assay is compatible with immunostaining. (A) Normal 48BR and XPGdeficient XP20BE fibroblasts were co-cultured on coverslips, UVC irradiated $(20J/m²)$, followed by 2h incubation with 10 µM EdU. Fixed cells were then immunostained with rabbit anti-XPG antibody (red), followed by conjugation of Alexa fluor 488-azide to incorporated EdU (green). (B) UV-induced EdU incorporation in quiescent cells. 48BR cells were UVC irradiated (20J/m²), followed by 2h incubation with 10 μ M EdU. Fixed cells were then immunostained with rabbit anti-ki67 antibody (red), followed by conjugation of Alexa fluor 488-azide to incorporated EdU (green). Triangle, normal arrow, and diamond arrow indicate Sphase, G0, and G1 (or G2/M), respectively. Detailed experimental conditions are described in Materials and Methods.

Figure 4. EdU assay in the presence of latex beads. Normal 48BR cells were pre-labeled with 0.5 µm diameter latex beads and co-cultured with indicated XP fibroblasts (no-beads) on coverslips. Cells were then UVC-irradiated $(20J/m²)$, followed by 2h incubation with 10 μ M EdU. Coverslips were then processed as in Figure 1. (A) Typical photos are displayed. White and red markings, arrows and arrow-heads represent 48BR and indicated XP cells, and non-S phase and S-phase cells, respectively. (**B-D**) Histograms of the UDS assay with the internal standard 48BR and (B) 48BR, normal; (C) XP12BR, XP-D; and (D) XP15BR, XP-A. Bars represent frequencies as described in Figure 1, with (black, 48BR; blue, XP cell-lines) or without (white, 48BR; red, XP cell-lines) UVC irradiation. Asterisks indicate the mean values of nuclear fluorescent intensities, which correspond the UDS levels. ∆ represents UDS differences with or without UVC irradiation.

It of the sensitivity by FdU supplement, longer in BR cells were cultured on coverslips, UVC irradia $μM EdU$ and $1uM FdU$ (Sigma) for 4 h. Cells we Acid extraction was performed using Bouin's fixa washing with PBS. Conjuga Figure 5. Improvement of the sensitivity by FdU supplement, longer incubation, and acid extraction. Normal 48BR cells were cultured on coverslips, UVC irradiated (20J/m²), followed by incubation with 10 µM EdU and 1uM FdU (Sigma) for 4 h. Cells were then fixed as indicated in Figure 2. Acid extraction was performed using Bouin's fixative (Sigma) for 30 min, followed by extensive washing with PBS. Conjugation of fluorescent-dye and detection of EdU signals were identical to the experiments in Figure 2. Bars represent frequencies of the fluorescence levels in indicated classes, with (black, identical to Figure 2A; blue, +FdU +4 h incubation +acid extraction) or without (white, identical to Figure 2A; red, +FdU +4 h incubation +acid extraction) UVC irradiation. Asterisks indicate the mean values of nuclear fluorescent intensities. ∆ represents UDS differences with or without UVC irradiation.

REFERENCES

- 1. Friedberg, E.C., Walker, G.C., Siede, W., Wood, R.D., Schultz, R.A. and Ellenberger, T. (2005) DNA Repair and Mutagenesis. 2 ed. ASM Press.
- 2. Bootsma, D., Kraemer, K.H., Cleaver, J.E. and Hoeijmakers, J.H. (2001) In Scriver, C. R., Beaudet, A. L., Sly, W. S. and Valle, D. (eds.), The Metabolic and Molecular Bases of Inherited Disease. 8 ed. McGraw-Hill, New York, pp. 677-703.
- 3. Gillet, L.C. and Scharer, O.D. (2006) Molecular mechanisms of mammalian global genome nucleotide excision repair. Chem Rev, 106, 253-276.
- 4. Kraemer, K.H., Patronas, N.J., Schiffmann, R., Brooks, B.P., Tamura, D. and DiGiovanna, J.J. (2007) Xeroderma pigmentosum, trichothiodystrophy and Cockayne syndrome: a complex genotype-phenotype relationship. Neuroscience, 145, 1388-1396.
- 5. Kleijer, W.J., Laugel, V., Berneburg, M., Nardo, T., Fawcett, H., Gratchev, A., Jaspers, N.G., Sarasin, A., Stefanini, M. and Lehmann, A.R. (2008) Incidence of DNA repair deficiency disorders in western Europe: Xeroderma pigmentosum, Cockayne syndrome and trichothiodystrophy. DNA Repair (Amst), 7, 744-750.
- 6. Masutani, C., Kusumoto, R., Yamada, A., Dohmae, N., Yokoi, M., Yuasa, M., Araki, M., Iwai, S., Takio, K. and Hanaoka, F. (1999) The XPV (xeroderma pigmentosum variant) gene encodes human DNA polymerase eta. Nature, 399, 700-704.
- 7. Lehmann, A.R. (2003) DNA repair-deficient diseases, xeroderma pigmentosum, Cockayne syndrome and trichothiodystrophy. Biochimie, 85, 1101-1111.
- 8. Stefanini, M., Lagomarsini, P., Giliani, S., Nardo, T., Botta, E., Peserico, A., Kleijer, W.J., Lehmann, A.R. and Sarasin, A. (1993) Genetic heterogeneity of the

excision repair defect associated with trichothiodystrophy. Carcinogenesis, 14, 1101-1105.

- 9. Laine, J.P. and Egly, J.M. (2006) When transcription and repair meet: a complex system. Trends Genet, 22, 430-436.
- 10. Sarasin, A. and Stary, A. (2007) New insights for understanding the transcriptioncoupled repair pathway. DNA Repair (Amst), 6, 265-269.
- 11. Cleaver, J.E. (2005) Cancer in xeroderma pigmentosum and related disorders of DNA repair. Nat Rev Cancer, 5, 564-573.
- 12. Lehmann, A.R. and Stevens, S. (1980) A rapid procedure for measurement of DNA repair in human fibroblasts and for complementation analysis of xeroderma pigmentosum cells. Mutat Res, 69, 177-190.
- 13. Lehmann, A.R., Thompson, A.F., Harcourt, S.A., Stefanini, M. and Norris, P.G. (1993) Cockayne's syndrome: correlation of clinical features with cellular sensitivity of RNA synthesis to UV irradiation. J Med Genet, 30, 679-682.
- 14. Vincent, F., Ceraline, J., Goldblum, S., Klein-Soyer, C. and Bergerat, J.P. (2001) A new flow cytometric method to follow DNA gap filling during nucleotide excision repair of UVc-induced damage. Cytometry, 45, 96-101.
- 15. Kleijer, W.J., van der Sterre, M.L., Garritsen, V.H., Raams, A. and Jaspers, N.G. (2007) Prenatal diagnosis of xeroderma pigmentosum and trichothiodystrophy in 76 pregnancies at risk. Prenat Diagn, 27, 1133-1137.
- 16. Stefanini, M., Giliani, S., Nardo, T., Marinoni, S., Nazzaro, V., Rizzo, R. and Trevisan, G. (1992) DNA repair investigations in nine Italian patients affected by trichothiodystrophy. Mutat Res, 273, 119-125.
- KINA synthesis to UV Irradiation. *J Med Genet,*
eraline, J., Goldblum, S., Klein-Soyer, C. and I
metric method to follow DNA gap filling durin-induced damage. Cytometry, 45, 96-101.
induced damage. Cytometry, 45, 96-101.
 17. Qiao, Y., Spitz, M.R., Guo, Z., Hadeyati, M., Grossman, L., Kraemer, K.H. and Wei, Q. (2002) Rapid assessment of repair of ultraviolet DNA damage with a modified host-cell reactivation assay using a luciferase reporter gene and correlation with polymorphisms of DNA repair genes in normal human lymphocytes. Mutat Res, 509, 165-174.
- 18. Droy, B.F., Miller, M.R., Freeland, T.M. and Hinton, D.E. (1988) Immunohistochemical detection of CCl4-induced, mitosis-related DNA synthesis in livers of trout and rat. Aquatic Toxicology, 13, 155-166.
- 19. Salic, A. and Mitchison, T.J. (2008) A chemical method for fast and sensitive detection of DNA synthesis in vivo. Proc Natl Acad Sci U S A, 105, 2415-2420.
- 20. Breinbauer, R. and Kohn, M. (2003) Azide-alkyne coupling: a powerful reaction for bioconjugate chemistry. Chembiochem, 4, 1147-1149.
- 21. Arlett, C.F., Green, M.H., Priestley, A., Harcourt, S.A. and Mayne, L.V. (1988) Comparative human cellular radiosensitivity: I. The effect of SV40 transformation and immortalisation on the gamma-irradiation survival of skin derived fibroblasts from normal individuals and from ataxia-telangiectasia patients and heterozygotes. Int J Radiat Biol, 54, 911-928.
- 22. Arlett, C.F., Plowman, P.N., Rogers, P.B., Parris, C.N., Abbaszadeh, F., Green, M.H., McMillan, T.J., Bush, C., Foray, N. and Lehmann, A.R. (2006) Clinical and cellular ionizing radiation sensitivity in a patient with xeroderma pigmentosum. Br J Radiol, 79, 510-517.
- 23. Moriwaki, S., Stefanini, M., Lehmann, A.R., Hoeijmakers, J.H., Robbins, J.H., Rapin, I., Botta, E., Tanganelli, B., Vermeulen, W., Broughton, B.C. et al. (1996) DNA repair and ultraviolet mutagenesis in cells from a new patient with xeroderma pigmentosum group G and cockayne syndrome resemble xeroderma pigmentosum cells. J Invest Dermatol, 107, 647-653.

>

- 24. Mallery, D.L., Tanganelli, B., Colella, S., Steingrimsdottir, H., van Gool, A.J., Troelstra, C., Stefanini, M. and Lehmann, A.R. (1998) Molecular analysis of mutations in the CSB (ERCC6) gene in patients with Cockayne syndrome. Am J Hum Genet, 62, 77-85.
	- 25. Kraemer, K.H., Coon, H.G., Petinga, R.A., Barrett, S.F., Rahe, A.E. and Robbins, J.H. (1975) Genetic heterogeneity in xeroderma pigmentosum: complementation groups and their relationship to DNA repair rates. Proc Natl Acad Sci U S A , 72, 59-63.

For Per Review

 $\mathbf 1$

Fig3

 $\mathbf 1$ \overline{c} $\overline{3}$ $\overline{\mathbf{4}}$ 5 6

C

*

Fluorescence intensity (a.u.)

ہی گھر ہوئی ہے۔ اس کے مطابق کی اس کے مطابق کے مطابق کے مطابق کی اس کے بعد اس کے مطابق کے مطابق کے مطابق کے مطا
اس کی اس کا اس

60

10 $\overline{5}$ $\ddot{\mathbf{0}}$

