

Sussex Research

Translesion DNA synthesis-assisted non-homologous end-joining of complex double-strand breaks prevents loss of DNA sequences in mammalian cells

Shay Covo, Jean-Pierre de Villartay, Penny Jeggo, Zvi Livneh

Publication date

01-11-2009

Licence

This work is made available under the **Copyright not evaluated** licence and should only be used in accordance with that licence. For more information on the specific terms, consult the repository record for this item.

Citation for this work (American Psychological Association 7th edition)

Covo, S., de Villartay, J.-P., Jeggo, P., & Livneh, Z. (2009). *Translesion DNA synthesis-assisted non-homologous end-joining of complex double-strand breaks prevents loss of DNA sequences in mammalian cells* (Version 1). University of Sussex. <https://hdl.handle.net/10779/uos.23313893.v1>

Published in

Nucleic Acids Research

Link to external publisher version

<https://doi.org/10.1093/nar/gkp703>

Copyright and reuse:

This work was downloaded from Sussex Research Open (SRO). This document is made available in line with publisher policy and may differ from the published version. Please cite the published version where possible. Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners unless otherwise stated. For more information on this work, SRO or to report an issue, you can contact the repository administrators at sro@sussex.ac.uk. Discover more of the University's research at <https://sussex.figshare.com/>

Translesion DNA synthesis-assisted non-homologous end-joining of complex double-strand breaks prevents loss of DNA sequences in mammalian cells

Article (Unspecified)

Covo, Shay, de Villartay, Jean-Pierre, Jeggo, Penny A and Livneh, Zvi (2009) Translesion DNA synthesis-assisted non-homologous end-joining of complex double-strand breaks prevents loss of DNA sequences in mammalian cells. *Nucleic Acids Research*, 37 (20). pp. 6737-6745. ISSN 0305-1048

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/2266/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**Translesion DNA Synthesis-Assisted Non-Homologous End-Joining of
Complex Double-Strand Breaks Prevents Loss of DNA Sequences
in Mammalian Cells**

Shay Covo^{1§}, Jean-Pierre de Villartay², Penny A. Jeggo³, and Zvi Livneh^{1*}

¹Department of Biological Chemistry, Weizmann Institute of Science, Rehovot 76100 Israel;

²Inserm, U768, Hôpital Necker-Enfants Malades, Paris, France; ³Genome Damage and Stability
Centre, University of Sussex, East Sussex BN1 9RQ, UK

[§]Current address: Laboratory of Molecular Genetics, National Institute of Environmental
Health Sciences, National Institutes of Health, Research Triangle Park, NC 27709, USA

*Incumbent of The Maxwell Ellis Professorial Chair in Biomedical Research. To whom
all correspondence should be addressed:

Dr. Zvi Livneh, Dept. of Biological Chemistry, Weizmann Institute of Science, Rehovot
76100, Israel.

Tel: 972-8-934-3203; Fax: 972-8-934-4169; Email: zvi.livneh@weizmann.ac.il

ABSTRACT

Double strand breaks (DSB) are severe DNA lesions, and if not properly repaired, may lead to cell death or cancer. While there is considerable data on the repair of simple DSB (sDSB) by non-homologous end-joining (NHEJ), little is known about the repair of complex DSBs (cDSB), namely breaks with a nearby modification, which precludes ligation without prior processing. To study the mechanism of cDSB repair we developed a plasmid-based shuttle assay for the repair of a defined site-specific cDSB in cultured mammalian cells. Using this assay we found that repair efficiency and accuracy of a cDSB with an abasic site in a 5' overhang was reduced compared to a sDSB. Translesion DNA synthesis (TLS) across the abasic site located at the break prevented loss of DNA sequences, but was highly mutagenic also at the template base next to the abasic site. Similar to sDSB repair, cDSB repair was totally dependent on XrccIV, and altered in the absence of Ku80. In contrast, Artemis appears to be specifically involved in cDSB repair. These results may indicate that mammalian cells have a damage control strategy, whereby severe deletions are prevented at the expense of the less deleterious point mutations during NHEJ.

INTRODUCTION

Double strand breaks (DSBs) are among the most detrimental types of DNA damage, and if not properly repaired, can lead to cell death or cancerous transformation (1). DSB are repaired in mammalian cells primarily by non-homologous end joining (NHEJ), in which the two broken parts of the chromosomes are ligated (For review see (2-4)). The core components in NHEJ include the Ku80/70 heterodimer, which protects DNA ends from degradation, and recruits other components of the repair machinery (5-8). The DNA-dependent protein kinase catalytic subunit (DNA-PKcs) joins Ku to form the DNA-PK holoenzyme, which connects the two termini of the break, and phosphorylates proteins involved in DNA damage response. Ligation is carried-out by the DNA ligase IV complex, composed of DNA Ligase IV, XRCC IV and Cernunnos/XLF all of which are essential for NHEJ (8-10). In addition to the core components of NHEJ, other enzymes participate in DSB repair mainly through DNA end processing prior to ligation. This includes the nuclease Artemis, which is activated by DNA-PK and involved in the repair of some DSBs (6,11,12), and DNA polymerases μ and λ , both implicated in DNA end-processing prior to ligation (6,13-16).

Most studies on NHEJ addressed simple DSB (sDSB), namely DNA breaks which did not include chemical modification of the base, sugar or phosphate moieties near the break point. However, some DNA damaging agents, most notably ionizing radiation, cause clusters of DNA damage, which often lead to modification in the vicinity of a DSB (17). For example, Winter and his colleagues have reported that abasic sites as well as other oxidized nucleotide derivatives reside in the vicinity of a DSB that was formed by ionizing radiation (18,19). The presence of modified bases in the vicinity of the DSB can interfere with its repair, and may therefore require special processing. Indeed, it has been observed that some DSB have slow repair kinetics in vivo

and in vitro, suggesting that their chemical structure hampers their repair (11,20). Despite their importance little is known on how cells cope with such complex DSB (cDSB), and specifically on how they are processed by the DSB repair machinery. Here we present a novel plasmid-based assay for the repair of a model cDSB in mammalian cells. We found that: (1) the presence of a single synthetic abasic site on a short 5'-overhang on one side of a DSB reduces the efficiency of DSB repair; (2) translesion DNA synthesis across the abasic site enables NHEJ with minimal loss of DNA bases; (3) remarkably, full-length TLS at the DNA terminus is highly mutagenic at the nucleotide following the abasic site, and it violates the A-rule at the abasic site; (4) *XrccIV* and Ku80 are involved in the repair of both sDSB and cDSB, whereas Artemis is involved specifically in the repair of cDSB in this assay system.

MATERIALS AND METHODS

Cell cultures and media

The human cell lines H1299 and PC3 were derived from a large cell lung carcinoma and a prostate carcinoma, respectively. They were cultured in RPMI 1640 supplemented with 10% FBS (GIBCO/BRL). 48BR normal, and Artemis-deficient CJ179, telomerase-immortalized fibroblasts (11,20) were grown in MEM medium with 10% FBS. Human Artemis deficient Guetel cells and their isogenic control Guetel/DA4 are skin fibroblasts transformed with SV40 and immortalized with telomerase (21). They were grown in RPMI 1640 10% FCS. The SV40-transformed NBS1 fibroblasts and their complemented controls (22) were grown in DMEM media supplemented with 10% fetal bovine serum, 2mM Glutamine, 100 units/ml penicillin and 100 µg/ml streptomycin. The CHO cell lines K1 (wild-type) and Xrs5 (*Ku80*^{-/-}) were purchased from ATCC, and XR-1 (*Xrcc4*^{-/-}) from Coriell cell repositories. All CHO cells were grown either

in MEM α or HAM medium with 10% FCS. All the cells were incubated at 37°C in a 5% CO₂ atmosphere.

DNA substrates

Construction of the linear plasmids with complex and simple DSB - The modified-linear plasmid LP41 was constructed by ligating a short duplex oligonucleotide carrying a site-specific synthetic abasic site to a restriction nuclease-cleaved plasmid (Fig. 1s). The short duplex oligonucleotide was prepared by annealing oligonucleotides 5'-AXCAGACCTGCGTGTACCG-3' (X is the abasic site) and 5'-ACACGCAGGTCTG-3' (200 pmol each) in 20 μ l of a solution containing 10 mM Tris-HCl (pH 7.5), 1 mM EDTA, and 150 mM NaCl by heating to 70° C for 10 min, and then cooling to room temperature over a period of 2-3 h. The control non-modified linear plasmid LP40 was prepared in a similar way, except that the ligation was to a control short duplex oligonucleotide prepared by annealing onligonucleotides 5'-AACAGACCTGCGTGTACCG-3' and 5'-ACACGCAGGTCTG-3'. The vector was the 3000 bp *Bst*XI-*Bgl*II fragment of plasmid pSKSL (23), to which EcoRV site was introduced (pSKSL-EcoRV-*Bgl*II) (Fig. 1s). The vector was obtained by cleavage of the plasmid with *Bst*XI and *Bgl*II (see Fig. 1s A), followed by fractionation on a 0.8% agarose gel, and purification by electroelution using the BIOTRAP* device (Schleicher & Schuell).

Preparative ligation of the insert to the vector was carried out in a 7.5:1 molar ratio, with 350 ng/ μ l vector DNA, and 1.5 unit/ μ l T4 ligase (New England Biolabs) at 16° C over night. Next, the ligase was heat inactivated at 65° C for 10 min, and the efficiency of ligation was examined. This was done by digesting 1 μ g of the linear plasmid with restriction nuclease Afe I for 3hr, followed by dephosphorelation, and then 5' end labeling with ³²P using polynucleotide

kinase and [γ - 32 P] ATP. The fragments were then separated on 6% PAGE. The ligated DNA fragment migrates slower than the non-ligated one. Ligation efficiency was at least 90% as determined by phosphorimaging (Fig. 2s). After confirming the ligation efficiency the linear plasmid was digested with EcoRV, which removed 10 nucleotides from one terminus, leaving a blunt end (Figs. 1B and 1s C). Finally, the DNA was again gel purified. Plasmid pSA26 (3,356 bp long; cm^R) was previously described (23).

NHEJ assay for cDSB in cultured cells

The assay for the repair of cDSB included the following steps (Fig. 1): (i) Transfection of the mammalian cells with a plasmid mixture containing the modified linear plasmid (LP41; kan^R), the internal control plasmid pSA26 (cm^R), and the carrier plasmid pUC18. As a control, a parallel transfection was performed with the non-modified linear plasmid, LP40, pSA26 and pUC18. (ii) Extraction of the plasmids from the mammalian cells. (iii) Transformation of an *E. coli* indicator strain with the plasmid mixture. (iv) Deduction of the extent of cDSB repair from the number of transformants. More specifically, mammalian cells were co-transfected with a DNA mixture containing 150-350 ng of linear plasmid, 3-24 ng of the internal control plasmid pSA26, and 10 μ g of the carrier plasmid pUC18. Transfection of H1299 cells was done using the JetPEI (Polyplus transfection), whereas other cells were transfected by electroporation using the Nucleofector system (Amaxa; Koln, Germany) with Nucleofector kits T (for CHO cells) and V and R (for other human cells). The cells were incubated in their appropriate medium for 16-24 hr at 37°C in a 5% CO₂ atmosphere. At the end of the incubation the cells were collected, washed 3 times with PBS and their plasmid content was extracted by using a plasmid purification kit based on alkaline lysis (Wizard miniprep, Promega). The recovered plasmids were electroporated into

150 μ l of competent *E. coli* JM109*recA* cells, followed by incubation in 1 ml SOC medium at 37°C for 1 h, and then plating in parallel on LB-agar plates containing kanamycin (50 μ g/ml) or chloramphenicol (30 μ g/ml). Relative repair of the cDSB was calculated based on the number of colonies, as described below. When desired, plasmids were extracted from kan^R colonies **originating from several independent transfection experiments**, and subjected to DNA sequence analysis at the vicinity of the break point.

RESULTS

Outline of the assay for cDSB repair in cultured cells

The cDSB repair assay measures the ability of cells in culture to circularize a transfected linearized plasmid that carries on one end a two-nucleotides 5' overhang with a site-specific abasic site next to the terminal nucleotide, whereas the other end is blunt (LP41, kan^R; Fig. 1). In order to enable quantification, the mammalian cells were co-transfected with a plasmid mixture containing the modified linearized plasmid (LP41, kan^R), an intact plasmid as an internal normalizing reference (pSA26; cm^R), and a carrier plasmid (pUC18). As a control, cells were transfected in parallel with a similar mixture, wherein the modified linearized plasmid was replaced with a non-modified linearized control plasmid (LP40 kan^R). The measurement of the linear plasmid repair was done in a subsequent step, in which the plasmid mixture was isolated from the mammalian cells, and used to transform an indicator *E. coli* strain. The transformants were plated in parallel on LB plates containing kanamycin, to select for cells harboring repaired plasmids that had originally carried a DSB, or chloramphenicol, to select for cells carrying the normalizing intact plasmid (Fig. 1A). Linear plasmids poorly transform *E. coli*, and therefore the *E. coli* colonies observed had been formed by a circular (repaired) plasmid molecule. To further

decrease the background transformation of linear plasmids, isolation of the plasmid content of the mammalian cells was done under mild alkaline conditions followed by neutralization, which cause denaturation of linear, but not covalently closed circular plasmids.

The efficiency of DSB repair in mammalian cells was calculated based on the number of *E. coli* transformants. First, the number of LP41 kan^R transformants was divided by the number of pSA26 cm^R transformants, to yield a normalized DSB repair efficiency for the cDSB. The same was done for the pair of LP40 and pSA26 transformants, to obtain the normalized DSB repair efficiency for the sDSB. We then divided the two to get the relative repair efficiency of the cDSB compared to sDSB. Importantly, none of the plasmids can be replicated within the mammalian cells, hence number of *E. coli* transformants correlates to the extent of repair within the mammalian cells.

A single abasic site near a DSB reduces both the efficiency and accuracy of DSB repair

To examine whether linearized plasmids with a cDSB are repaired in human cells, we performed a series of experiments with the human lung cancer cell line H1299. As can be seen in Table 1, which presents the results of 7 independent experiments, the number of kan^R and cm^R *E. coli* transformants obtained after extraction of the plasmids mixture from the human cells varied considerably among experiments. However, the relative repair of cDSB (relative to the sDSB) was remarkably reproducible, yielding a value of 20%±2%. Thus, the assay yields quantitative and reproducible measurements of the repair of the cDSB relative to the sDSB. The value obtained indicates that the presence of a single modification near the break points significantly decreased its repair by 5-fold relative to a sDSB.

To analyze the accuracy of DSB repair we performed DNA sequence analysis of repaired descendents of the two types of linearized plasmids. As can be seen in Fig. 2, 90% (45/50 events) of the sDSB (substrate LP40) were accurately repaired in H1299 cells, with restoration of the original DNA sequence. Largely accurate repair of sDSB was observed also in human 48BR cells (80%), and to lesser extent in the CHO K1 cells (57%; Fig. 2). The picture dramatically changed for cDSB (substrate LP41). In H1299 cells only 16% (19/123 events) of the repair events preserved the correct length of the plasmid, whereas 84% contained either deletions or insertions. Similarly, in human 48BR and CHO K1 cells the original DNA length was rarely preserved, and the vast majority of events included deletions and insertions (>96% and 95%, respectively; Fig 2).

Spectrum of repair events of cDSB in human cells

DNA sequence analysis of 123 plasmid isolates that have resulted from the repair of cDSB in human H1299 cells revealed a wide variety of mutations types (Fig. 3). The most frequent class of events comprised very small deletions of up to 5 bp, and accounted for 43% of all repair events. Of these, 8 contained minus 2 deletions, indicating precise elimination of the 5' overhang in plasmid LP41 (line 4 in Fig. 3B). **Minus 1 deletions and full length restoration comprised 39 and 19 events, respectively (Fig. 3B). These events are the signature of the activity of a DNA polymerase(s) that skipped over the lesion and replicated the next template nucleotide, or performed full-length TLS, respectively (see below).** Another mechanism that contributed to formation of deletions is microhomology-directed end joining (MHEJ). According to this mechanism the ligation of the two ends is mediated by short homologies of 1-4 bases, and therefore the DNA is chewed until the microhomology is found (line 5 in Fig. 3B). Ten out of 13 of larger deletion events are consistent with such a mechanism. Insertions constituted 29% of all

the repair events (36/123) and could be divided into two subclasses: 1) Insertion of 50-200 bp DNA fragments originating most likely from bovine genomic DNA that is present in the serum of the cells growth medium (24 out of 36 insertion events). 2) Hybrids with pUC18 that had served as carrier DNA in the assay.

To examine whether treatment of the cells by DNA damaging agents might affect the repair of cDSB, we repeated the experiments with γ -irradiated human H1299 cells. We observed no effect of the radiation on the relative efficiency, nor the accuracy of cDSB repair (data not shown). This is consistent with the lack of effect of pre-irradiating cells on another plasmid-based repair assay used in our laboratory, which measures translesion DNA synthesis (24). It may indicate that pre-induction of the DNA damage response does not affect NHEJ of cDSB, or else that the transfection process itself, being stressful to the cells, induces the relevant genes (25).

Full-length TLS across an abasic sites near a DSB lacks template sequence instruction

Those cDSB whose repair resulted restoration of full DNA length or a -1 deletion opposite the abasic site, must have occurred via a mechanism in which the abasic site was tolerated rather than removed. Since the substrate contained a 2-nucleotides 5' overhang on one side, and a blunt end on the other side, the simplest mechanism would have been TLS by a DNA polymerase, to generate a blunt end that is sealable. A total of 102 out of 232 repair events (44%) in H1299 cells, which involved loss of one (64 events) or no (38 events) nucleotides, are likely to have occurred by such a mechanism. The DNA sequences of the 38 full-length TLS products are shown in Fig. 3 C, D. Each of the four possible nucleotides was found at the position corresponding to the abasic site, suggesting that bypass synthesis across this site occurred

without any particular specificity (Fig. 3 C, D). This is different from the preferential insertion of a dAMP opposite an abasic site observed by most DNA polymerases (the so-called A-rule; see Discussion). Surprisingly, the nucleotide next to the abasic site was highly mutated too. Only in 13 out of 38 cases (34%) was the correct nucleotide T inserted opposite the template A (Fig. 3 C, D). The majority of the insertion events at this site were mutagenic, with no strong bias for a specific nucleotide. This suggests that the polymerase(s) that perform the bypass synthesis acts in a manner that is template independent. However, the precise length synthesized (2 nt) suggests that if such a mechanism had operated, it is template length-dependent.

TLS prevents sequence losses during repair of a cDSB in various cell lines

To examine whether the TLS observed at the cDSB in H1299 cells is a general phenomenon, we analyzed DNA sequence changes caused during the repair of cDSB in several other cell lines. Table 3 presents the results of such experiments performed with the human prostate carcinoma cell line PC3, an SV40-transformed fibroblast cell line from an NBS1 patient, and the latter stably complemented with the NBS1 gene. In each cell line a significant fraction of repaired plasmids included the signature of TLS, primarily of full-length products and -1 deletions. One case of a single nucleotide insertion, which led to a net one-nucleotide increase in sequence length was observed too (Table 3), and is likely to result from TLS activity. The fraction of TLS events varied from 17% (NBS1⁺ cells) to 53% (PC3 cells), averaging at 34% for the 4 cell lines, of which about a 1/3 were full length TLS events and the rest primarily -1 deletions, consistent with skipping across the abasic site.

The repair of linear plasmids is dependent on XrccIV and altered in the absence of Ku80

In order to examine whether the repair of linear plasmids in our system depends on the known component of the NHEJ machinery, we performed the in vivo assay in *XrccIV*-deficient CHO XR1 cells, which are impaired in NHEJ, and sensitive to ionizing radiation (26,27). As a control we used the parent CHO K1 cells. As can be seen in Table 2, in K1 cells the numbers of kan^R colonies, originating from repaired linearized substrates, was similar to that obtained in other cell types, yielding a relative repair of 56% for the cDSB relative to the sDSB. In the XR1 *XrccIV*-deficient cells the results were totally different. The number of cm^R colonies was normal (hundreds/plate), however very few kan^R colonies were observed (typically 0-6) for both the sDSB or cDSB substrates, indicating that these cells were severely deficient in the repair of both sDSB and cDSB (Table 2).

We examined the repair of sDSB and cDSB also in *Ku80*-deficient CHO XRS5 cells. Unlike in the *XrccIV*-deficient cells, DSB repair did occur in the *Ku80*-deficient cells, and the relative repair of the cDSB was similar to that observed in the *Ku80*-proficient K1 cells (Table 2). DNA sequence analysis of plasmids that underwent DSB repair showed that in *Ku80*-proficient cells, most repair events at sDSB did not cause any nucleotides loss at either the protruding or the blunt ends of the break (Fig. 4A and Fig. 3s). The pattern of repair of cDSB was different: While about half of the blunt ends (53%) were preserved, most protruding ends were degraded (95% $P=0.001$; Fig. 4A and Fig. 3s), indicating that the blunt end was selectively protected compared to the complex protruding end. This selective protection was lost in the *Ku80*^{-/-} cells, where sequences were lost from both sides of the cDSB, as well as the sDSB (Fig. 4B and Fig. 3s). Analysis of the number of deletions larger than 10 bp showed a significant increase in *Ku80*^{-/-} cells compared to *Ku80*^{+/+} cells for both sDSB (from 18% to 73% $P=0.0008$)

and cDSB (from 26% to 79% $P=0.003$) (Fig. 4 and Fig. 3s). Thus, although Ku80 does not affect the relative efficiency of repair of the cDSB, it does protect its ends from nucleolytic degradation.

Deficiency in the Artemis nuclease modestly reduces the relative repair efficiency of cDSB

Since most of the cDSB repair events involved a nuclease activity, we examined whether the Artemis nuclease was involved. As can be seen in Table 2, the relative repair of cDSB was reduced by nearly twofold in Artemis-deficient cells compared to Artemis-complemented cells ($18\pm3\%$ and $35\pm10\%$ respectively; Table 2). DNA sequence analysis of linear plasmids with cDSB that underwent repair revealed similar patterns of repair events in Artemis-proficient and -deficient cells (Fig. 4s). However, the mean size of deletions larger than 2bp was 78% longer in Artemis-proficient cells compared to Artemis-deficient cells (52.4bp versus 29.4bp, respectively; Fig. 4s). Similar effects were obtained when another normal human cell (48BR) was compared to the Artemis-deficient cell line CJ179: The relative repair in the Artemis-deficient cells was twofold lower than in the normal cells (Table 2), and the average size of deletions larger than 2 bp was 58% longer in Artemis-proficient cells (42.2 bp) compared to Artemis-deficient cells (26.7 bp) (Fig. 5s).

DISCUSSION

The presence of chemical modifications near DSB may interfere with mechanisms that function to seal the break, such as NHEJ. As a result, both the efficiency and the accuracy of the process may diminish. Obviously, nucleolytic processing can in principle eliminate chemically damaged DNA regions, and generate simple and ligatable ends, however, this will cause loss of DNA

sequences near the break point. Although NHEJ is generally considered to be an error-prone repair process, mechanisms that minimize the mutagenic outcome of NHEJ **and/or its severity** may exist in mammalian cells. Indeed, several studies have shown that NHEJ is capable of aligning, annealing and patching partially complementary overhangs, thereby minimizing, or even avoiding loss of DNA sequences (6,13-16). However, the effect of chemical modifications on such processes is just beginning to emerge (28,29).

For such a purpose a plasmid assay system offers significant advantages, since it enables engineering of the DNA ends, including the incorporation of damaged nucleotides. Being a model assay system it is not expected to fully mimic the chromosomal repair of cDSB, but its relevance is indicated by its response to proteins known to be involved in chromosomal DSB repair, and its power is the ability to report on individual repair events of cDSB at a single nucleotide resolution in a large variety of mammalian cells. In the past, several predictions that were made by plasmid-based assays were later confirmed by chromosomal-based assays. For example the use of microhomology directed repair in NHEJ (15,30,31) and the end protection from degradation by the Ku complex (5,32).

DSB repair in our assay system was totally dependent on the XrccIV protein for both cDSB and sDSB. This is consistent with the critical role of the XrccIV-LigIV in NHEJ (26,27). As for Ku80, we observed similar extents of repair in Ku80-deficient and proficient cells, consistent with previous results with plasmid and chromosomal assays (5,26,32). However, degradation of the DNA ends was more extensive in Ku80-deficient cells, and the fraction of deletions longer than 10 bp significantly increased, consistent with the loss of the protection usually endowed by the Ku proteins. Interestingly, Ku80 suppressed the potentially deleterious repair-associated deletions also in cDSB repair, when accurate repair was not possible. Recently,

it was reported that a DNA ligase IV- independent alternative end joining (AEJ) pathway is robust in mammalian cells (33-35). This pathway depends to great extent on long microhomology DNA termini, and therefore relies on the sequence context. It is possible that in some sequence contexts AEJ can repair cDSB with high efficiency but with a great risk to genomic stability. Interestingly, human cells deficient in the Artemis nuclease were specifically defective in the repair of cDSB. This was associated with shorter deletions compared to Artemis-complemented cells (Fig. 4s and 5s).

The DNA substrates that we have used carried DSB that were blunt on one side, and had a short two-nucleotides 5' overhang on the other side. Such a configuration enables DNA synthesis to convert the protruding terminus into a blunt end, and subsequent ligation could lead to DSB repair with a minimal or no loss of DNA sequences (Fig. 5). In human cells most of the repair events of sDSB indeed involved template-directed DNA synthesis, clearly indicating that the repair process that preserves the original DNA length was favored over the one that causes loss of DNA sequences, consistent with previous results (30). The presence of a single abasic site in the overhang caused a significant 5-fold decrease in the efficiency of DSB repair compared to the sDSB in H1299 cells. While abasic sites inhibit DNA synthesis in at least some mammalian cells, the significant decrease in repair efficiency was surprising, since a simple nucleolytic cleavage of the problematic overhang and beyond, might have easily solved the problem, and enabled ligation. The reason for this significant effect of a single abasic site on the relative repair of the DSB is not clear, however it might hint on the presence of molecular protection against loss of sequences from the end.

With the abasic site present in the 5' overhang, 44% of the DSB repair events in H1299 cells occurred without loss of DNA sequences or with a minimal -1 deletion at the site

corresponding to the lesion. These sequences outcomes are hallmarks of TLS, representing full-length bypass, or skipping over the lesion bypass, respectively. A similar picture was observed also for the prostate cancer cell line PC3, in which 53% of DSB repair events could be attributed to TLS-assisted NHEJ, whereas in the SV40-immortalized fibroblasts examined TLS was involved in 17-21% of NHEJ repair events (Table 3). Still, in some cell lines we observed only a small fraction of TLS. This may reflect a limitation of our plasmid assay, in which the DNA ends are likely to be less protected from accidental nucleolytic degradation compared to chromosomal cDSB.

Among the TLS events, skipping over the lesion was generally more abundant than full-length TLS. This is different from TLS across an abasic site opposite a gap, where the fraction of -1 deletions was smaller than full-length bypass (36). This difference might be attributed to the substrates difference, since a higher DNA flexibility of a single-stranded overhang compared to a short ssDNA segment embedded in a duplex DNA might facilitate misalignment and skipping by the polymerase. Alternatively, this may reflect the activity of the DNA polymerase(s) involved. In most skipping events, the template A 5' to the abasic site was accurately copied. However, there were events in which skipping was accompanied by a mutation in the next nucleotide, which is generally consistent with the mutations caused at that site during TLS without skipping. Currently there are no effective methods to study the repair of chromosomal cDSB at high resolution. However, it is possible that in chromosomes, where flexibility of the DNA is more restricted, the dominant TLS event during NHEJ of cDSB will be full-length bypass. In any case, the outcome of these events is a point mutation (base substitution or -1 deletion), which is considerably less severe than long deletions.

An abasic site is an inherently miscoding lesion, and therefore in theory, any of the four nucleotides could be inserted opposite it by DNA polymerases. However, it is well documented that many prokaryotic and mammalian purified DNA polymerases tend to insert dAMP opposite abasic sites (the so-called ‘A-rule’)(37-39). Moreover, it was shown both in our laboratory and by others that dAMP is preferentially inserted opposite abasic sites in human cells (36,40), including H1299 cells used in this study (23). This implies that when operating near a cDSB, TLS in H1299 cells violates the A-rule. Interestingly, nucleotide insertion opposite the terminal nucleotide of the overhang appears to have occurred at random, without template instruction. Taken together with the events opposite the abasic site, it appears that TLS at the cDSB occurred via template-directed but sequence-independent polymerization activity. Several mammalian DNA polymerases were shown to be able to bypass an abasic site (38,39,42-45), however we do not know yet which DNA polymerase carries out the TLS step during the NHEJ events monitored in this study. A potential candidate is DNA polymerase μ , which was implicated to be involved in NHEJ, and was shown by us to possess a template-dependent, but sequence-independent activity during TLS across a synthetic basic site (46). However, an abasic site is a strongly blocking lesion, which is bypassed slowly in human cells, and likely to require more than one polymerase (39). Specifically, *in vivo* experiments have shown that pol ζ (39) as well as an aphidiclin-sensitive polymerase (22) are involved, and *in vitro* experiments have implicated also REV1 (47) and pol θ (48). The high mutability of TLS across and near an abasic site may be relevant to somatic hypermutation in the immune system, where abasic sites generated by the combined action of AID and UDG play a major role (49).

In summary, using a DNA substrate with a single abasic site on a 5’ overhang had a surprisingly big inhibitory effect on NHEJ in human cells, and uncovered a TLS-assisted NHEJ

mechanism, that allowed repair of cDSB with diminished loss of DNA sequences, albeit with the concomitant formation of point mutations. To our knowledge this is the first demonstration of TLS-assisted NHEJ in mammalian cells. Further studies are needed in order to elucidate the mechanism and regulation of this process, which might represent a damage control strategy, whereby severe deletions are prevented at the expense of less deleterious point mutations.

SUPPLEMENTARY DATA

Appended

ACKNOWLEDGEMENTS

ZL is the incumbent of the Maxwell Ellis Professorial Chair in Biomedical Research. We thank Yosef Shiloh (Tel Aviv University, Israel) for *NBS1* and complemented cells,

FUNDING

This work was supported by grants from the Flight Attendant Medical Research Institute, Florida, USA, and the Israel Science Foundation (no. 564/04 and 1136/08).

REFERENCES

1. Friedberg, E.C., Walker, G.C., Siede, W., Wood, R.D., Schultz, R.A. and Ellenberger, T. (2006) *DNA Repair and Mutagenesis*. 2nd edition ed. ASM Press, Washington DC.
2. Bassing, C.H. and Alt, F.W. (2004) The cellular response to general and programmed DNA double strand breaks. *DNA Repair (Amst)*, **3**, 781-796.

3. Lieber, M.R., Lu, H., Gu, J. and Schwarz, K. (2008) Flexibility in the order of action and in the enzymology of the nuclease, polymerases, and ligase of vertebrate non-homologous DNA end joining: relevance to cancer, aging, and the immune system. *Cell Res.*, **18**, 125-133.
4. Weterings, E. and Chen, D.J. (2008) The endless tale of non-homologous end-joining. *Cell Res.*, **18**, 114-124.
5. Liang, F. and Jasin, M. (1996) Ku80-deficient cells exhibit excess degradation of extrachromosomal DNA. *J. Biol. Chem.*, **271**, 14405-14411.
6. Ma, Y., Lu, H., Tippin, B., Goodman, M.F., Shimazaki, N., Koiwai, O., Hsieh, C.L., Schwarz, K. and Lieber, M.R. (2004) A biochemically defined system for mammalian nonhomologous DNA end joining. *Mol. Cell*, **16**, 701-713.
7. Nick McElhinny, S.A., Snowden, C.M., McCarville, J. and Ramsden, D.A. (2000) Ku recruits the XRCC4-ligase IV complex to DNA ends. *Mol. Cell. Biol.*, **20**, 2996-3003.
8. Li, Z., Otevrel, T., Gao, Y., Cheng, H.L., Seed, B., Stamato, T.D., Taccioli, G.E. and Alt, F.W. (1995) The XRCC4 gene encodes a novel protein involved in DNA double-strand break repair and V(D)J recombination. *Cell*, **83**, 1079-1089.
9. Ahnesorg, P., Smith, P. and Jackson, S.P. (2006) XLF interacts with the XRCC4-DNA ligase IV complex to promote DNA nonhomologous end-joining. *Cell*, **124**, 301-313.
10. Buck, D., Malivert, L., de Chasseval, R., Barraud, A., Fondanèche, M.C., Sanal, O., Plebani, A., Stéphan, J.L., Hufnagel, M., le Deist, F. *et al.* (2006) Cernunnos, a novel nonhomologous end-joining factor, is mutated in human immunodeficiency with microcephaly. *Cell*, **124**, 287-299.

11. Riballo, E., Kuhne, M., Rief, N., Doherty, A., Smith, G.C., Recio, M.J., Reis, C., Dahm, K., Fricke, A., Krempler, A. *et al.* (2004) A pathway of double-strand break rejoining dependent upon ATM, Artemis, and proteins locating to gamma-H2AX foci. *Mol. Cell*, **16**, 715-724.
12. Moshous, D., Callebaut, I., de Chasseval, R., Corneo, B., Cavazzana-Calvo, M., Le Deist, F., Tezcan, I., Sanal, O., Bertrand, Y., Philippe, N. *et al.* (2001) Artemis, a novel DNA double-strand break repair/V(D)J recombination protein, is mutated in human severe combined immune deficiency. *Cell*, **105**, 177-186.
13. Mahajan, K.N., Nick McElhinny, S.A., Mitchell, B.S. and Ramsden, D.A. (2002) Association of DNA polymerase mu (pol mu) with Ku and ligase IV: role for pol mu in end-joining double-strand break repair. *Mol. Cell. Biol.*, **22**, 5194-5202.
14. Lee, J.W., Blanco, L., Zhou, T., Garcia-Diaz, M., Bebenek, K., Kunkel, T.A., Wang, Z. and Povirk, L.F. (2004) Implication of DNA polymerase lambda in alignment-based gap filling for nonhomologous DNA end joining in human nuclear extracts. *J. Biol. Chem.*, **279**, 805-811.
15. Capp, J.P., Boudsocq, F., Bertrand, P., Laroche-Clary, A., Pourquier, P., Lopez, B.S., Cazaux, C., Hoffmann, J.S. and Canitrot, Y. (2006) The DNA polymerase lambda is required for the repair of non-compatible DNA double strand breaks by NHEJ in mammalian cells. *Nucleic Acids Res.*, **34**, 2998-3007.
16. Capp, J.P., Boudsocq, F., Besnard, A.G., Lopez, B.S., Cazaux, C., Hoffmann, J.S. and Canitrot, Y. (2007) Involvement of DNA polymerase mu in the repair of a specific subset of DNA double-strand breaks in mammalian cells. *Nucleic Acids Res.*, **35**, 3551-3560.

17. Ward, J.F. (2000) Complexity of damage produced by ionizing radiation. *Cold Spring Harbor Symp. Quant. Biol.*, **65**, 377-382.
18. Datta, K., Neumann, R.D. and Winters, T.A. (2005) Characterization of complex apurinic/apyrimidinic-site clustering associated with an authentic site-specific radiation-induced DNA double-strand break. *Proc. Natl. Acad. Sci. USA*, **102**, 10569-10574.
19. Datta, K., Jaruga, P., Dizdaroglu, M., Neumann, R.D. and Winters, T.A. (2006) Molecular analysis of base damage clustering associated with a site-specific radiation-induced DNA double-strand break. *Radiation Res.*, **166**, 767-781.
20. Nikjoo, H., O'Neill, P., Terrissol, M. and Goodhead, D.T. (1999) Quantitative modelling of DNA damage using Monte Carlo track structure method. *Radiat. Environ. Biophys.*, **38**, 31-38.
21. Poinsignon, C., de Chasseval, R., Soubeyrand, S., Moshous, D., Fischer, A., Hache, R.J.G. and de Villartay, J.P. (2004) Phosphorylation of Artemis following irradiation-induced DNA damage. *Eur. J. Immunol.*, **34**, 3146-3155.
22. Tauchi, H., Kobayashi, J., Morishima, K., Matsuura, S., Nakamura, A., Shiraishi, T., Ito, E., Masnada, D., Delia, D. and Komatsu, K. (2001) The forkhead-associated domain of NBS1 is essential for nuclear foci formation after irradiation but not essential for hRAD50[middle dot]hMRE11[middle dot]NBS1 complex DNA repair activity. *J Biol Chem*, **276**, 12-15.
23. Avkin, S., Adar, S., Blander, G. and Livneh, Z. (2002) Quantitative Measurement of Translesion Replication in Human Cells: Evidence for Bypass of Abasic Sites by a Replicative DNA Polymerase. *Proc. Natl. Acad. Sci. USA*, **99**, 3764-3769.

24. Avkin, S., Sevilya, Z., Toubé, L., Geacintov, N.E., Chaney, S.G., Oren, M. and Livneh, Z. (2006) p53 and p21 regulate error-prone DNA repair to yield a lower mutation load. *Mol. Cell*, **22**, 407-413.
25. Renzing, J. and Lane, D.P. (1995) p53-dependent growth arrest following calcium phosphate-mediated transfection of murine fibroblasts. *Oncogene*, **10**, 1865-1868.
26. Schulte-Uentrop, L., El-Awady, R.A., Schliecker, L., Willers, H. and Dahm-Daphi, J. (2008) Distinct roles of XRCC4 and Ku80 in non-homologous end-joining of endonuclease- and ionizing radiation-induced DNA double-strand breaks. *Nucleic Acids Res.*, **36**, 2561-2569.
27. Giaccia, A.J., MacLaren, R.A., Denko, N., Nicolaou, D. and Stamato, T.D. (1990) Increased sensitivity to killing by restriction enzymes in the XR-1 DNA double-strand break repair-deficient mutant. *Mutat. Res.*, **236**, 67-76.
28. Chen, S., Inamdar, K.V., Pfeiffer, P., Feldmann, E., Hannah, M.F., Yu, Y., Lee, J.W., Zhou, T., Lees-Miller, S.P. and Povirk, L.F. (2001) Accurate in vitro end joining of a DNA double strand break with partially cohesive 3'-overhangs and 3'-phosphoglycolate termini: effect of Ku on repair fidelity. *J Biol. Chem.*, **276**, 24323-24330.
29. Zhou, R.Z., Blanco, L., Garcia-Diaz, M., Bebenek, K., Kunkel, T.A. and Povirk, L.F. (2008) Tolerance for 8-oxoguanine but not thymine glycol in alignment-based gap filling of partially complementary double-strand break ends by DNA polymerase lambda in human nuclear extracts. *Nucleic Acids Res.*, **36**, 2895-2905.
30. Roth, D.B. and Wilson, J.H. (1986) Nonhomologous recombination in mammalian cells: role for short sequence homologies in the joining reaction. *Mol. Cell. Biol.*, **6**, 4295-4304.

31. Hazkani-Covo, E. and Covo, S. (2008) Numt-mediated double-strand break repair mitigates deletions during primate genome evolution. *PLoS Genet.*, **4**, e1000237.
32. Guirouilh-Barbat, J., Huck, S., Bertrand, P., Pirzio, L., Desmaze, C., Sabatier, L. and Lopez, B.S. (2004) Impact of the KU80 pathway on NHEJ-induced genome rearrangements in mammalian cells. *Mol. Cell*, **14**, 611-623.
33. Yan, C.T., Boboila, C., Souza, E.K., Franco, S., Hickernell, T.R., Murphy, M., Gumaste, S., Geyer, M., Zarrin, A.A., Manis, J.P. *et al.* (2007) IgH class switching and translocations use a robust non-classical end-joining pathway. *Nature*, **449**, 478-482.
34. Corneo, B., Wendland, R.L., Deriano, L., Cui, X., Klein, I.A., Wong, S.Y., Arnal, S., Holub, A.J., Weller, G.R., Pancake, B.A. *et al.* (2007) Rag mutations reveal robust alternative end joining. *Nature*, **449**, 483-486.
35. Soulas-Sprauel, P., Le Guyader, G., Rivera-Munoz, P., Abramowski, V., Olivier-Martin, C., Goujet-Zalc, C., Charneau, P. and de Villartay, J.P. (2007) Role for DNA repair factor XRCC4 in immunoglobulin class switch recombination. *J. Exp. Med.*, **204**, 1717-1727.
36. Shachar, S., Ziv, O., Avkin, S., Adar, S., Wittschieben, J., Reisner, T., Chaney, S.G., Friedberg, E.C., Wang, Z., Carell, T. *et al.* (2009) Two-polymerase mechanisms dictate error-free and error-prone translesion DNA synthesis in mammals. *EMBO J.*, **28**, 383-393.
37. Strauss, B.S. (1991) The 'A rule' of mutagen specificity: a consequence of DNA polymerase bypass of non-instructional lesions? *Bioessays*, **13**, 79-84.
38. Daube, S.S., Arad, G. and Livneh, Z. (2000) Translesion replication by DNA polymerase beta is modulated by sequence context and stimulated by fork-like flap structures in DNA. *Biochemistry*, **39**, 397-405.

39. Daube, S.S., Tomer, G. and Livneh, Z. (2000) Translesion replication by DNA polymerase delta depends on processivity accessory proteins and differs in specificity from DNA polymerase beta. *Biochemistry*, **39**, 348-355.
40. Takeshita, M. and Eisenberg, W. (1994) Mechanism of mutation on DNA templates containing synthetic abasic sites: study with double strand vector. *Nucleic Acids Res.*, **22**, 1897-1902.
42. Ohashi, E., Ogi, T., Kusumoto, R., Iwai, S., Masutani, C., Hanaoka, F. and Ohmori, H. (2000) Error-prone bypass of certain DNA lesions by the human DNA polymerase kappa. *Genes Dev.*, **14**, 1589-1594.
43. Efrati, E., Tocco, G., Eritja, R., Wilson, S.H. and Goodman, M.F. (1997) Abasic translesion synthesis by DNA polymerase beta violates the "A-rule". *J. Biol. Chem.*, **272**, 2559-2569.
44. Maga, G., Shevelev, I., Ramadan, K., Spadari, S. and Hubscher, U. (2002) DNA polymerase theta purified from human cells is a high-fidelity enzyme. *J. Mol. Biol.*, **319**, 359-369.
45. Maga, G., Villani, G., Ramadan, K., Shevelev, I., Le Gac, N.T., Blanco, L., Blanca, G., Spadari, S. and Hubscher, U. (2002) Human DNA polymerase lambda functionally and physically interacts with proliferating cell nuclear antigen in normal and translesion DNA synthesis. *J. Biol. Chem.*, **277**, 48434-48440.
46. Covo, S., Blanco, L. and Livneh, Z. (2004) Lesion bypass by human DNA polymerase mu reveals a template-dependent sequence-independent nucleotidyl transferase activity. *J. Biol. Chem.*, **279**, 859-865.

47. Lin, W., Xin, H., Zhang, Y., Wu, X., Yuan, F. and Wang, Z. (1999) The human REV1 gene codes for a DNA template-dependent dCMP transferase. *Nucleic Acids Res.*, **27**, 4468-4475.
48. Seki, M., Masutani, C., Yang, L.W., Schuffert, A., Iwai, S., Bahar, I. and Wood, R.D. (2004) High-efficiency bypass of DNA damage by human DNA polymerase Q. *EMBO J.*, **23**, 4484-4494.
49. Di Noia, J.M. and Neuberger, M.S. (2007) Molecular mechanisms of antibody somatic hypermutation. *Annu. Rev. Biochem.*, **76**, 1-22.

Table 1. Repair of complex DSB (Substrate LP41) relative to simple DSB (Substrate LP40) in human H1299 cells

Experiment no.		Number of colonies		Kan ^R /Cm ^R	Relative Repair ^a (%)
		Kan ^R	Cm ^R		
No passage ^b	LP41	0	3200	< 3x10 ⁻⁴	NA
	LP40	0	2272	< 5x10 ⁻⁴	
1	LP41	182	834	0.22	21
	LP40	1860	1782	1.04	
2	LP41	94	1448	0.065	16
	LP40	147	352	0.41	
3	LP41	359	283	1.27	20
	LP40	416	65	6.4	
4	LP41	62	100	0.62	19
	LP40	285	88	3.2	
5	LP41	233	311	0.75	17
	LP40	325	72	4.5	
6	LP41	584	896	0.65	22
	LP40	2784	962	2.9	
7	LP41	90	638	0.14	22
	LP40	361	572	0.63	
Average Relative repair:					20±2

Human H1299 cells were transfected with a mixture of linear plasmid LP41 with an abasic site in a 5' overhang, the normalizing intact plasmid pSA26, and the carrier plasmid pUC18. A parallel transfection was conducted with a mixture of the control linear plasmid LP40, which had no abasic site, along with the control and carrier plasmids. After 16 hours the plasmids were extracted using an alkaline procedure, and used to transform *E. coli recA* cells, which were then seeded in parallel on LB plates containing kanamycin (to select for repaired LP41 or LP40 plasmids) or chloramphenicol (to select for the normalizing plasmid pSA26). The repair of cDSB

relative to sDSB was calculated by dividing the kan^R/cm^R colonies ratio obtained for LP41, by the ratio obtained for LP40. See text for details.

^a Relative repair of cDSB relative to sDSB.

^b The plasmid mixture was used to directly transform the *E. coli recA* cells without prior passage through the H1299 human cells.

Table 2. Repair of complex DSB (Substrate LP41) relative to simple DSB (Substrate LP40) in hamster and human cells defective in NHEJ genes

Cell type/vector	Number of colonies (DNA amount ^b)		Relative Repair ^a %
	Kan ^R	Cm ^R	
CHO K1 (wild-type)			
LP41	203 (350)	253 (24)	56.1±7.0
LP40	393 (350)	260 (24)	
CHO XR1 (<i>XrccIV</i> ^{-/-})			
LP41	6 (350)	636 (24)	NA
LP40	2 (350)	364 (24)	
CHO XRS5 (<i>Ku80</i> ^{-/-})			
LP41	106 (200)	1228 (24)	51.0±4.9
LP40	203 (200)	1122 (24)	
Human Guetel/DA4 (Artemis ⁺)			
LP41	106 (200)	1072 (12)	35.3±9.6
LP40	278 (200)	784 (12)	
Human Guetel (Artemis ⁻)			
LP41	54 (200)	1164 (12)	18.5±3.4
LP40	242 (200)	1084 (12)	
Human 48BR (Artemis ⁺)			
LP41	28 (300)	666 (24)	15±4.9
LP40	59 (300)	205 (24)	
Human CJ179 (Artemis ⁻)			
LP41	28 (300)	237 (24)	7±2
LP40	108 (300)	80 (24)	

NHEJ assays were performed with the indicated cell lines as described in the legend to Table 1 and under Materials and Methods. The colonies numbers shown represent a typical experiment, whereas the relative repair represents the average obtained from 3-6 experiments.

^a Relative repair of cDSB relative to sDSB.

^b The numbers in the parentheses show the amount of DNA (in ng) used to transfect the mammalian cells.

Table 3. Abundance of TLS-assisted NHEJ events in various cell lines

Cell line	TLS event	DNA sequence	Occurrence
PC3	Full-length	5' CAGGTCTG <u>TT</u> ↓ATCCGGTATC	2
	Skipping	5' CAGGTCTG- <u>T</u> ATCCGGTATC	5
	Skipping	5' CAGGTCTG- <u>A</u> ATCCGGTATC	1
	Skipping	5' CAGGTCTG- <u>C</u> ATCCGGTATC	1
	Skipping	5' CAGGTCTG <u>GGG</u> ATCCGGTATC	1
	Total TLS events:		10/19 (53%)
NBS1 ⁺	Full length	5' CAGGTCTG <u>CA</u> ↓ATCCGGTATC	2
	Full length	5' CAGGTCTG <u>TT</u> ATCCGGTATC	1
	Skipping	5' CAGGTCTG- <u>T</u> ATCCGGTATC	2
	Total TLS events:		5/29 (17%)
NBS1 ⁻	Full length	5' CAGGTCTG <u>TA</u> ↓ATCCGGTATC	2
	Full length	5' CAGGTCTG <u>AT</u> ATCCGGTATC	1
	Skipping	5' CAGGTCTG- <u>T</u> ATCCGGTATC	2
	Total TLS events:		5/24 (21%)
H1299*	Full length	5' CAGGTCTG <u>NM</u> ↓ATCCGGTATC*	38
	Skipping	5' CAGGTCTG- <u>T</u> ATCCGGTATC	54?
	Skipping	5' CAGGTCTG- <u>C</u> ATCCGGTATC	10?
	Total TLS events:		102/232 (44%)

NHEJ experiments were conducted with the indicated cell lines, after which descendants of plasmids with a cDSB were isolated and subjected to DNA sequence analysis.

* The detailed sequence changes in the 38 NHEJ events associated with full length TLS (schematically presented as NM) are presented in Fig. 3C, D.

Figure Legends

Figure 1. Quantitative assay for the repair of cDSB in cultured mammalian cells. **(A)** Flow chart of the assay. Cells were transfected with a DNA mixture containing a modified linear plasmid with an abasic site on a 5'-overhang (LP41, kan^R), a normalizing intact plasmid (pSA26, cm^R) and a carrier plasmid (pUC18, amp^R). In parallel, cells were transfected with a control mixture containing a non-modified linear plasmid with a 5'-overhang but no abasic site (LP40, kan^R). Cells were incubated to allow repair, after which the plasmids were extracted and electroporated into indicator *E. coli* cells. Finally, the bacteria cells were seeded in parallel on LB plates containing kanamycin or chloramphenicol. The relative repair of the cDSB relative to the sDSB was deduced from the colonies count, as described in the text. LP40 and LP41 descendent plasmids were recovered from the bacteria colonies, and subjected to DNA sequence analysis at the vicinity of the original break point. **(B)** The break points of the linear plasmids carrying a cDSB (LP41) and a sDSB (LP40).

Figure 2. Repair of sDSB and cDSB without loss of DNA sequences. The proportion of DSB repair events with no deletions or insertions was determined for sDSB and cDSB in the indicated cell lines. In human H1299 cells full-length repair accounted for 45/50 all sequenced sDSB repair isolates and 19/123 cDSB repair isolates; in human 48BR cells the respective fractions were 20/25 and 0/25, and the Chinese hamster ovary cells CHO-K1 13/22 and 1/19. The data was obtained from experiments performed as described in the legends to Table 1. The sequences for each cell line were obtained from at least three different experiments.

Figure 3. Spectrum of mutations at repaired sites of cDSB in human H1299 cells. (A) DNA sequence analysis was performed for 123 repair events of cDSB, and the results were classified according to repair scenario. (B) Examples of DNA sequences at the repaired cDSB. The numbers in parenthesis show the occurrence of each event out of 123 repair events. (C) DNA sequence analysis of plasmids with cDSB that had been repaired without any loss of nucleotides, suggestive of a mechanism of TLS-assisted NHEJ. The number of occurrence of each sequence is presented in the parentheses. The nucleotides inserted opposite the original 5'-overhang are underlined. (D) Summary of the identity and occurrence of the nucleotides inserted opposite and 3' to the abasic site during the NHEJ of the linear plasmids with the cDSB.

Figure 4. Accuracy of sDSB and cDSB repair in *Ku80*^{+/+} and *Ku80*^{-/-} cells. Occurrence of accurate and inaccurate repair events in CHO K1 *Ku80*^{+/+} cells (A) and Xrs5 *Ku80*^{-/-} cells (B). The percentage of each type of repair event was calculated out of the total number of repair events including hybrids with pUC18, based on sequences presented in Fig. 3s.

Figure 5. Model for TLS-assisted NHEJ in mammalian cells. TLS across a DNA lesion located on a 5' overhang in one side of a cDSB leads to the formation of a blunt end. If the other side of the DSB is blunt, this will enable repair of the cDSB without loss of nucleotides. See text for details.





Figure 1
Covo et al, 2009




Figure 2
Covo et al, 2009




Figure 3
Covo et al, 2009





Figure 4
Covo et al, 2009




Figure 5
Covo et al, 2009