

# Sussex Research

## The factor structure of the twelve item General Health Questionnaire (GHQ-12): the result of negative phrasing?

Matthew Hankins

### Publication date

24-04-2008

### Licence

This work is made available under the **Copyright not evaluated** licence and should only be used in accordance with that licence. For more information on the specific terms, consult the repository record for this item.

### Citation for this work (American Psychological Association 7th edition)

Hankins, M. (2008). *The factor structure of the twelve item General Health Questionnaire (GHQ-12): the result of negative phrasing?* (Version 1). University of Sussex. <https://hdl.handle.net/10779/uos.23316950.v1>

### Published in

Clinical Practice and Epidemiology in Mental Health

### Link to external publisher version

<https://doi.org/10.1186/1745-0179-4-10>

### Copyright and reuse:

This work was downloaded from Sussex Research Open (SRO). This document is made available in line with publisher policy and may differ from the published version. Please cite the published version where possible. Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners unless otherwise stated. For more information on this work, SRO or to report an issue, you can contact the repository administrators at [sro@sussex.ac.uk](mailto:sro@sussex.ac.uk). Discover more of the University's research at <https://sussex.figshare.com/>

## The factor structure of the twelve item General Health Questionnaire (GHQ-12): the result of negative phrasing?

Article (Unspecified)

**Citation:**

Hankins, Matthew (2008) The factor structure of the twelve item General Health Questionnaire (GHQ-12): the result of negative phrasing? Clinical Practice and Epidemiology in Mental Health, 4 (10). ISSN 1745-0179

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/7495/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

**Copyright and reuse:**

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Research

Open Access

## The factor structure of the twelve item General Health Questionnaire (GHQ-12): the result of negative phrasing?

Matthew Hankins<sup>1,2,3</sup>

Address: <sup>1</sup>King's College London, Department of Psychology (at Guy's), Institute of Psychiatry, London, UK, <sup>2</sup>Department of Primary Care & Public Health, Brighton & Sussex Medical School, Brighton, UK and <sup>3</sup>Brighton & Sussex University Hospitals NHS Trust, Royal Sussex County Hospital, Brighton, UK

Email: Matthew Hankins - m.c.hankins@bsms.ac.uk

Published: 24 April 2008

Received: 4 February 2008

*Clinical Practice and Epidemiology in Mental Health* 2008, **4**:10 doi:10.1186/1745-0179-4-10

Accepted: 24 April 2008

This article is available from: <http://www.cpementalhealth.com/content/4/1/10>

© 2008 Hankins; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** The 12-item General Health Questionnaire (GHQ-12) is used routinely as a unidimensional measure of psychological morbidity. Many factor-analytic studies have reported that the GHQ-12 has two or three dimensions, threatening its validity. It is possible that these 'dimensions' are the result of the wording of the GHQ-12, namely its division into positively phrased (PP) and negatively phrased (NP) statements about mood states. Such 'method effects' introduce response bias which should be taken into account when deriving and interpreting factors.

**Methods:** GHQ-12 data were obtained from the 2004 cohort of the Health Survey for England (N = 3705). Following exploratory factor analysis (EFA), the goodness of fit indices of one, two and three factor models were compared with those of a unidimensional model specifying response bias on the NP items, using structural equation modelling (SEM). The hypotheses were (1) the variance of the responses would be significantly higher for NP items than for PP items because of response bias, and (2) that the modelling of response bias would provide the best fit for the data.

**Results:** Consistent with previous reports, EFA suggested a two-factor solution dividing the items into NP and PP items. The variance of responses to the NP items was substantially and significantly higher than for the PP items. The model incorporating response bias was the best fit for the data on all indices (RMSEA = 0.068, 90%CL = 0.064, 0.073). Analysis of the frequency of responses suggests that the response bias derives from the ambiguity of the response options for the absence of negative mood states.

**Conclusion:** The data are consistent with the GHQ-12 being a unidimensional scale with a substantial degree of response bias for the negatively phrased items. Studies that report the GHQ-12 as multidimensional without taking this response bias into account risk interpreting the artefactual factor structure as denoting 'real' constructs, committing the methodological error of reification. Although the GHQ-12 seems unidimensional as intended, the presence of such a large response bias should be taken into account in the analysis of GHQ-12 data.

## Background

The 12-item General Health Questionnaire (GHQ-12) is a self-report measure of psychological morbidity, intended to detect "psychiatric disorders...in community settings and non-psychiatric settings" [1]. It is widely used in both clinical practice [2], epidemiological research [3] and psychological research [4].

The GHQ-12 has been extensively evaluated in terms of its validity and reliability as a unidimensional index of severity of psychological morbidity [5-9]. Many studies, however, have reported that the GHQ-12 is not unidimensional, but instead assesses psychological morbidity in two or three dimensions [10-19]. Several two- and three-dimensional models have been proposed (see Martin & Newell 2005 for review [20]), and to date no study examining the factor structure of the GHQ-12 has found it to be unidimensional. These various factors have been interpreted as substantive psychological constructs such as 'Anxiety', 'Psychological distress', 'Social Dysfunction', 'Positive Health', for example. When these competing models have been compared [10,20], confirmatory factor analysis suggests that the best fitting model is the three-dimensional model of Graetz [21], which proposes that the GHQ-12 measures three distinct constructs of 'Anxiety', 'Social dysfunction' and 'Loss of confidence'.

The GHQ-12 was validated on the assumption that it was a unidimensional and generic measure of psychiatric morbidity: if it truly is multidimensional, measuring psychological functioning in three specific domains, then the validation data may be questioned, and hence the use of the GHQ-12 in clinical practice and research contexts.

Despite the apparently consistent evidence that the GHQ-12 is multifactorial, the findings of these studies could be interpreted as demonstrations of a phenomenon long-known in the psychometric literature: that measure comprising a mixture of positive and negative statements can produce an entirely artefactual division into factors [22-27]. Such artefacts, introduced as they are by the method of measurement, are known as 'method effects'.

The GHQ-12 itself comprises six items that are positive descriptions of mood states (e.g. "felt able to overcome difficulties") and six that are negative descriptions of mood states (e.g. "felt like a worthless person"). For brevity these will be referred to as 'negatively phrased items' (NP items) and 'positively phrased items' (PP items) respectively. Consistent with the hypothesis that these factors derive from method effects, a casual inspection of the two and three factor structures previously reported reveals that they comprise most (or all) of the NP items and most (or all) of the PP items. For example, of the four two factor solutions included in the review by Newell & Martin, two

comprise all of the NP items vs. all of the PP items, and two comprise all of the NP items plus one PP item vs. the remaining PP items. Of the three three factor solutions reviewed, the additional factor comprises either two NP items or two PP items, i.e. the division between NP and PP items is maintained. In addition, when reported, the correlations between dimensions are very high; this is again consistent with the hypothesis that the GHQ-12 is unidimensional, and that the apparent two- or three-dimensional structure is artefactual. Hence the question arises of whether the factors identified in these studies truly reflect clinically distinct dimensions of psychological morbidity. If the dimensions identified by these studies are simply due to the measurement bias introduced by method effects, then the constructs identified do not exist (a methodological error known as 'reification' [28]).

The analyses so far reported employed either exploratory factor analysis (EFA) to derive the factor structure or confirmatory factor analysis (CFA) to confirm the factor structure of the GHQ-12. EFA cannot in principle distinguish between genuine multidimensional structure and spurious factors generated by method effects. CFA, used appropriately, can model such method effects and so test the hypothesis that the apparent factor structure is artefactual. Unfortunately, studies employing CFA have only tested the models originally generated by EFA; used in this way, CFA is also in principle incapable of distinguishing between a genuine factor structure and an artefactual one.

The modelling of the proposed method effect is, however, relatively straightforward. Various mechanisms have been proposed for the general phenomenon of NP and PP items separating into factors, focusing on respondents' difficulties in processing negatively-phrased items. It has been suggested that negatively phrased items are more difficult to process because of inattention, variation in education, or an aversion to negative emotional content [22,24,25]. Whatever the cause, since response bias introduces variation in responses not due to variation in the measured construct, nor to random measurement error (both of which should apply equally to all items), then it may be hypothesised that response bias creates additional variation in negatively-phrased item scores, and that this additional variance is common to the negatively phrased items but not the positively phrased items. Two hypotheses follow from this: (1) NP items should have significantly higher variance than PP items and (2) a unidimensional model incorporating response bias on the NP items should fit the data better than the two- or three-factor models proposed.

The aim of this study was therefore to explore the possibility that the previously-reported factor structures of the

GHQ-12 were due to method effects, namely a response bias on the negatively phrased items.

## Methods

GHQ-12 data were obtained from the 2004 cohort of the Health Survey for England, a longitudinal general population conducted in the UK. Sampling and methodological details are in the public domain [29]. For the purposes of this study, a single adult was selected from each household to maintain independence of data. Of the 4000 such adults, 3705 provided complete data for the GHQ-12. The Likert scoring method was used, with each of the twelve items scored in the range 1 to 4, 4 indicating the most negative mood state. For clarity, the six NP items were labelled n1 to n6 and the six PP items p1 to p6.

The variances of all items were computed with 95% confidence limits, the hypothesis being that the variances on the negative questions would be uniformly greater than those on the positive questions. In addition, Pearson correlation coefficients were calculated between all items (producing a correlation matrix). Exploratory factor analysis was first conducted to explore whether the data would replicate either the two- or three-factor solutions previously reported; in the context of this study, this was a necessary step, since in the absence of a multifactorial solution there would be no method effect to explain. Consistent with most of the previous EFA analyses, the principal components method was used, with orthogonal (Varimax) rotation.

Following EFA, four models were tested for goodness of fit (CFA) using the structural equation modelling package AMOS 6.0 (maximum likelihood method):

1. Unidimensional: one factor, i.e. the intended GHQ-12 measurement model
2. Confirmatory analysis of the EFA solution
3. Three dimensional: three correlated factors, derived from the Graetz [21] model of three factors measured by 6 items ('anxiety', p1 to p6), 4 items ('social dysfunction', n1 to n4) and two items ('loss of confidence', n5 and n6)
4. Unidimensional with method effects: one factor (essentially Model 1) with correlated error terms on the NP items

Models 1 to 3 are typical of the EFA/CFA approach taken to date. Model 4 is the model hypothesised to have the best fit for the data. Path diagrams for all models are presented in Figure 1. As recommended by Byrne [30] a range of goodness of fit indices were computed for model comparison. These indices differ primarily in their treatment

of sample size, model parsimony and the range of values considered to indicate a well-fitting model. The goodness-of-fit index (GFI), adjusted goodness-of-fit index (AGFI), normed fit index (NFI) and comparative fit index (CFI) all increase towards a maximum value of 1.00 for a perfect fit, with values around 0.950 indicating a good fit for the data. In contrast, the values of the root mean square error of approximation (RMSEA) and the expected cross-validation index (ECVI) decrease with increasingly good fit, and are not limited to the range zero to one. The ECVI indicates how well the model will cross-validate to another sample, while the RMSEA provides a 'rule of thumb' cutoff for model adequacy of less than 0.08.

## Results

### Exploratory factor analysis (EFA)

Table 1 shows the Pearson correlation coefficients for all items, with correlations between similarly-phrased items highlighted in bold. Exploratory factor analysis (principal components analysis with Varimax rotation) suggested a two factor solution explaining 59.0% of the total variance in items (factor 1 eigenvalue = 3.9; factor 2 eigenvalue = 3.1). Consistent with the hypotheses of this study, the first factor comprised all of the NP items and the second factor all of the PP items. Table 2 shows the rotated component matrix for all items, illustrating the division between NP and PP items. This two-factor model was further examined by CFA, below (hypothesis 2, model 2).

### Hypothesis 1: Comparison of the variances of NP and PP items


Table 3 shows the variances of all items with 95% confidence limits (see also Figure 2 for a graphical representation of the variances of the items). It can be readily seen that the variances of the NP items (median 0.515) were uniformly higher than those of the PP items (median 0.215), with no overlap in the 95% confidence limits. This supports the first hypothesis that the NP items would have increased variances due to additional error variance caused by response bias.

### Hypothesis 2: Comparison of models

Table 4 shows the goodness-of-fit measures for all models. Of the previously reported models the best fitting was the Graetz three dimensional model; indeed, of models 1 to 3 this was the only model with an acceptable fit (RMSEA = 0.073, 90%CL (0.069, 0.077); ECVI = 0.301, 90%CL (0.274, 0.331). Model 4 was, however, superior in all fit measures to model 3 (RMSEA = 0.068, 90%CL (0.064, 0.073; ECVI = 0.214, 90%CL (0.191, 0.238)), thus confirming the second hypothesis.

### Exploration of NP response bias

Figure 3 shows frequency histograms for the six PP items. It can be seen that most of the respondents indicated the



**Figure 1**  
**Model specification (models 1 to 4).**

presence of positive mood states by responding on point 2 of the four point Likert type scale, indicating that these positive mood states were experienced with the 'Same' frequency as usual (i.e. a lack of pathology was indicated by

a PP item score of 2). In contrast, Figure 4 shows the frequency histograms for the six NP items. It can be seen that the same respondents indicated an absence of negative mood states for items n1 to n4 with a response of either 1

**Table 1: Pearson correlations between all items (PP items p1 to p6; NP items n1 to n6)**

	p2	p3	p4	p5	p6	n1	n2	n3	n4	n5	n6
p1	<b>0.43</b>	<b>0.41</b>	<b>0.47</b>	<b>0.43</b>	<b>0.39</b>	0.37	0.38	0.38	0.39	0.40	0.36
p2	-	<b>0.49</b>	<b>0.43</b>	<b>0.40</b>	<b>0.38</b>	0.27	0.25	0.34	0.33	0.38	0.39
p3	-	-	<b>0.36</b>	<b>0.47</b>	<b>0.38</b>	0.24	0.24	0.33	0.28	0.36	0.32
p4	-	-	-	<b>0.50</b>	<b>0.41</b>	0.32	0.37	0.40	0.39	0.38	0.35
p5	-	-	-	-	<b>0.48</b>	0.33	0.36	0.42	0.41	0.42	0.41
p6	-	-	-	-	-	0.39	0.40	0.42	0.51	0.48	0.49
n1	-	-	-	-	-	-	<b>0.58</b>	<b>0.54</b>	<b>0.57</b>	<b>0.51</b>	<b>0.44</b>
n2	-	-	-	-	-	-	-	<b>0.60</b>	<b>0.60</b>	<b>0.53</b>	<b>0.44</b>
n3	-	-	-	-	-	-	-	-	<b>0.60</b>	<b>0.60</b>	<b>0.54</b>
n4	-	-	-	-	-	-	-	-	-	<b>0.69</b>	<b>0.58</b>
n5	-	-	-	-	-	-	-	-	-	-	<b>0.71</b>

or 2 on the four-point scale. These correspond to responses of 'Not at all' and 'No more than usual'. Hence, respondents wishing to indicate an absence of negative mood states were forced to choose between two response options due to the ambiguous wording, and the distribution was split between the two scale points. Items n5 and n6 correspond to the extreme negative mood states of 'Losing confidence' and 'Feeling worthless', for which the response 'No more than usual' makes little sense in the absence of such mood states. Hence for these items there was less ambiguity and correspondingly more respondents opted for point 1 of the scale.

The additional variance in the NP items may therefore be explained: the ambiguous response options for the NP items shift many responses from point 2 to point 1, increasing the scale variance. The appearance of two factors in previous studies is also explained in this way, since the shift in the distribution results in the NP items being more correlated with each other than with the PP items (see Table 1: median NP inter-item correlation  $r = 0.58$ ; median PP inter-item correlation  $r = 0.43$ ; median NP inter-item correlation partialling PP items  $r_p = 0.43$ ), thus generating a spurious two factor solution comprising NP

and PP items. The Graetz three factor model may also be attributed to this mechanism, with an additional factor generated by the two least ambiguous NP items (n5 and n6 correspond to the factor of 'Losing confidence' reported by Graetz).

## Discussion


Simple exploratory factor analysis replicated previous studies, suggesting a two factor solution comprising exclusively NP and PP items. Confirmatory factor analysis, however, suggested that this model was not a good fit for the data (RMSEA = 0.086; 90% CL = 0.082, 0.090). The best fitting model was the hypothesised model with response bias on the NP items (RMSEA = 0.068; 90% CL = 0.064, 0.073). Interpreting the simple EFA solution as signalling the existence of substantive constructs (for example, factor 1 as 'depression' and factor 2 as 'social dysfunction') was therefore not warranted by the data, and would represent an example of the methodological error of reification. That the best fitting model was the model incorporating method effects suggests that the most parsimonious interpretation of the data is that the

**Table 2: Rotated component matrix for exploratory factor analysis**

Item	Factor 1	Factor 2
<b>n4:</b> Been feeling unhappy and depressed	0.82	0.25
<b>n2:</b> Felt constantly under strain	0.78	0.16
<b>n5:</b> Been losing confidence in self	0.77	0.32
<b>n1:</b> Lost sleep over worry	0.75	0.16
<b>n3:</b> Felt couldn't overcome difficulties	0.75	0.29
<b>n6:</b> Been thinking of self as worthless	0.68	0.34
<b>p3:</b> Felt capable of making decisions	0.10	0.76
<b>p2:</b> Felt playing useful part in things	0.16	0.74
<b>p5:</b> Been able to face problems	0.30	0.69
<b>p4:</b> Able to enjoy day-to-day activities	0.29	0.66
<b>p1:</b> Able to concentrate	0.31	0.64
<b>p6:</b> Been feeling reasonably happy	0.47	0.52

**Table 3: Variances (95%CL) for PP and NP items**

PP items	Variance	95%CL for variance
p5	0.15	0.14, 0.16
p3	0.16	0.16, 0.17
p1	0.19	0.19, 0.20
p6	0.24	0.23, 0.25
p2	0.25	0.24, 0.27
p4	0.26	0.25, 0.27
<b>NP items</b>		
n6	0.43	0.41, 0.45
n3	0.45	0.43, 0.47
n2	0.51	0.48, 0.53
n1	0.52	0.49, 0.54
n5	0.54	0.51, 0.56
n4	0.59	0.56, 0.61



**Figure 2**  
Item variances (PP items p1 to p6; NP items n1 to n6).

GHQ-12 is largely or wholly unidimensional, but with substantial response bias on the NP items. While this finding is in accord with the intended function of the GHQ-12 (as a unidimensional, non-specific index of psychiatric morbidity), it should be remembered that the accuracy of measurement depends on low measurement error and the absence of substantial response bias. One assumption of this study is that the random measurement error is approximately equal across all items, and that any difference in variance between NP and PP items is largely due to response bias. Although this assumption is confirmed in the analysis of the variance of the items and the pattern of inter-item correlations, it is of course also consistent with a genuine two or three factor solution in which the factors themselves have different variances. This alternative explanation is, however, even less parsimonious,


since it hypothesises not only that multiple factors represent substantive constructs, but that those constructs differ in variance along the same lines as the NP and PP items. Further, since no previous study has hypothesised in advance that (a) multiple factors exist and (b) they will differ significantly in their variance, justification of this kind would be *ex post facto*, in contrast with the *a priori* hypothesis tested in this study.

In contrast to the causal mechanisms proposed in previous studies, the response bias seems not to result from inattention, lack of education or aversion to negative phrasing, but from the ambiguous response frame used for the NP items. Respondents wishing to indicate an absence of a negative mood state are faced with two possible choices: 'Not at all' or 'No more than usual'. For


**Table 4: Fit measures for all models**

Model	GFI	AGFI	NFI	CFI	RMSEA (90%CL)	ECVI (90%CL)
1. Unidimensional	0.843	0.774	0.846	0.848	0.125 (0.121, 0.128)	0.867 (0.818, 0.917)
2. Two-dimensional	0.931	0.898	0.927	0.929	0.086 (0.082, 0.090)	0.419 (0.386, 0.454)
3. Three dimensional	0.953	0.928	0.948	0.951	0.073 (0.069, 0.077)	0.301 (0.274, 0.331)
<b>4. Unidimensional with correlated errors</b>	<b>0.969</b>	<b>0.938</b>	<b>0.965</b>	<b>0.967</b>	<b>0.068 (0.064, 0.073)</b>	<b>0.214 (0.191, 0.238)</b>





**Figure 3**  
Frequency histograms of PP item responses (items p1 to p6).



**Figure 4**  
Frequency histograms of NP item responses (items n1 to n6).

respondents not suffering from psychiatric problems, either of these options reflects their current status. This ambiguity increases the variance of the NP items and adds a correlated component to the error term for these items, thus generating spurious factors. It remains to be seen if this method effect applies in populations with a higher prevalence of psychiatric disturbance.

## Conclusion

While many studies report the factor structure of the GHQ-12 as two- or three-dimensional, it appears that these are spurious findings resulting from a neglect of method effects, specifically a response bias on the items expressing negative mood states. This in turn seems to be due to the ambiguous response options for indicating the absence of negative mood states. While the results presented here are consistent with the GHQ-12 being unidimensional as originally intended, the presence of substantial response bias on the negatively phrased items required careful attention in the analysis of clinical epidemiological data. This is especially true of the assessment of the reliability of the GHQ-12, since the computation of reliability coefficients such as Cronbach's alpha assumes no such bias, and the resulting reliability may be over- or under-estimated.

## Competing interests

The author declares that they have no competing interests.

## Authors' contributions

MH is the sole author

## References

- Goldberg DP, Williams P: *A User's Guide to the General Health Questionnaire* Windsor: nferNelson; 1988.
- Richardson A, Plant H, Moore S, Medina J, Cornwall A, Ream E: **Developing supportive care for family members of people with lung cancer: a feasibility study.** *Supportive Care in Cancer* 2007, **15**(11):1259-69.
- Henkel V, Mergl R, Kohnen R, Maier W, Möller HJ, Hegerl U: **Identifying depression in primary care: a comparison of different methods in a prospective cohort study.** *BMJ* 2003, **326**:200-201.
- Jones M, Rona RJ, Hooper R, Wessely S: **The burden of psychological symptoms in UK Armed Forces.** *Occupational Medicine* 2006, **56**(5):322-328.
- Hardy GE, Shapiro DA, Haynes CE, Rick JE: **Validation of the General Health Questionnaire-12: Using a sample of employees from England's health care services.** *Psychological Assessment* 1999, **11**(2):159-165.
- Navarro P, Ascaso C, García-Estevé L, Aguado J, Torres A, Martín-Santos R: **Postnatal psychiatric morbidity: A validation study of the GHQ-12 and the EPDS as screening tools.** *General Hospital Psychiatry* 2007, **29**(1):1-7.
- Quek KF, Low WY, Razack AH, Loh CS: **Reliability and validity of the General Health Questionnaire (GHQ-12) among urological patients: A Malaysian study.** *Psychiatry Clin Neurosci* 2001, **55**(5):509-513.
- Tait RJ, French DJ, Hulse GK: **Validity and psychometric properties of the General Health Questionnaire-12 in young Australian adolescents.** *Aust N Z J Psychiatry* 2003, **37**(3):374-381.
- Hahn D, Reuter K, Harter M: **Screening for affective and anxiety disorders in medical patients: Comparison of HADS, GHQ-12 and brief-PHQ.** *GMS Psycho-Social-Medicine* 2006, **3**:1-11.
- Shevlin M, Adamson G: **Alternative Factor Models and Factorial Invariance of the GHQ-12: A Large Sample Analysis Using Confirmatory Factor Analysis.** *Psychological Assessment* 2005, **17**(2):231-236.
- Claes R, Fraccaroli F: **The General Health Questionnaire (GHQ-12): Factorial invariance in different language versions.** *Bollettino di Psicologia Applicata* 2002, **237**:25-35.
- Pennikilampi-Kerola V, Miettinen J, Ebeling H: **A comparative assessment of the factor structures and psychometric properties of the GHQ-12 and the GHQ-20 based on data from a Finnish population-based sample.** *Scandinavian Journal of Psychology* 2006, **47**(5):431-440.
- Ip WY, Martin CR: **Factor structure of the Chinese version of the 12-item General Health Questionnaire (GHQ-12) in pregnancy.** *Journal of Reproductive and Infant Psychology* 2006, **24**(2):87-98.
- Ip WY, Martin CR: **Psychometric properties of the 12-item General Health Questionnaire (GHQ-12) in Chinese women during pregnancy and in the postnatal period.** *Psychology, Health & Medicine* 2006, **11**(1):60-69.
- Vanheule S, Bogaerts S: **Short Communication: The factorial structure of the GHQ-12.** *Stress and Health: Journal of the International Society for the Investigation of Stress* 2005, **21**(4):217-222.
- Lopez-Castedo A, Fernandez L: **Psychometric Properties of the Spanish Version of the 12-Item General Health Questionnaire in Adolescents.** *Percept Mot Skills* 2005, **100**(3 (Pt 1)):676-680.
- Kalliath TJ, O'Driscoll MP, Brough P: **A confirmatory factor analysis of the General Health Questionnaire-12.** *Stress and Health: Journal of the International Society for the Investigation of Stress* 2004, **20**(1):11-20.
- Werneke U, Goldberg DP, Yalcin I, Ustun BT: **The stability of the factor structure of the General Health Questionnaire.** *Psychological Medicine* 2000, **30**(4):823-829.
- Hu Y, Stewart-Brown S, Twigg L, Weich S: **Can the 12-item General Health Questionnaire be used to measure positive mental health?** *Psychological Medicine* 2007, **37**(7):1005-1013.
- Martin CR, Newell RJ: **Is the 12-item General Health Questionnaire (GHQ-12) confounded by scoring method in individuals with facial disfigurement?** *Psychology and Health* 2005, **20**:651-659.
- Graetz B: **Multidimensional properties of the General Health Questionnaire.** *Soc Psychiatry Psychiatr Epidemiol* 1991, **26**(3):132-138.
- Schmitt N, Stults DM: **Factors defined by negatively keyed items: The results of careless respondents?** *Applied Psychological Measurement* 1985, **9**:367-373.
- Marsh HW: **The bias of negatively worded items in rating scales for young children: A cognitive-developmental phenomenon.** *Developmental Psychology* 1986, **22**:37-49.
- Bagozzi RP: **An examination of the psychometric properties of measures of negative affect in the PANAS scales.** *Journal of Personality and Social Psychology* 1993, **65**:836-851.
- Cordery JL, Sevastos PP: **Responses to the original and revised job diagnostic survey: Is education a factor in responses to negatively worded items?** *Journal of Applied Psychology* 1993, **78**:141-143.
- Bagozzi RP, Heatherton TF: **A general approach to representing multifaceted personality constructs: application to state self-esteem.** *Structural Equation Modeling* 1994, **1**:35-67.
- Marsh HW: **Positive and negative global self-esteem: A substantively meaningful distinction or artifacts?** *J Pers Soc Psychol* 1996, **70**(4):810-819.
- Gould SJ: *The mismeasure of man* New York: Norton; 1981.
- National Centre for Social Research and University College London: Department of Epidemiology and Public Health, Health Survey for England; 2004. [computer file]. Colchester, Essex: UK Data Archive [distributor], July 2006. SN: 5439
- Byrne B: *Structural Equation Modeling With AMOS: Basic Concepts, Applications, and Programming* Lawrence Erlbaum Associates; 2001.