

# Sussex Research

## Genetic profiling in acute myeloid leukemia

Gillian Horne, John Brewin, Timothy Chevassut

#### **Publication date**

14-06-2012

#### Licence

This work is made available under the Copyright not evaluated licence and should only be used in accordance with that licence. For more information on the specific terms, consult the repository record for this item.

### Citation for this work (American Psychological Association 7th edition)

Horne, G., Brewin, J., & Chevassut, T. (2012). *Genetic profiling in acute myeloid leukemia* (Version 1). University of Sussex. https://hdl.handle.net/10779/uos.23389517.v1

#### Published in

New England Journal of Medicine

#### Link to external publisher version

https://doi.org/10.1056/NEJMc1204662

#### Copyright and reuse:

This work was downloaded from Sussex Research Open (SRO). This document is made available in line with publisher policy and may differ from the published version. Please cite the published version where possible. Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners unless otherwise stated. For more information on this work, SRO or to report an issue, you can contact the repository administrators at <a href="mailto:sro@sussex.ac.uk">sro@sussex.ac.uk</a>. Discover more of the University's research at <a href="mailto:https://sussex.figshare.com/">https://sussex.figshare.com/</a>

#### CORRESPONDENCE



# **Genetic Profiling in Acute Myeloid Leukemia**

TO THE EDITOR: In their article on the prognostic relevance of integrated genetic profiling in patients with acute myeloid leukemia (AML), Patel et al. (March 22 issue)1 propose an elaborate riskstratification system for refining prognosis for patients with intermediate-risk AML. This stratification is based on mutational analysis by DNA sequencing of 10 individual leukemia genes in addition to standard karyotyping. However, even ignoring the impracticality of such an analysis, we consider this risk stratification to be overly complicated and unjustified. Instead, on the basis of the report's supplementary data, we believe that only two genes are worthy of mutational screening, DNMT3A and MLL. Mutations in either of these genes predict adverse outcomes independent of other mutations, including internal tandem duplication in FLT3 (FLT3-ITD), as reported previously.2 Moreover, DNMT3A and MLL mutations define a biologic subgroup of AML patients typically presenting with myelomonocytic or blastic morphology and marked leukocytosis3 who may benefit from escalation of induction chemotherapy with dose-intensified daunorubicin.4 We propose that rapid identification of unfavorable mutations in DNMT3A and partial tandem duplication in MLL (MLL-PTD) alone is required for guiding optimal treatment in patients with newly diagnosed AML.

Gillian Horne, M.D. John Brewin, M.D. Timothy Chevassut

Timothy Chevassut, M.D., Ph.D.

Brighton and Sussex Medical School Brighton, United Kingdom t.chevassut@bsms.ac.uk

No potential conflict of interest relevant to this letter was reported.

1. Patel JP, Gönen M, Figueroa ME, et al. Prognostic relevance of integrated genetic profiling in acute myeloid leukemia. N Engl J Med 2012;366:1079-89.

- **2.** Ley TJ, Ding L, Walter MJ, et al. *DNMT3A* mutations in acute myeloid leukemia. N Engl J Med 2010;363:2424-33.
- **3.** Yan XJ, Xu J, Gu ZH, et al. Exome sequencing identifies somatic mutations of DNA methyltransferase gene DNMT3A in acute monocytic leukemia. Nat Genet 2011;43:309-15.
- 4. Godley LA. Profiles in leukemia. N Engl J Med 2012;366: 1152-3.

THE AUTHORS REPLY: As the correspondents suggest, the addition of diagnostic testing for mutations in DNMT3A and MLL-PTD to karyotypic evaluation would improve prognostication in patients with AML. However, restricting molecular studies to these two genes would not capture all patients with molecularly defined unfavorablerisk disease. Specifically, 41% of patients with risk that is unfavorable on the basis of our integrated genetic classification would remain in the intermediate-risk cohort without additional mutational data. We recognize that full-length resequencing of a set of informative genes in the clinical setting is a labor-intensive effort. However, with the plummeting cost of sequencing, focused nextgeneration sequencing is now a potentially viable

#### THIS WEEK'S LETTERS

- 2321 Genetic Profiling in Acute Myeloid Leukemia
- 2322 Device Closure for Stroke with Patent Foramen Ovale
- 2324 The Next GME Accreditation System
- 2325 A Man with Pain and Weakness in the Legs
- 2326 Playing Politics with the Doctor–Patient Relationship
- 2327 Central-Airway Necrosis after Stereotactic Body-Radiation Therapy

option for mutational studies. The goal of our study was to identify all genes known to be mutated in AML with potential clinical relevance when integrated with current cytogenetic evaluation and to validate their prognostic relevance. Given the increasing affordability of sequencing technology, we believe that the cost of identifying each additional patient who would benefit from more detailed mutational profiling, as compared with the more limited diagnostic eval-

uation suggested by the correspondents, is completely justifiable.

Jay P. Patel Mithat Gönen, Ph.D. Ross L. Levine, M.D.

Memorial Sloan-Kettering Cancer Center New York, NY leviner@mskcc.org

Since publication of their article, the authors report no further potential conflict of interest.

## **Device Closure for Stroke with Patent Foramen Ovale**

**TO THE EDITOR:** Furlan et al. (March 15 issue)<sup>1</sup> report no benefit from the closure of a patent foramen ovale in patients with cryptogenic stroke. Several issues deserve comment.

First, it is unclear how many study patients truly had cryptogenic stroke (30% had hypertension and 20% smoked, which suggests that some may have had lacunar strokes). It is of interest that most of the recurrent strokes (81%) were not considered to be clearly cryptogenic (including some that may have been explained by lacunar infarction in patients with risk factors). Second, among patients with shunts of moderate or substantial size, the estimated reduction in risk of stroke is 35%. The decision to include patients at low risk may have diluted the actual protective effect of the device in patients at higher risk.

Furthermore, the protocol specified the use of antiplatelet agents in the closure group, although 97 of the patients in this group had previously been taking warfarin. Was warfarin withdrawn in the closure group while being allowed in some patients in the medical-therapy group? Patent foramen ovale closure is associated with a risk of atrial fibrillation that is eight times as high as that among patients assigned to medical therapy. This issue may be critical, since double antiplatelet therapy has been shown to be inferior to warfarin in patients with atrial fibrillation.<sup>2</sup>

Carlos Guijarro, M.D., Ph.D.

Hospital Universitario Fundación Alcorcón Madrid, Spain cguijarro@fhalcorcon.es

No potential conflict of interest relevant to this letter was reported.

- 1. Furlan AJ, Reisman M, Massaro J, et al. Closure or medical therapy for cryptogenic stroke with patent foramen ovale. N Engl J Med 2012;366:991-9.
- 2. Connolly S, Pogue J, Hart R, et al. Clopidogrel plus aspirin versus oral anticoagulation for atrial fibrillation in the Atrial fibrillation Clopidogrel Trial with Irbesartan for prevention of Vascular Events (ACTIVE W): a randomised controlled trial. Lancet 2006;367:1903-12.

TO THE EDITOR: Several factors could limit the translation of CLOSURE I (Evaluation of the STARFlex Septal Closure System in Patients with a Stroke and/or Transient Ischemic Attack due to Presumed Paradoxical Embolism through a Patent Foramen Ovale) into clinical practice. Some patients with large redundant atrial septal aneurysms were excluded, although such aneurysms are an important risk factor for recurrent embolism.<sup>1</sup> The STARFlex device (NMT Medical) is no longer available; the rates of residual shunt and thrombus associated with this device were higher than those associated with other devices.2 Patients at highest risk for recurrence may have been treated outside the trial or may have received other devices. In addition, the complication rate is much higher and the success rate lower in CLOSURE I than has been previously reported.<sup>3</sup> Finally, it is an underpowered trial; thus, the possibility of clinical benefit from closure cannot be ruled out.

In our opinion, this trial demonstrates the importance of patient selection (i.e., excluding patients with alternative causes of stroke) and the need to concentrate the use of these interventions at high-volume centers with low rates of procedural complications and high success rates. Such centers can complete the ongoing randomized trials that will contribute information additional to that provided by CLOSURE I