

Sussex Research

On the convergence of finite element methods for Hamilton-Jacobi-Bellman equations

Max Jensen, Iain Smears

Publication date

01-01-2013

Licence

This work is made available under the Copyright not [evaluated](https://rightsstatements.org/page/CNE/1.0/?language=en) licence and should only be used in accordance with that licence. For more information on the specific terms, consult the repository record for this item.

Citation for this work (American Psychological Association 7th edition)

Jensen, M., & Smears, I. (2013). On the convergence of finite element methods for Hamilton-Jacobi-Bellman equations (Version 1). University of Sussex. https://hdl.handle.net/10779/uos.23396588.v1

Published in

SIAM Journal on Numerical Analysis (SINUM)

Link to external publisher version

<https://doi.org/10.1137/110856198>

Copyright and reuse:

This work was downloaded from Sussex Research Open (SRO). This document is made available in line with publisher policy and may differ from the published version. Please cite the published version where possible. Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners unless otherwise stated. For more information on this work, SRO or to report an issue, you can contact the repository administrators at [sro@sussex.ac.uk.](mailto:sro@sussex.ac.uk) Discover more of the University's research at <https://sussex.figshare.com/>

Sussex Research Online

On the convergence of finite element methods for Hamilton-Jacobi-Bellman equations

Article (Unspecified)

Jensen, Max and Smears, Iain (2013) On the convergence of finite element methods for Hamilton-Jacobi-Bellman equations. SIAM Journal on Numerical Analysis (SINUM), 51 (1). pp. 137-162. ISSN 0036-1429

This version is available from Sussex Research Online: http://sro.sussex.ac.uk/id/eprint/45507/

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

On the Convergence of Finite Element Methods for Hamilton-Jacobi-Bellman Equations

Max Jensen[∗] , Iain Smears†

November 24, 2011

Abstract

In this note we study the convergence of monotone *P*1 finite element methods on unstructured meshes for fully nonlinear Hamilton-Jacobi-Bellman equations arising from stochastic optimal control problems with possibly degenerate, isotropic diffusions. Using elliptic projection operators we treat discretisations which violate the consistency conditions of the framework by Barles and Souganidis. We obtain strong uniform convergence of the numerical solutions and, under non-degeneracy assumptions, strong L^2 convergence of the gradients.

I Introduction

Hamilton-Jacobi-Bellman (HJB) equations, which are of the form

$$
-\partial_t v + \sup_{\alpha} (L^{\alpha} v - d^{\alpha}) = 0,
$$
 (1)

where the L^{α} are linear first- or second order operators and $d^{\alpha} \in L^2$, characterise the value function of optimal control problems. Indeed, one possibility to introduce the notion of solution of (1) is via the underlying optimal control structure. An alternative approach is to use the monotonicity properties of the operator which leads to the concept of viscosity solutions. While these perceptions are essentially equivalent [17, p.72] both views have been instructive for the design and analysis of numerical methods.

The former approach, based on the discretisation of the optimal control problem before employing the Dynamic Programming Principle, has been proposed in the setting of finite elements in [26, 7, 8], see also the review article [22] and the references therein. Regarding finite difference methods we refer to the book [23]. The latter approach, which is also adopted in this note, was firmly established with the contribution [3] by Barles and Souganidis in 1991, providing an abstract framework for the convergence to viscosity solutions. Starting with [20, 21] techniques were developed to quantify the rate of convergence; more recent works are [1, 13]. A third direction was opened by the method of vanishing moments which neither enforces discrete maximum principles nor makes use of the underlying optimal control structure but relies on a higher order regularisation [16]. For a more comprehensive review of the state-of-the-art in the numerical solution of fully non-linear second order equations we refer to [15].

In the traditional finite element analysis the multiplicative testing with hat functions is viewed as the discrete analogue of the multiplicative testing procedure to define weak solutions of the (variational) differential equation. While elements of this viewpoint are implicitly used in Section VII on gradient convergence, we would like to stress a second interpretation: multiplication with hat functions as regularisation of the residual. Consider for a moment the linear problem $-a(x)\Delta u(x) = f(x)$ with smooth functions *a* and *u* as well as a hat function ϕ at the node y^{ℓ} . Let *P* be the orthogonal projection onto the approximation space with respect to the scalar product $\langle v, w \rangle = \int \nabla v \cdot \nabla w \, dx$ (given suitable boundary

[∗]Department of Mathematical Sciences, University of Durham, South Road, Durham DH1 3LE, England, m.p.j.jensen@durham.ac.uk

[†]Mathematical Institute, University of Oxford, 24-29 St. Giles', Oxford OX1 3LB, England, iain.smears@maths.ox.ac.uk

conditions). If *y* is near y^ℓ then on a fine mesh

$$
-a(y)\Delta u(y) = -\int a(y)\Delta u(y)\hat{\phi}(x) dx \approx -a(y^{\ell})\int \Delta u(x)\hat{\phi}(x) dx = a(y^{\ell})\int \nabla u(x)\cdot \nabla \hat{\phi}(x) dx = a(y^{\ell})\int \nabla Pu(x)\cdot \nabla \hat{\phi}(x) dx.
$$

since $\hat{\phi}:=\phi/\|\phi\|_{L^1(\Omega)}$ approximates a Dirac Delta as the element size is decreased. In contrast, on general meshes,

$$
-a(y)\Delta u(y) \not\approx a(y^{\ell})\int \nabla \mathfrak{I} u(x) \cdot \nabla \hat{\phi}(x) dx, \qquad (\mathfrak{I} \text{ nodal interpolant})
$$

even in the limit as the mesh is refined (see Example 1 below). This indicates that the orthogonality properties of the projection of the exact solution into the approximation space play an important role for the understanding of the (pointwise) consistency of the finite element scheme. Furthermore, this interpretation may serve as a starting point in selecting a discretisation of the HJB operator.

Our analysis combines the following key elements in a single finite element framework:

Treatment of nodally inconsistent discretisations and uniform convergence: The consistency condition (see [3, eqn.(2.4)] or [17, p.332]) of Barles and Souganidis is based on a limit involving pointwise values of smooth test functions. This condition is not satisfied by finite element methods, even for linear equations. Based on an alternative consistency condition we show the uniform convergence of finite element solutions to the viscosity solution.

Gradient convergence: We demonstrate how the coercivity of the linear operators under the supremum is recovered by the finite element method in order to control the gradient of the numerical solutions. In a uniformly elliptic setting, this leads to strong convergence in $L^2([0,T],H^1(\Omega))$.

Operators of non-negative characteristic form: The presented analysis includes the treatment of partially and fully deterministic optimal control problems, corresponding to degenerate elliptic operators under the supremum of the Hamiltonian.

Unstructured meshes: In the spirit of finite element methods the computational domain may be triangulated with an unstructured mesh, allowing to capture complex domains more easily than in a finite difference setting. Typically, weaker conditions on the mesh than quasi-uniformity can be made.

Regularisation with second order operators: We highlight that the regularisation with second order elliptic operators is sufficient to achieve convergence to the viscosity solution. Indeed, in the example of the method of artificial diffusion, we illustrate how the regularisation in the second order fully non-linear case is of the same kind and order as for first order linear operators.

Unconditional time step: Our analysis permits explicit, semi-implicit and fully implicit discretisations in time. Fully implicit discretisations in time lead to unconditionally stable schemes.

The structure of the article is as follows: In Section II we introduce a framework of finite element methods. In Section III we study the well-posedness of the discrete systems of equations and describe how these systems are solvable by a known globally convergent, locally superlinearly convergent algorithm. Section IV establishes consistency properties of elliptic projection operators. This enables us to demonstrate in Section V that the upper and lower envelopes of the numerical solutions are sub- and supersolutions. Uniform convergence to the viscosity solution is derived in Section VI and is then built upon to analyse the convergence of the gradient in Section VII. We provide a concrete specimen of a scheme belonging to our framework by describing the method of artificial diffusion in Section VIII.

II Problem statement and definition of the numerical method

Let Ω be a bounded Lipschitz domain in R *d* , *d* ≥ 2. Let *A* be a compact metric space and

$$
A \to C(\overline{\Omega}) \times C(\overline{\Omega}, \mathbb{R}^d) \times C(\overline{\Omega}) \times C(\overline{\Omega}), \ \alpha \to (a^{\alpha}, b^{\alpha}, c^{\alpha}, d^{\alpha})
$$

be continuous, such that the families of functions $\{a^a\}_{a\in A}$, $\{b^a\}_{a\in A}$, $\{c^a\}_{a\in A}$ and $\{d^a\}_{a\in A}$ are equi-continuous. Consider the bounded linear operators of non-negative characteristic form [25]

$$
L^{\alpha}: H_0^1(\Omega) \to H^{-1}(\Omega), \ w \mapsto -a^{\alpha} \Delta w + b^{\alpha} \cdot \nabla w + c^{\alpha} w
$$

where α belongs to A. Furthermore, suppose that pointwise $d^{\alpha} \ge 0$. Then

$$
\sup_{\alpha \in A} \|(a^{\alpha}, b^{\alpha}, c^{\alpha}, d^{\alpha})\|_{C(\overline{\Omega}) \times C(\overline{\Omega}, \mathbb{R}^d) \times C(\overline{\Omega}) \times C(\overline{\Omega})} < \infty, \qquad \sup_{\alpha \in A} \|L^{\alpha}\|_{C^2(\overline{\Omega}) \to C(\overline{\Omega})} < \infty.
$$
\n(2)

We assume that the final-time boundary data $v_T \in C(\overline{\Omega})$ is non-negative: $v_t \ge 0$ on $\overline{\Omega}$. For smooth *w* let

$$
Hw := \sup_{\alpha} (L^{\alpha} w - d^{\alpha}),
$$

where the supremum is applied pointwise. The HJB equation considered is

$$
-\partial_t v + Hv = 0 \qquad \text{in } (0, T) \times \Omega,
$$
 (3a)

$$
v = 0 \qquad \text{on } (0, T) \times \partial \Omega,
$$
 (3b)

$$
v = v_T \qquad \text{on } \{T\} \times \overline{\Omega}.\tag{3c}
$$

Definition 1 ([2, 17]). An upper semi-continuous (respectively lower semi-continuous) function $v : [0, T] \times \overline{\Omega} \to \mathbb{R}$ is a vis*cosity subsolution (respectively supersolution) of*

$$
-\partial_t v + Hv = 0 \tag{4}
$$

 on (0, T) × Ω *if for any w* ∈ $C^\infty(\R$ × $\R^d)$ such that v−*w has a strict local maximum (respectively minimum) at* (*t*,*x*) ∈ (0, T) × Ω with $v(t,x) = w(t,x)$, gives $-\partial_t w(t,x) + Hw(t,x) \le 0$, (respectively greater than or equal to 0). If $v \in C(\mathbb{R} \times \mathbb{R}^d)$ of equation (4) *is a viscosity subsolution and supersolution, then v is called a viscosity solution.*

Let V_i be a sequence of piecewise linear shape-regular finite element spaces with nodes y_i^{ℓ} and associated hat functions ϕ_i^{ℓ} . Here ℓ is the index ranging over the nodes of the finite element mesh. Let V_i^0 be the subspace of functions which satisfy homogeneous Dirichlet conditions. It is convenient to assume that $y_i^{\ell} \in \Omega$ for $\ell \leq N := \dim V_i^0$; i.e. the index ℓ first ranges over internal and then over external nodes. Set $\hat{\phi}^\ell_i:=\phi^\ell_i/\|\phi^\ell_i\|_{L^1(\Omega)}.$ The mesh size, i.e. the largest diameter of an element, is denoted $(\Delta x)_i$. It is assumed that $(\Delta x)_i \to 0$ as $i \to \infty$.

Let h_i be the (uniform) time step used in conjunction with V_i , with $\frac{T}{h_i} \in \mathbb{N}$, and let s_i^k be the *k*th time step at the refinement level *i*. The set of time steps is

$$
S_i := \{s_i^k : k = 0, ..., \frac{T}{h_i}\}.
$$

Let the ℓ th entry of $d_i w(s_i^k, \cdot)$ be

$$
(d_i w(s_i^k, \cdot))_\ell = \frac{w(s_i^{k+1}, y_i^\ell) - w(s_i^k, y_i^\ell)}{h_i}
$$

For each α and i find an approximate splitting $L^{\alpha} \approx E_i^{\alpha} + I_i^{\alpha}$ into linear operators

$$
\begin{split} E_i^\alpha&\colon H_0^1(\Omega)\to H^{-1}(\Omega),\;w\mapsto -\bar a_i^\alpha\,\Delta\,w+\bar b_i^\alpha\cdot\nabla\,w+\bar c_i^\alpha\,w,\\ I_i^\alpha&\colon H_0^1(\Omega)\to H^{-1}(\Omega),\;w\mapsto -\bar{\bar a}_i^\alpha\,\Delta\,w+\bar{\bar b}_i^\alpha\cdot\nabla\,w+\bar{\bar c}_i^\alpha\,w, \end{split}
$$

with continuous

$$
A \to C(\overline{\Omega}) \times C(\overline{\Omega}, \mathbb{R}^d) \times C(\overline{\Omega}), \ \alpha \to (\bar{a}_i^{\alpha}, \bar{b}_i^{\alpha}, \bar{c}_i^{\alpha}),
$$

$$
A \to C(\overline{\Omega}) \times C(\overline{\Omega}, \mathbb{R}^d) \times C(\overline{\Omega}), \ \alpha \to (\bar{a}_i^{\alpha}, \bar{b}_i^{\alpha}, \bar{c}_i^{\alpha})
$$
 (5)

.

such that \bar{c}_i^{α} and \bar{c}_i^{α} are non-negative and for some $\gamma \in \mathbb{R}$ and all $\alpha \in A$,

$$
\gamma \ge \|\bar{c}_i^{\alpha}\|_{L^{\infty}} + \|\bar{\bar{c}}_i^{\alpha}\|_{L^{\infty}}.
$$
\n
$$
(6)
$$

Also find for each *i* a non-negative d_i^{α} which approximates d^{α} : $d_i^{\alpha} \approx d^{\alpha}$. These consistency conditions $L^{\alpha} \approx E_i^{\alpha} + I_i^{\alpha}$ and $d^{\alpha} \approx d_i^{\alpha}$ are made precise as follows:

Assumption 1. *For all sequences of nodes* $(y_i^\ell)_{i\in\mathbb{N}}$ *, where in general* $\ell = \ell(i)$ *depends on i:*

$$
\lim_{i \to \infty} \sup_{\alpha \in A} \|a^{\alpha} - (\bar{a}_i^{\alpha}(y_i^{\ell}) + \bar{a}_i^{\alpha}(y_i^{\ell}))\|_{L^{\infty}(\text{supp}\,\hat{\phi}_i^{\ell})} + \|b^{\alpha} - (\bar{b}_i^{\alpha} + \bar{b}_i^{\alpha})\|_{L^{\infty}(\overline{\Omega},\mathbb{R}^d)} + \|c^{\alpha} - (\bar{c}_i^{\alpha} + \bar{c}_i^{\alpha})\|_{L^{\infty}(\overline{\Omega})} + \|d^{\alpha} - d_i^{\alpha}\|_{L^{\infty}(\overline{\Omega})} = 0.
$$

Define, for *w* ∈ *H*¹(Ω), ℓ ∈ {1,..., *N* = dim V_i^0 },

$$
(\mathsf{E}_{i}^{\alpha} w)_{\ell} := \bar{a}_{i}^{\alpha} (y_{i}^{\ell}) \langle \nabla w, \nabla \hat{\phi}_{i}^{\ell} \rangle + \langle \bar{b}_{i}^{\alpha} \cdot \nabla w + \bar{c}_{i}^{\alpha} w, \hat{\phi}_{i}^{\ell} \rangle, \tag{7a}
$$

$$
(\mathrm{I}_i^{\alpha} w)_{\ell} := \bar{\bar{a}}_i^{\alpha} (y_i^{\ell}) \langle \nabla w, \nabla \hat{\phi}_i^{\ell} \rangle + \langle \bar{\bar{b}}_i^{\alpha} \cdot \nabla w + \bar{\bar{c}}_i^{\alpha} w, \hat{\phi}_i^{\ell} \rangle, \tag{7b}
$$

$$
(\mathsf{C}_i^{\alpha})_\ell := \langle d_i^{\alpha}, \hat{\phi}_i^{\ell} \rangle. \tag{7c}
$$

On the restriction to V_i we identify the $E_i^a w$ and $I_i^a w$ with their matrix representations with respect to the nodal basis $\{\phi_i^{\ell}\}_{\ell}$. Similarly the nodal evaluation operator corresponds then to the identity matrix ld.

 ${\bf Definition \ 2.}$ $\,$ *An operator* $F:V\to \R^N$ *is said to satisfy the Local Monotonicity Property (LMP) property if for all* $v\in V_i$ *such that v has a non-positive minimum at the internal node* y_i^ℓ *,* $\ell \in \{1,\dots,N\}$ *, we have* $(Fv)_\ell \leq 0$ *. The operator F satisfies the weak Discrete Maximum Principle (wDMP) provided that:*

if
$$
(Fw)_{\ell} \ge 0
$$
 for all $\ell \in \{1, ..., N\}$, then $\min_{\Omega} w \ge \min{\{\min_{\partial \Omega} w, 0\}}$. (8)

More explicit alternative formulations of the wDMP are discussed, for example, in [5] and [6]. Note that also Id and 0 satisfy this LMP property. It is clear that if *F* satisfies the LMP and $v \in V_i$ has a *negative* minimum at the internal node y_i^ℓ then $((F + \varepsilon) \cdot \mathbf{d}) \cdot \mathbf{v}$ \in 0 for all ε > 0. This implies for all ε > 0 that $F + \varepsilon$ ld satisfies the wDMP.

 A ssumption 2. A ssume for each $\alpha \in A$ that E^α_i restricted to V_i has non-positive off-diagonal entries. Let h_i be small enough so that all h_i E $_i^a$ – Id are monotone, i.e. so that all entries of all h_i E $_i^a$ – Id are non-positive. Assume that for each α \in A that I *α i satisfies the LMP property.*

Obtain the numerical solution $v_i(T, \cdot) \in V_i$ by interpolation of v_T . Then $v_i(s_i^k, \cdot) \in V_i^0$ at time s_i^k is defined, inductively, by

$$
-d_i v_i(s_i^k, \cdot) + \sup_{\alpha} \left(E_i^{\alpha} v_i(s_i^{k+1}, \cdot) + I_i^{\alpha} v_i(s_i^k, \cdot) - C_i^{\alpha} \right) = 0. \tag{9}
$$

If all I_i^{α} are 0 then (9) is an explicit scheme, otherwise implicit. Notice that the monotonicity assumption on $h_i E_i^{\alpha}$ – ld is a time step restriction if E_i^{α} has positive diagonal entries.

III Well-posedness of the discrete HJB equations

Let $\alpha_i^{\ell,k}$ $i_i^{(\ell,\kappa)}(w)$ be a control α which maximises

$$
\sup_{\alpha} \left(E_i^{\alpha} w(s_i^{k+1}, \cdot) + I_i^{\alpha} w(s_i^k, \cdot) - C_i^{\alpha} \right)_{\ell}.
$$
\n⁽¹⁰⁾

Let $I^{k,w}_i$ k, w and $E_i^{k,w}$ $\binom{k}{i}$ be the matrices whose ℓ th row is equal to that of

$$
\mathsf{I}_{i}^{\alpha_{i}^{\ell,k}(w)} \quad \text{and} \quad \mathsf{E}_{i}^{\alpha_{i}^{\ell,k}(w)},
$$

respectively. Also let the ℓ th entry of $\zeta^{k,w}_i$ $\int_{i}^{\kappa, w}$ be

$$
\mathsf{C}_i^{\alpha_i^{\ell,k}(w)}.
$$

Thus, informally speaking, the $\mathsf{E}^{k,w}_i$ $\int_i^{k,w}$, $\int_i^{k,w}$ $\int_{i}^{k,w}$ and $C_i^{k,w}$ $a_i^{k,w}$ are gained by 'reshuffling' the rows of the E_i^{α} , I_i^{α} and C_i^{α} , respectively. Notice that the maximising control in (10) may be non-unique.

Where no ambiguity can arise we simply write I_i^w , E_i^w and C_i^w without explicitly referring to *k*. We will make use of the partial ordering of R *n* :

for
$$
x, y \in \mathbb{R}^n
$$
; $x \ge y$ if and only if $x_\ell \ge y_\ell$, $\forall \ell \in \{1, ..., n\}$.

For a collection $\{x^{\alpha}\}_{\alpha \in A} \subset \mathbb{R}^n$, we define the operator $\sup_{\alpha \in A}$ componentwise: $(\sup_{\alpha \in A} x^{\alpha})_{\ell} = \sup_{\alpha \in A} x^{\alpha}_{\ell}$. The following lemma shows that in the linear case the wDMP turns, for functions which vanish on the boundary, into an M-matrix property.

Lemma 1. The matrices $h_i \mathsf{E}_i^{k,w}$ *i*^{*k*,*w*} − Id *are monotone. The matrices of* h_i ^{*,<i>w*} $a_i^{k,w}$ + Id *restricted to* V_i^0 *are invertible diagonally dominant M-matrices for all w* \in *C*([0, *T*] × $\overline{\Omega}$). *The operators* $I^{k,w}_i$ $\int_i^{k,w}$ and h_i $\Big|_i^{k,w}$ k,w_t + Id *satisfy the LMP and wDMP, respectively*.

Proof. Monotonicity of $h_i \mathsf{E}_i^{k,u}$ $a_i^{k,w}$ – Id is a straightforward consequence of the non-positivity of the entries of $h_i \mathsf{E}_i^{\alpha}$ – Id for all $\alpha \in A$. The LMP property of \int_i^α for the node y_i^ℓ only imposes a condition on the ℓ -th row of the matrix of \int_i^α . Hence it is easily checked that the $I^{k,w}_i$ $\int_{i}^{k,w}$ and the h_i ^{$\int_{i}^{k,w}$} $a_i^{k,w}$ + ld, which are composed row-wise from the I_i^{α} and $h_iI_i^{\alpha}$ + ld, satisfy the LMP and wDMP respectively when all I_i^{α} satisfy the LMP property.

The LMP property also implies that the matrix representations of the I^α_i restricted to V^0_i are weakly diagonally dominant for all $\alpha \in A$. This is because taking $\nu = -\sum_{\ell=1}^{N} \phi_i^{\ell}$ yields

$$
0 \geq \left(\begin{matrix} \mathbf{l}_{i}^{\alpha} \boldsymbol{v} \end{matrix}\right)_{\ell} = -\left(\begin{matrix} \mathbf{l}_{i}^{\alpha} \end{matrix}\right)_{\ell \ell} - \sum_{j \neq \ell}^{N} \left(\begin{matrix} \mathbf{l}_{i}^{\alpha} \end{matrix}\right)_{\ell j},
$$

using the fact that *v* attains a non-positive minimum at each internal node. For $j \neq \ell$ the hat function ϕ^j_i $\frac{J}{i}$ attains a nonpositive minimum at y_i^{ℓ} , giving $\left(\begin{smallmatrix} 1^{\alpha} \\ i \end{smallmatrix}\right)_{\ell}$ \leq 0. This yields

$$
\left(\mathsf{I}^{\alpha}_i\right)_{\ell\ell} - \sum_{j\neq\ell}^N \left|\left(\mathsf{I}^{\alpha}_i\right)_{\ell j}\right| \geq 0.
$$

Therefore h_i ^{$\mathsf{l}^{k,u}_i$} $i^{k,w}$ + Id restricted to V_i^0 is strictly diagonally dominant and thus invertible. Furthermore, since $(h_i|_{i}^{k,w})$ $\frac{k}{i}$ ^u + Id) + *ε* ld is similarly invertible for all *ε* ≥ 0 and all off-diagonal entries are non-positive, [18, p. 114] shows that $h_i l_i^{k,w}$ $i^{k,w}$ + Id restricted to V_i^0 is represented by an invertible M-matrix. \Box

Corollary 1. *The non-linear operators* $w \mapsto I_i^{k,w}$ $\int_{i}^{k,w} w$ and $w \mapsto (h_i|_{i}^{k,w})$ $\int_{i}^{k,w}$ + ld) *w* satisfy the LMP and wDMP, respectively. More*over,* $w \mapsto -(h_i \mathsf{E}_i^{k,w})$ *^{k,<i>w*} − Id) *w is positive: if* w ≥ 0 *then* −(h_i E $_i^{k,u}$ $i^{k,w}$ – Id) $w \ge 0$.

We record a constructive proof of existence of a solution $v_i \in S_i \times V_i^0$ to (9) for all $k \in \{0, 1, 2, ..., T/h_i - 1\}$ with the below Algorithm 1. This algorithm, which can be traced back to [19], is found in the continuous setting in [24] which provides the proof of convergence and existence of solutions. In [4] it is shown that in the discrete setting it is a semi-smooth Newton method that converges superlinearly.

The algorithm to solve the non-linear problem (9) at a given time level is the following.

Algorithm 1. Given $k \in \mathbb{N}$ and $v_i(s_i^{k+1},\cdot) \in V_i^0$, choose an arbitrary $\alpha \in A$ and find $w_0 \in V_i^0$ such that

$$
(h_i I_i^{\alpha} + \mathsf{Id}) w_0 = h_i C_i^{\alpha} - (h_i E_i^{\alpha} - \mathsf{Id}) v_i (s_i^{k+1}, \cdot).
$$

For m \in {0,1,2,...}*, inductively find* $w_{m+1} \in V_i^0$ *such that*

$$
(h_i|_{i}^{w_m} + \text{Id}) w_{m+1} = h_i C_i^{w_m} - (h_i E_i^{w_m} - \text{Id}) v_i (s_i^{k+1}, \cdot). \tag{11}
$$

Theorem 1. *The numerical solution* v_i *exists, is unique, solves the linear systems*

$$
(h_i|_i^{k,\nu_i} + \text{Id})\,\nu_i(s_i^k,\cdot) = -(h_i \mathsf{E}_i^{k,\nu_i} - \text{Id})\,\nu_i(s_i^{k+1},\cdot) + h_i \mathsf{C}_i^{\nu_i} \quad \forall k \in \{0, 1, 2, \dots, \frac{T}{h_i} - 1\};\tag{12}
$$

and is non-negative. Given $k \in \{0,1,2,...\}$ and $v_i(s_i^{k+1},.) \in V_i^0$, the iterates of Algorithm 1 converge superlinearly to the unique solution $v_i(s_i^k, \cdot)$ of (9): $\lim_m w_m = v_i(s_i^k, \cdot)$. Any numerical solution v_i^α of the linear evolution problem associated *to a fixed α with homogeneous Dirichlet conditions, that is*

$$
(h_i|_i^{\alpha} + \text{Id}) v_i^{\alpha}(s_i^k, \cdot) = -(h_i \mathsf{E}_i^{\alpha} - \text{Id}) v_i^{\alpha}(s_i^{k+1}, \cdot) + h_i \mathsf{C}_i^{\alpha}, k \in \{1, 2, 3, \ldots\}, \quad with \quad v_i^{\alpha}(T, \cdot) = v_i(T, \cdot),
$$

$$
v_i(s_i^k, \cdot) \in V_i^0 \quad \text{for all } k \in \{0, 1, \ldots, \frac{T}{h_i} - 1\};
$$
 (13)

is an upper bound: $v_i \leq v_i^{\alpha}$ *on* $S_i \times \overline{\Omega}$ *.*

Figure 1: (a) illustrates a mesh that leads to a FEM discretisation of the Laplacian that is pointwise consistent with respect to the interpolant. This is no longer the case for the mesh depicted by (b) . In (b) , T_1 denotes the upper-right element.

Proof. [4, Theorem 2.1] shows existence and uniqueness of a solution $v_i(s_i^k, \cdot)$ given *k* and $v_i(s_i^{k+1}, \cdot)$ and superlinear convergence of the algorithm: their Assumption (H1) is ensured by Lemma 1 and their Assumption (H2) is guaranteed by equation (5). Existence of a solution *vⁱ* is then obtained by induction over *k*.

Also $v_i \ge 0$ on $S_i \times \overline{\Omega}$ follows from induction over *k*. By assumption $v_i(T, \cdot) \ge 0$ on $\overline{\Omega}$. Since all entries of $h_i E_i^{v_i}$ – Id are non-positive, all entries of $C_i^{\nu_i}$ are non-negative, and $\nu_i(s_i^{k+1},\cdot) \geq 0$, (12) shows

$$
(h_i|_{i}^{v_i} + \text{Id}) v_i(s_i^k, \cdot) = -(h_i \mathsf{E}_i^{v_i} - \text{Id}) v_i(s_i^{k+1}, \cdot) + h_i \mathsf{C}_i^{v_i} \ge 0.
$$

Hence by inverse positivity of $h_i|_i^{v_i}$ + ld, we deduce that $v_i(s_i^k, \cdot) \ge 0$ on $\overline{\Omega}$.

Finally, we prove that $v_i \le v_i^{\alpha}$ for all $\alpha \in A$. Fix $\alpha \in A$. Firstly, $v_i(T, \cdot) = v_i^{\alpha}(T, \cdot)$. For given $k \in \mathbb{N}$ assume that $v_i(s_i^{k+1}, \cdot) \le v_i^{k+1}$. $v_i^{\alpha}(s_i^{k+1}, \cdot)$. Then (9) implies

$$
\left(h_i\vert_{i}^{\alpha} + \mathsf{Id}\right)v_i(s_i^k, \cdot) \le h_i\mathsf{C}_i^{\alpha} - \left(h_i\mathsf{E}_i^{\alpha} - \mathsf{Id}\right)v_i(s_i^{k+1}, \cdot).
$$

Subtracting (13) from the above inequality and using monotonicity of $h_i \mathsf{E}_i^{\alpha}$ – Id yields

$$
\left(h_i\vert_{i}^\alpha+\mathsf{Id}\right)\left(v_i(s_i^k,\cdot)-v_i^\alpha(s_i^k,\cdot)\right)\leq \left(h_i\mathsf{E}^\alpha_i-\mathsf{Id}\right)\left(v_i^\alpha(s_i^{k+1},\cdot)-v_i(s_i^{k+1},\cdot)\right)\leq 0.
$$

Thus by inverse positivity of $h_i|_i^{\alpha}$ + ld we conclude that $v_i(s_i^k, \cdot) \le v_i^{\alpha}(s_i^k, \cdot)$ on Ω , which completes the induction. \Box

IV Consistency properties of elliptic projections

The Barles-Souganidis argument requires the existence of a projection operator onto the discrete function space that satisfies two properties. First, the projections of a smooth function must be convergent in a sufficiently strong sense, for example in $W^{1,\infty}$. Second, the discretisations of the partial differential operators must be pointwise consistent when applied to the projections of a smooth function, i.e. the values of the operators applied to the projections converge to the values of the continuous operator applied to the smooth function. In the context of classical finite difference methods, the interpolant to the grid satisfies these properties trivially because the operators are designed to be consistent with respect to interpolation. However, in the case of FEM, the nodal interpolant may fail to satisfy the consistency condition, even for reasonable meshes. We illustrate this behaviour in Example 1.

Example 1. *For a fixed point x in a domain, consider two sequences of meshes, such that the elements neighbouring x are as* depicted in Figure 1. Denote $\hat{\phi}_i$ and $\hat{\phi}_i$ the L 1 -normalised hat functions associated with the node x for the meshes depicted *respectively by (a) and (b). Let w be a smooth function; let* $\mathcal{I}_a w$ and $\mathcal{I}_b w$ be the nodal interpolants of w respectively on the *two meshes. We show that the mesh type of (a) leads to a FEM discretisation of the Laplacian that is strongly consistent with respect to interpolation, whereas the mesh type of (b) does not.*

For the mesh of Figure 1(a), it is well known that the FEM discretisation of the Laplacian coincides with a finite difference discretisation and that

$$
\langle \nabla \mathcal{I}_a w, \nabla \hat{\phi}_i \rangle = \frac{1}{(\Delta x)^2_i} \left(4w(x) - w(y^2_i) - w(y^3_i) - w(y^4_i) - w(y^5_i) \right) = -\Delta w(x) + O((\Delta x)^2_i).
$$

For the mesh of Figure 1(b), we sketch the calculation: first we have

$$
\left\|\varphi_i\right\|=\frac{2}{3}(\Delta x)_i^2;\quad \nabla \hat{\varphi}_i|_{T_1}=\frac{3}{2(\Delta x)_i^3}\begin{pmatrix}-1\\-1\end{pmatrix};\quad \nabla \mathcal{I}_b w|_{T_1}=\frac{1}{(\Delta x)_i}\begin{pmatrix}w(y_i^3)-w(x)\\w(y_i^2)-w(x)\end{pmatrix};
$$

thus

$$
\int_{T_1} \nabla \mathcal{I}_b w \cdot \nabla \hat{\varphi}_i dx = \frac{3}{4(\Delta x)_i^2} \left(2w(x) - w(y_i^3) - w(y_i^2) \right).
$$

Doing a similar calculation for the other elements shows that

$$
\langle \nabla \mathcal{I}_b w, \nabla \hat{\varphi}_i \rangle = \frac{3}{2(\Delta x)_i^2} \left(4 w(x) - w(y_i^2) - w(y_i^3) - w(y_i^4) - w(y_i^5) \right) = -\frac{3}{2} \Delta w(x) + O((\Delta x)_i^2).
$$

We overcome this difficulty by using a different projection operator in the Barles-Souganidis argument. Given *w* ∈ $C([0, T], H^1(\Omega))$, denote by $P_i w$ a linear mapping into $[0, T] \times V_i$ which satisfies for all $\hat{\phi}_i^{\ell} \in V_i^0$

$$
\langle \nabla P_i w(t, \cdot), \nabla \hat{\phi}_i^{\ell} \rangle = \langle \nabla w(t, \cdot), \nabla \hat{\phi}_i^{\ell} \rangle \quad \forall \, t \in [0, T]. \tag{14}
$$

Notice that P_i coincides with the classical elliptic projection of the Laplacian if $P_i w$ is chosen to interpolate w on the boundary.

Assumption 3. There are mappings P_i satisfying (14) and there is a constant C ≥ 0 such that for every $w \in C^\infty(\mathbb{R}^d)$ and *i* ∈ N*,*

$$
\|P_i w\|_{W^{1,\infty}(\Omega)} \le C \|w\|_{W^{1,\infty}(\Omega)} \quad \text{and} \quad \lim_{i \to \infty} \|P_i w - w\|_{W^{1,\infty}(\Omega)} = 0. \tag{15}
$$

The conditions under which the above assumption holds for the elliptic projection typically include a condition on the mesh grading and on the domain. In [12], it is shown that (3) holds when Ω is a bounded convex polyhedral domain in $ℝ^d$, *d* ∈ {2,3}, when the mesh satisfies a local quasi-uniformity condition and when the test functions vanish on the boundary. To apply the result for non-convex domains Ω and general $w\in C^\infty(\R\times\R^d)$, consider for example a convex polyhedral domain *B* containing Ω and assume there is a locally quasi-uniform mesh on *B* which coincides with the original mesh on Ω. Let *η* be a smooth cut-off function with compact support in *B* such that $η \equiv 1$ on $Ω$. Then the classical elliptic projection on *B*, acting on $\eta w : B \to \mathbb{R}$, has the required properties. Given this construction for P_i , it is natural to refer to it as an elliptic projection.

Lemma 2. *Let* $w \in C^{\infty}(\mathbb{R} \times \mathbb{R}^d)$ *and let* $\{s_i^{k(i)}\}$ $\{a_i^{(i)}\}_{i=1}^{\infty}$ tend to $t \in [0, T)$ *. Then*

$$
\lim_{i \to \infty} d_i P_i w(s_i^{k(i)}, \cdot) = \partial_t w(t, \cdot) \ in \ W^{1, \infty}(\Omega). \tag{16}
$$

 \Box

Proof. By linearity of P_i and (15), the result follows in the limit $i \rightarrow \infty$ from

$$
\|d_i P_i w(s_i^{k(i)}, \cdot) - \partial_t w(t, \cdot) \|_{W^{1,\infty}(\Omega)} \leq \|d_i P_i w(s_i^{k(i)}, \cdot) - d_i P_i w(t, \cdot) \|_{W^{1,\infty}(\Omega)} + \|d_i P_i w(t, \cdot) - P_i \partial_t w(t, \cdot) \|_{W^{1,\infty}(\Omega)} + \|P_i \partial_t w(t, \cdot) - \partial_t w(t, \cdot) \|_{W^{1,\infty}(\Omega)} \leq C \|d_i w(s_i^{k(i)}, \cdot) - d_i w(t, \cdot) \|_{W^{1,\infty}(\Omega)} + C \|d_i w(t, \cdot) - \partial_t w(t, \cdot) \|_{W^{1,\infty}(\Omega)} + \|P_i \partial_t w(t, \cdot) - \partial_t w(t, \cdot) \|_{W^{1,\infty}(\Omega)},
$$

where $d_i P_i w(t, \cdot) = (P_i w(t + h_i, \cdot) - P_i w(t, \cdot))/h_i$, assuming *i* is sufficiently large to ensure $t + h_i \leq T$.

Lemma 3. *Let* $w \in C^{\infty}(\mathbb{R} \times \mathbb{R}^{d})$ *and let* $\{s_i^{k(i)}\}$ $\{y_i^{(i)}\}_{i=1}^{\infty}$ tend to $t \in [0, T]$, $\{y_i^{(\ell(i))}\}_{i=1}^{\infty}$ $\int_{i}^{\ell(i)} \int_{i=1}^{\infty}$ tend to $x \in \Omega$. Then

$$
\lim_{i \to \infty} \left(\mathsf{E}_{i}^{\alpha} P_{i} w(s_{i}^{k(i)+1}, \cdot) + \mathsf{I}_{i}^{\alpha} P_{i} w(s_{i}^{k(i)}, \cdot) - \mathsf{C}_{i}^{\alpha} \right)_{\ell(i)} = L^{\alpha} w(t, x) - d^{\alpha}(x) \quad \text{uniformly over all } \alpha \in A. \tag{17}
$$

Proof. For ease of notation, the dependence of *k* and *ℓ* on *i* is made implicit. From the definition of *P*^{*i*} and integration by parts,

$$
\begin{split} &\left|\bar{\bar{a}}_{i}^{\alpha}(y_{i}^{\ell})\langle\nabla P_{i}w(s_{i}^{k},\cdot),\nabla\hat{\phi}_{i}^{\ell}\rangle+\bar{a}_{i}^{\alpha}(y_{i}^{\ell})\langle\nabla P_{i}w(s_{i}^{k+1},\cdot),\nabla\hat{\phi}_{i}^{\ell}\rangle-a^{\alpha}(y_{i}^{\ell})\langle\nabla w(t,\cdot),\nabla\hat{\phi}_{i}^{\ell}\rangle\right|\\ &=\left|\bar{\bar{a}}_{i}^{\alpha}(y_{i}^{\ell})\langle\nabla w(s_{i}^{k},\cdot),\nabla\hat{\phi}_{i}^{\ell}\rangle+\bar{a}_{i}^{\alpha}(y_{i}^{\ell})\langle\nabla w(s_{i}^{k+1},\cdot),\nabla\hat{\phi}_{i}^{\ell}\rangle-a^{\alpha}(y_{i}^{\ell})\langle\nabla w(t,\cdot),\nabla\hat{\phi}_{i}^{\ell}\rangle\right|\\ &\leq\left|\left(a^{\alpha}(y_{i}^{\ell})-\bar{\bar{a}}_{i}^{\alpha}(y_{i}^{\ell})-\bar{a}_{i}^{\alpha}(y_{i}^{\ell})\right)\langle-\Delta w(t,\cdot),\hat{\phi}_{i}^{\ell}\rangle\right|+\left|\bar{\bar{a}}_{i}^{\alpha}(y_{i}^{\ell})\langle\Delta w(t,\cdot)-\Delta w(s_{i}^{k},\cdot),\hat{\phi}_{i}^{\ell}\rangle\right|+\left|\bar{a}_{i}^{\alpha}(y_{i}^{\ell})\langle\Delta w(t,\cdot)-\Delta w(s_{i}^{k+1},\cdot),\hat{\phi}_{i}^{\ell}\rangle\right|.\end{split}
$$

Using Assumption 1 and the continuity of $w: [0, T] \mapsto C^2(\overline{\Omega})$ together with uniform boundedness of $\left\{ |\bar{a}_i^{\alpha}(y_i^{\ell})| \right\}_{\alpha \in A}$ and $\left\{\left|\bar{a}_{i}^{\alpha}(y_{i}^{\ell})\right|\right\}_{\alpha \in A}$, we conclude that

$$
\lim_{i\to\infty}\sup_{\alpha\in A}\Big|\bar{a}_i^\alpha(y_i^\ell)\langle \nabla P_i w(s_i^k,\cdot), \nabla\hat{\phi}_i^\ell\rangle+\bar{a}_i^\alpha(y_i^\ell)\langle \nabla P_i w(s_i^{k+1},\cdot), \nabla\hat{\phi}_i^\ell\rangle-a^\alpha(y_i^\ell)\langle \nabla w(t,\cdot), \nabla\hat{\phi}_i^\ell\rangle\Big|=0.
$$

Owing to the Heine-Cantor theorem for all *ε* > 0, there is a *δ* > 0 such that |∆*w*(*t*,*x*)−∆*w*(*t*, *y*)| < *ε* if |*x* − *y*| < *δ*. Since, for *i* sufficiently large, the support of $\hat{\phi}_i^{\ell}$ is contained in the ball $B(x,\delta)$ and since $\|\hat{\phi}_i^{\ell}\|_{L^1(\Omega)} = 1$ as well as $\hat{\phi}_i^{\ell} \ge 0$, we find

$$
\left|\Delta w(t,x)-\langle\Delta w(t,\cdot),\hat{\phi}_i^{\ell}\rangle\right|<\varepsilon.
$$

As $\left\{a^\alpha\right\}_{\alpha\in A}$ is an equi-continuous family of functions, we conclude that

$$
\lim_{i \to \infty} \sup_{\alpha \in A} \left| a^{\alpha}(y_i^{\ell}) \langle \Delta w(t, \cdot), \hat{\phi}_i^{\ell} \rangle - a^{\alpha}(x) \Delta w(t, x) \right| = 0;
$$

thus showing that

$$
\lim_{i \to \infty} \sup_{\alpha \in A} \left| \bar{a}_i^{\alpha}(y_i^{\ell}) \langle \nabla P_i w(s_i^k, \cdot), \nabla \hat{\phi}_i^{\ell} \rangle + \bar{a}_i^{\alpha}(y_i^{\ell}) \langle \nabla P_i w(s_i^{k+1}, \cdot), \nabla \hat{\phi}_i^{\ell} \rangle - \left(-a^{\alpha}(x) \Delta w(t, x) \right) \right| = 0. \tag{18}
$$

Using Assumption 3 and regularity of w , we see that $P_i w(s_i^k, \cdot)$ and $P_i w(s_i^{k+1}, \cdot)$ converge to $w(t, \cdot)$ in $W^{1,\infty}(\Omega)$. It can then be shown by analogous estimates and by using the equi-continuity of $\{b^\alpha\}_{\alpha\in A},$ $\{c^\alpha\}_{\alpha\in A}$ and $\{d^\alpha\}_{\alpha\in A}$, that

$$
\lim_{i \to \infty} \sup_{\alpha \in A} \left| \langle \bar{b}_i^{\alpha} \cdot \nabla P_i w(s_i^k, \cdot), \hat{\phi}_i^{\ell} \rangle + \langle \bar{b}_i^{\alpha} \cdot \nabla P_i w(s_i^{k+1}, \cdot), \hat{\phi}_i^{\ell} \rangle - b^{\alpha}(x) \cdot \nabla w(t, x) \right| = 0,
$$
\n(19a)

$$
\lim_{i \to \infty} \sup_{\alpha \in A} \left| \langle \bar{c}_i^{\alpha} P_i w(s_i^k, \cdot), \hat{\phi}_i^{\ell} \rangle + \langle \bar{c}_i^{\alpha} P_i w(s_i^{k+1}, \cdot), \hat{\phi}_i^{\ell} \rangle - c^{\alpha}(x) w(t, x) \right| = 0,
$$
\n(19b)

$$
\lim_{i \to \infty} \sup_{\alpha \in A} \left| \langle d_i^{\alpha}, \hat{\phi}_i^{\ell} \rangle - d^{\alpha}(x) \right| = 0.
$$
\n(19c)

Combining equations (18) and (19) yields (17).

V Sub- and supersolution

Set

$$
\nu^*(t,x)=\sup_{(s_i^k,y_i^\ell)\to (t,x)}\limsup_{i\to\infty}\nu_i(s_i^k,y_i^\ell),\qquad \nu_*(t,x)=\inf_{(s_i^k,y_i^\ell)\to (t,x)}\liminf_{i\to\infty}\nu_i(s_i^k,y_i^\ell)
$$

where the limit superior and limit inferior are taken over all sequences of nodes in $[0,T] \times \overline{\Omega}$ which converge to $(t, x) \in$ [0, T] $\times \overline{\Omega}$. By construction, v^* is upper and v_* lower semi-continuous. With the use of elliptic projection operators key steps of the convergence proof in [3], which is stated there in a suitable form for finite difference methods, are transferred to finite element schemes, which do not satisfy the consistency condition in [3].

Theorem 2. *The function v*[∗] *is a viscosity subsolution of* (4) *and v*∗ *is a viscosity supersolution of* (4)*.*

Proof. Step 1 (v^{} is a subsolution).* To show that *v*^{*} is a viscosity subsolution, suppose that $w \in C^\infty(\mathbb{R} \times \mathbb{R}^d)$ is a test function such that *v* [∗]−*w* has a strict local maximum at (*s*, *y*) ∈ (0,*T*)×Ω, with *v* ∗ (*s*, *y*) = *w*(*s*, *y*). Consider a neighbourhood $B := \{(t, x) \in (0, T) \times \Omega : |t - s| + |x - y| \le \delta\}$ with $\delta > 0$ such that

$$
v^*(s, y) - w(s, y) > v^*(t, x) - w(t, x) \quad \forall (t, x) \in B \setminus (s, y).
$$

 \Box

Choose *i* sufficiently large for *B* to contain nodes. Let (s_i^k, y_i^ℓ) denote the position where $v_i(s_i^k, y_i^\lambda) - P_i w(s_i^k, y_i^\lambda)$ attains the maximum among all nodes $(s_i^k, y_i^{\lambda}) \in B$. Let us pass to a subsequence $\{(s_{i(j)}^k, y_{i(j)}^\ell)\}_{j \in \mathbb{N}}$ of $\{(s_i^k, y_i^\ell)\}_{i \in \mathbb{N}}$ for which $\{v_i(s_{i(j)}^k, y_{i(j)}^\ell)\}_{j \in \mathbb{N}}$ converges to the limit superior of $\{v_i(s_i^k, y_i^\ell)\}_{i \in \mathbb{N}}$. By compactness of B, there is a subsequence of $\{(s_{i(j)}^k, y_{i(j)}^\ell)\}_{j\in\mathbb{N}}$ converging to a point $(\tilde{s}, \tilde{y}) \in B$. Then $P_i w(s_{i(j)}^k, y_{i(j)}^\ell) \to w(\tilde{s}, \tilde{y})$ from (15) and by continuity of w. As the (s_i^k, y_i^ℓ) are maximisers, one has

$$
\nu^*(\tilde{s}, \tilde{y}) - w(\tilde{s}, \tilde{y}) = \limsup_{i \to \infty} \nu_i(s_{i(j)}^k, y_{i(j)}^k) - P_i w(s_{i(j)}^k, y_{i(j)}^k) = \nu^*(s, y) - w(s, y);
$$

hence $(\tilde{s}, \tilde{y}) = (s, y)$ since (s, y) is a strict maximiser of $v^* - w$ on *B*. Thus there is a subsequence of maximising nodes converging to (s, y) to which we now pass without change of notation: $(s_i^k, y_i^\ell) \rightarrow (s, y)$. It follows that

$$
\nu_i(s_i^k, y_i^\ell) - P_i w(s_i^k, y_i^\ell) \to \nu^*(s, y) - w(s, y) = 0.
$$
\n(20)

.

Moreover, because of $(s_i^k, y_i^\ell) \to (s, y)$, the neighbours of the (s_i^k, y_i^ℓ) eventually also belong to B: For *i* sufficiently large we have $(s_i^k, y_i^{\lambda}) \in B$ if $\kappa \in \{k, k+1\}$ and $y_i^{\lambda} \in \text{supp }\hat{\phi}_i^{\ell}$; in which case

$$
v_i(s_i^k, y_i^{\lambda}) - P_i w(s_i^k, y_i^{\lambda}) \le v_i(s_i^k, y_i^{\ell}) - P_i w(s_i^k, y_i^{\ell}) \qquad \Leftrightarrow \qquad P_i w(s_i^k, y_i^{\lambda}) + \mu_i \ge v_i(s_i^k, y_i^{\lambda}),
$$

with $\mu_i = v_i(s_i^k, y_i^\ell) - P_i w(s_i^k, y_i^\ell)$. Notice that $\mu_i \to 0$ as $i \to \infty$ because of (20).

Since the matrices E_i^{α} have non-zero off diagonal entries $(E_i^{\alpha})_{\ell\lambda}$ only if y_i^{λ} ϵ supp $\hat{\phi}_i^{\ell}$, we have for all $\alpha \in A$

$$
\left((h_i \mathsf{E}_i^{\alpha} - \mathsf{Id}) \left[P_i w(s^{k+1}, \cdot) + \mu_i \right] \right)_{\ell} \leq \left((h_i \mathsf{E}_i^{\alpha} - \mathsf{Id}) v_i(s_i^{k+1}, \cdot) \right)_{\ell}.
$$

By the LMP property and linearity of I_i^{α} , since $P_i w(s_i^k, \cdot) + \mu_i - v_i(s_i^k, \cdot)$ has a non-positive minimum at y_i^{ℓ} ,

$$
\left((h_i|_i^\alpha + \mathsf{Id}) \left[P_i w(s_i^k, \cdot) + \mu_i \right] \right)_{\ell} \le \left((h_i|_i^\alpha + \mathsf{Id}) v_i(s_i^k, \cdot) \right)_{\ell}
$$

From the definition of the scheme,

$$
0 = -d_i v_i (s_i^k, y_i^\ell) + \sup_{\alpha \in A} \left(E_i^\alpha v_i (s_i^{k+1}, \cdot) + I_i^\alpha v_i (s_i^k, \cdot) - C_i^\alpha \right)_\ell
$$
 (21a)

$$
\geq -d_i \left(P_i w(s_i^k, y_i^\ell) + \mu_i \right) + \sup_{\alpha \in A} \left(E_i^\alpha \left(P_i w(s_i^{k+1}, \cdot) + \mu_i \right) + I_i^\alpha \left(P_i w(s_i^k, \cdot) + \mu_i \right) - C_i^\alpha \right)_{\ell} \tag{21b}
$$

$$
= -d_i P_i w(s_i^k, y_i^\ell) + \sup_{\alpha \in A} \left[\left(\mathsf{E}_i^{\alpha} P_i w(s_i^{k+1}, \cdot) + \mathsf{I}_i^{\alpha} P_i w(s_i^k, \cdot) - \mathsf{C}_i^{\alpha} \right)_{\ell} + \mu_i \langle \bar{c}_i^{\alpha} + \bar{c}_i^{\alpha}, \hat{\phi}_i^{\ell} \rangle \right] \tag{21c}
$$

$$
\geq -d_i P_i w(s_i^k, y_i^\ell) + \sup_{\alpha \in A} \left(\mathsf{E}_i^{\alpha} P_i w(s_i^{k+1}, \cdot) + \mathsf{I}_i^{\alpha} P_i w(s_i^k, \cdot) - \mathsf{C}_i^{\alpha} \right)_{\ell} - \gamma |\mu_i| \,. \tag{21d}
$$

Since

$$
\left| \sup_{\alpha \in A} \left(E_i^{\alpha} P_i w(s_i^{k+1} s, \cdot) + I_i^{\alpha} P_i w(s_i^k, \cdot) - C_i^{\alpha} \right)_\ell - \sup_{\alpha \in A} \left(L^{\alpha} w(s, y) - d^{\alpha}(y) \right) \right|
$$

\n
$$
\leq \sup_{\alpha \in A} \left| \left(E_i^{\alpha} P_i w(s_i^{k+1} s, \cdot) + I_i^{\alpha} P_i w(s_i^k, \cdot) - C_i^{\alpha} \right)_\ell - \left(L^{\alpha} w(s, y) - d^{\alpha}(y) \right) \right|,
$$

Lemmas 2 and 3 show that after taking the limit $i \to \infty$ in inequality (21d) and recalling that $\mu_i \to 0$, we obtain

$$
0 \ge -\partial_t w(s, y) + \sup_{\alpha \in A} \left(L^{\alpha} w(s, y) - d^{\alpha}(y) \right). \tag{22}
$$

Therefore v^* is a viscosity subsolution.

Step 2 (v∗ *is a supersolution).* Arguments similar to those above show that *v*∗ is a viscosity supersolution, where the principal changes are that one considers $w\in C^\infty(\R\times\R^d)$ such that $v_*-\overline{w}$ has a strict local minimum at some $(s,y)\in$ $(0, T) \times \Omega$ with $v_*(s, y) = w(s, y)$. Using analogous notation, inequality (21d) corresponds to

$$
0 \leq -d_i P_i w(s_i^k, y_i^\ell) + \sup_{\alpha \in A} \left(\mathsf{E}_i^{\alpha} P_i w(s_i^{k+1}, \cdot) + \mathsf{I}_i^{\alpha} P_i w(s_i^k, \cdot) - \mathsf{C}_i^{\alpha} \right)_{\ell} + \gamma \left| \mu_i \right|,
$$

i.e. there is a slight asymmetry in the argument due to the last sign in (21d). Nevertheless it is then deduced that

$$
0 \leq -\partial_t w(s, y) + \sup_{\alpha \in A} \left(L^{\alpha} w(s, y) - d^{\alpha}(y) \right).
$$

Thus *v*∗ is a viscosity supersolution.

 \Box

VI Uniform convergence

We now turn to the initial and boundary conditions. Together with the sub- and supersolution property we appeal to a comparison principle to obtain uniform convergence of the numerical solutions.

For each $\alpha \in A$, define

$$
v^{\alpha,*}(t,x) = \sup_{(s_i^k, y_i^\ell) \to (t,x)} \limsup_{i \to \infty} v_i^{\alpha}(s_i^k, y_i^\ell);
$$

where the v_i^{α} are as in (13) and the limit superior is taken over all sequences of nodes which converge to $(t, x) \in [0, T] \times \overline{\Omega}$.

Assumption 4. *Suppose that for each* $(t, x) \in [0, T] \times \partial \Omega$

$$
\inf_{\alpha \in A} v^{\alpha,*}(t, x) = 0. \tag{23}
$$

Before further considerations, let us motivate Assumption 4 with a simple example. As a side remark, this example also illustrates how in some settings Kushner-Dupuis finite difference schemes, as described in [23, 17], may be interpreted as finite element methods in the framework of this paper.

Example 2. *Consider the backward time-dependent equation in one spatial dimension*

$$
-v_t + |v_x| = 1 \quad on (0,1) \times (-1,1), \tag{24}
$$

with boundary conditions v = 0 *on* [0,1]×{−1,1} ∪{1} ×[−1,1]*. Equation* (24) *may be re-written in HJB form as*

$$
-v_t + \sup_{\alpha \in \{-1,1\}} (\alpha u_x - 1) = 0.
$$

The viscosity solution is v = min(1−*t*,1−|*x*|)*. We choose a uniform mesh with element size* 2(∆*x*)*ⁱ and we use a fully explicit discretisation, where monotonicity will be achieved by using the method of artificial diffusion, as described in [5]. Thus we have*

$$
\left(\mathsf{E}_{i}^{\alpha}w\right)_{\ell}=\varepsilon\langle\partial_{x}w,\partial_{x}\hat{\phi}_{i}^{\ell}\rangle+\alpha\langle\partial_{x}w,\hat{\phi}_{i}^{\ell}\rangle,
$$

where ε is the artificial diffusion parameter to be chosen to obtain a monotone scheme. Calculating the entries shows that the E_i^{α} *are of the form*

$$
\varepsilon \begin{pmatrix}\n\frac{2}{(\Delta x)_i^2} - \frac{1}{(\Delta x)_i^2} & & & \\
\ddots & \ddots & \ddots & & \\
& -\frac{1}{(\Delta x)_i^2} \frac{2}{(\Delta x)_i^2} - \frac{1}{(\Delta x)_i^2} & \\
& \ddots & \ddots & \ddots \\
& & -\frac{1}{(\Delta x)_i^2} \frac{2}{(\Delta x)_i^2}\n\end{pmatrix} + \alpha \begin{pmatrix}\n0 & \frac{1}{2(\Delta x)_i} & & \\
\ddots & \ddots & \ddots & \\
& -\frac{1}{2(\Delta x)_i} & 0 & \frac{1}{2(\Delta x)_i} \\
& \ddots & \ddots & \ddots \\
& & -\frac{1}{2(\Delta x)_i} & 0\n\end{pmatrix}.
$$

For monotonicity we require that all off-diagonal terms of the E_i^a *be non-positive, i.e. we require* $\varepsilon \geq (\Delta x)_i/2$ *. For example the special choice* $\varepsilon = (\Delta x)_i / 2$ *yields*

$$
E_i^1 = \begin{pmatrix} \frac{1}{(\Delta x)_i} & 0 & & & \\ \ddots & \ddots & & & \\ & -\frac{1}{(\Delta x)_i} & \frac{1}{(\Delta x)_i} & 0 & \\ & & \ddots & \ddots & \\ & & & -\frac{1}{(\Delta x)_i} & \frac{1}{(\Delta x)_i} \end{pmatrix}; \quad E_i^{-1} = \begin{pmatrix} \frac{1}{(\Delta x)_i} - \frac{1}{(\Delta x)_i} & & & \\ & \ddots & \ddots & \\ & & 0 & \frac{1}{(\Delta x)_i} - \frac{1}{(\Delta x)_i} \\ & & & \ddots & \ddots \\ & & & & 0 & \frac{1}{(\Delta x)_i} \end{pmatrix}.
$$

This is equivalent to discretising the spatial part of $- v_t + v_x$ *with backward finite differences and discretising the spatial part of* −*v^t* − *v^x with forward finite differences, as can be done in applying a Kushner-Dupuis scheme. It can then be deduced,* whilst using appropriate time steps, that v^1_i approximates the solution of

$$
-v_t + v_x = 1 \text{ on } (0,1) \times (-1,1), \qquad v = 0 \text{ on } (0,T) \times \{-1\} \cup \{1\} \times (-1,1);
$$

while v−¹ *i approximates the solution of*

$$
-v_t - v_x = 1 \text{ on } (0,1) \times (-1,1), \qquad v = 0 \text{ on } (0,T) \times \{1\} \cup \{1\} \times (-1,1).
$$

Consequently, Assumption 4 is enforced by $v^{1,*}$ *on* [0,1] × {-1} *and by* $v^{-1,*}$ *on* [0,1] × {1}*.*

Recall from Theorem 1 that

$$
0 \le v_i \le v_i^{\alpha} \quad \text{for all } \alpha \in A,
$$

and note that by construction $0 \le v_* \le v^*$. Assumption 4 thus implies that $v_*|_{[0,T] \times \partial \Omega} = v^*|_{[0,T] \times \partial \Omega} = 0$. Observe that because (23) holds in particular for all $(t, x) \in \{T\} \times \partial \Omega$, Assumption 4 implicitly enforces that the initial condition v_T vanishes on $\partial Ω$ as the v_i^{α} interpolate v_T at the final time.

Lemma 4. *The sub and super-solutions v*[∗] *and v*[∗] *satisfy*

$$
\nu^*(T, \cdot) = \nu_*(T, \cdot) = \nu_T \quad \text{ on } \overline{\Omega}.
$$
 (25)

Proof. Fix $\varepsilon > 0$ and choose a $v_T^{\varepsilon} \in C^{\infty}(\mathbb{R}^d)$ such that $v_T - 2\varepsilon \ge v_T^{\varepsilon} \ge v_T - 3\varepsilon$. Owing to Assumption 3 there is $n \in \mathbb{N}$ such that $||P_i v_T^{\varepsilon} - v_T^{\varepsilon}||_{L^{\infty}(\Omega)} \leq \varepsilon$ and $||\mathcal{I}_i v_T - v_T||_{L^{\infty}(\Omega)} \leq \varepsilon$ for all $i \geq n$. Hence, for $i \geq n$,

$$
\nu_i(T, \cdot) = \mathcal{I}_i \nu_T \ge P_i \nu_T^{\varepsilon} \ge \nu_T - 4\varepsilon. \tag{26}
$$

Recalling (14) and as $v_T^{\varepsilon} \in C^{\infty}(\mathbb{R}^d)$, it is clear that there exists $K = K(\varepsilon) \ge 0$ which bounds

$$
\left| \left(\left(\mathsf{E}_{i}^{\alpha} + \mathsf{I}_{i}^{\alpha} \right) P_{i} v_{T}^{\varepsilon} - \mathsf{C}_{i}^{\alpha} \right)_{\ell} \right| = \left| - \left(\bar{a}_{i}^{\alpha} (y_{i}^{\ell}) + \bar{a}_{i}^{\alpha} (y_{i}^{\ell}) \right) \left\langle \Delta v_{T}^{\varepsilon}, \hat{\phi}_{i}^{\ell} \right\rangle + \left\langle \left(\bar{b}_{i}^{\alpha} (y_{i}^{\ell}) + \bar{b}_{i}^{\alpha} (y_{i}^{\ell}) \right) \cdot \nabla P_{i} v_{T}^{\varepsilon} + \left(\bar{c}_{i}^{\alpha} (y_{i}^{\ell}) + \bar{c}_{i}^{\alpha} (y_{i}^{\ell}) \right) P_{i} v_{T}^{\varepsilon}, \hat{\phi}_{i}^{\ell} \right\rangle - \left(\mathsf{C}_{i}^{\alpha} \right)_{\ell} \right|
$$

for all $i \in \mathbb{N}$, $\ell \in \{1, ..., N\}$ and $\alpha \in A$. Define $w_i = P_i v_T^{\varepsilon} - K(T - t)$. To show that $v_i(s_i^k, \cdot) \ge w_i(s_i^k, \cdot)$ assume $v_i(s_i^{k+1}, \cdot) \ge w_i(s_i^k, \cdot)$ $w_i(s_i^{k+1}, \cdot)$, noting (26) for $s_i^{k+1} = T$. Fix an *i* and ℓ and let $\alpha = \alpha_i^{k,\ell}$ $i^{k,\ell}(v_i)$ as for (10). From

$$
-d_i w_i(s_i^k, y_i^{\ell}) + \left(\mathsf{E}_i^{\alpha} w_i(s_i^{k+1}, \cdot) + \mathsf{I}_i^{\alpha} w_i(s_i^{k+1}, \cdot)\right)_{\ell} = -K + \left(\left(\mathsf{E}_i^{\alpha} + \mathsf{I}_i^{\alpha}\right) P_i v_{T}^{\epsilon}\right)_{\ell} - K(T - s_i^{k+1}) \langle \bar{c}_i^{\alpha}, \hat{\phi}_i^{\ell} \rangle - K(T - s_i^{k}) \langle \bar{c}_i^{\alpha}, \hat{\phi}_i^{\ell} \rangle
$$

$$
\leq \left(\mathsf{C}_i^{\alpha}\right)_{\ell} \stackrel{(12)}{=} -d_i v_i(s_i^k, y_i^{\ell}) + \left(\mathsf{E}_i^{v_i} v_i(s_i^{k+1}, \cdot) + \mathsf{I}_i^{v_i} v_i(s_i^{k+1}, \cdot)\right)_{\ell}
$$

we may deduce that

$$
\left(\left(h_i\big|_{i}^{v_i} + \mathsf{Id}\right)\left[v_i(s_i^k,\cdot) - w_i(s_i^k,\cdot)\right]\right)_{\ell} \ge \left(\left(h_i\mathsf{E}_i^{v_i} - \mathsf{Id}\right)\left[v_i(s_i^k,\cdot) - w_i(s_i^k,\cdot)\right]\right)_{\ell} \ge 0.
$$

Note that $v_i(s_i^k, \cdot) \in V_i^0$ vanishes on $\partial \Omega$ and $w_i(s_i^k, \cdot) \le 0$ on $\partial \Omega$. Thus Lemma 1 and (8) imply $v_i(s_i^k, \cdot) \ge w_i(s_i^k, \cdot)$ on $\overline{\Omega}$. Because K is independent of i and $P_i v_T^{\varepsilon} \to v_T^{\varepsilon}$ as $i \to \infty$, we have for any sequence $(s_i^k, y_i^{\ell}) \to (T, x)$, $x \in \Omega$,

$$
\liminf_{i \to \infty} \nu_i(s_i^k, y_i^\ell) \ge \liminf_{i \to \infty} \nu_i(s_i^k, y_i^\ell) \ge \nu_T(x) - 4\varepsilon.
$$

So $v_*(T, \cdot) \ge v_T - 4\varepsilon$. Since ε was arbitrary, $v_*(T, \cdot) \ge v_T$. The argument for showing that $v^* \le v_T$ is analogous with $w_i = P_i v_T^{\varepsilon} + K(T - t)$ and $v_T + 2\varepsilon \le v_T^{\varepsilon} \le v_T + 3\varepsilon$. To conclude, $v_T \le v_*(T, \cdot) \le v^*(T, \cdot) \le v_T$, which proves (25). \Box

The proof of Lemma 4 is related to the arguments in [17, p. 335]. In the next assumption we draw upon one of the building blocks of the theory of viscosity solutions, namely the extension of classical comparison principles to spaces of semi-continuous functions, cf. [10, Sec. 5] and [17, p. 219].

Assumption 5. Let \overline{v} be a lower semi-continuous supersolution with $\overline{v}|_{[0,T]\times\partial\Omega} = 0$ and $\overline{v}(T,\cdot) = v_T$. Similarly, let <u>v</u> be an *upper semi-continuous subsolution with* $v|_{[0,T]\times\Omega} = 0$ *and* $v(T, \cdot) = v_T$. Then $v \leq \overline{v}$.

Let $t = \partial s_i^k + (1-\theta)s_i^{k+1} \in [s_i^k, s_i^{k+1}]$ lie between two time steps, $\theta \in [0,1]$. Then we interpret $v_i(t, \cdot)$ as the linear interpolant between $v_i(s_i^k, \cdot)$ and $v_i(s_i^{k+1}, \cdot)$:

$$
v_i(t, \cdot) = \partial v_i(s_i^k, \cdot) + (1 - \partial) v_i(s_i^{k+1}, \cdot). \tag{27}
$$

Theorem 3. One has $v_* = v^* = v$, where v is the unique viscosity solution of equation (4) with $v(T, \cdot) = v_T$ and $v|_{[0,T] \times \partial \Omega} = 0$. *Furthermore*

$$
\lim_{i \to \infty} ||v_i - v||_{L^{\infty}((0, T) \times \Omega)} = 0.
$$
\n(28)

Proof. The previous assumption implies that $v_* \ge v^*$ thus $v^* = v_* = v$. Select for each $i \in \mathbb{N}$ a point $(t_i, x_i) \in [0, T] \times \overline{\Omega}$ such that

$$
||v_i - v||_{L^{\infty}((0,T)\times\Omega)} = |v_i - v|(t_i, x_i).
$$

Such (*tⁱ* ,*xi*) exist as *vⁱ* − *v* is a continuous function on a compact domain. Let *xⁱ* belong to (the closure of) the element *T* of the finite element mesh and $t \in [s_i^k, s_i^{k+1}]$; then $v_i(t_i, x_i)$ is a weighted average of the values of v_i at the corners of the slab $[s_i^k, s_i^{k+1}] \times \overline{T}$. Thus there is a corner (s_i^k, y_i^ℓ) of the slab such that

$$
||v_i - v||_{L^{\infty}((0,T)\times\Omega)} \leq |v_i(s_i^k, y_i^{\ell}) - v(t_i, x_i)|.
$$

If (28) was wrong we could select a subsequence and an $\varepsilon > 0$ such that

$$
\liminf_{j\to\infty} |v_{i(j)}(s_{i(j)}^k, y_{i(j)}^\ell) - v(t_{i(j)}, x_{i(j)})| \geq \varepsilon.
$$

By possibly passing to a further subsequence we may assume that $\{(t_{i(j)},x_{i(j)})\}_j$ converges to an $(t,x)\in[0,T]\times\overline{\Omega}.$ However, this contradicts

$$
\nu(t,x) = \nu_*(t,x) \le \liminf_{j \to \infty} \nu_{i(j)}(s_{i(j)}^k, y_{i(j)}^\ell) \le \limsup_{i \to \infty} \nu_{i(j)}(s_{i(j)}^k, y_{i(j)}^\ell) \le \nu^*(t,x) = \nu(t,x).
$$

Thus (28) holds.

VII Gradient convergence

For shorthand, let $W = W^{1,\infty}((0,T) \times \Omega)$. It is convenient to introduce the discrete spaces

 $W_i := \{ v \in C([0, T], V_i^0) : v|_{[s_i^k, s_i^{k+1}] \times \Omega} \text{ is affine in time} \},$

which means that functions in W_i have between two time-steps the form of (27). Observe that $W_i \subset W$ for all $i \in \mathbb{N}$.

Fix an arbitrary $\alpha \in A$. It is convenient to view E^{α}_i and I^{α}_i as bilinear forms on $H^1(\Omega) \times V_i$. Functions $u \in V_i$ have the nodal representation

$$
u(y) = \sum_{\ell} u(y_i^{\ell}) \phi_i^{\ell}(y).
$$

To test with functions other than $\hat{\phi}^\ell_i$ we introduce the following bilinear form as a partially discrete pivot: for $w\in H^1(\Omega)$ and $u \in V_i$

$$
\langle \mathsf{E}_i^{\alpha} w, u \rangle \mathrel{\mathop:}= \sum_{\ell} u(y_i^{\ell}) \big(\bar{a}_i^{\alpha} (y_i^{\ell}) \langle \nabla w, \nabla \phi_i^{\ell} \rangle + \langle \bar{b}_i^{\alpha} \cdot \nabla w + \bar{c}_i^{\alpha} w, \phi_i^{\ell} \rangle \big).
$$

We use corresponding interpretation for $\langle \mathcal{U}_i^{\alpha} w, u \rangle$ and also

$$
\langle\!\langle w,u\rangle\!\rangle = \langle\!\langle \mathsf{Id}\,w,u\rangle\!\rangle = \sum_{\ell} w(y_i^{\ell})\,u(y_i^{\ell})\|\phi_i^{\ell}\|_{L^1(\Omega)} \qquad \text{and} \qquad \langle\!\langle \mathsf{C}_i^{\alpha},u\rangle\!\rangle = \sum_{\ell} u(y_i^{\ell})\,\langle d_i^{\alpha},\phi_i^{\ell}\rangle = \langle d_i^{\alpha},u\rangle.
$$

Assume that for the chosen *α*:

$$
|w|_{L^{2}([0,T],H^{1}(\Omega))}^{2} \lesssim \sum_{k=0}^{(T/h_{i})-1} \Biggl(\langle\!\langle (h_{i} \mathsf{E}_{i}^{\alpha} - \mathsf{Id}) w(s_{i}^{k+1}, \cdot) + (h_{i} I_{i}^{\alpha} + \mathsf{Id}) w(s_{i}^{k}, \cdot), w(s_{i}^{k}, \cdot) \rangle\!\rangle \Biggr) + \frac{1}{2} \langle\!\langle w(T, \cdot), w(T, \cdot) \rangle + \|w(T, \cdot)\|_{H^{1}(\Omega)}^{2}
$$
\n
$$
\stackrel{\left(\frac{\ast}{2}\right)}{=} \sum_{k=0}^{(T/h_{i})-1} \Biggl(h_{i} \langle\!\langle \mathsf{E}_{i}^{\alpha} w(s_{i}^{k+1}, \cdot) + \mathsf{I}_{i}^{\alpha} w(s_{i}^{k}, \cdot), w(s_{i}^{k}, \cdot) \rangle\!\rangle + \frac{1}{2} \langle\!\langle w(s_{i}^{k+1}, \cdot) - w(s_{i}^{k}, \cdot), w(s_{i}^{k+1}, \cdot) - w(s_{i}^{k}, \cdot) \rangle\!\rangle \Biggr) \tag{29}
$$
\n
$$
+ \frac{1}{2} \langle\!\langle w(0, \cdot), w(0, \cdot) \rangle\!\rangle + \|w(T, \cdot)\|_{H^{1}(\Omega)}^{2}
$$

 \Box

for all $w \in W_i$ with $w \ge 0$ and $i \in \mathbb{N}$, where (*) is a simple reformulation in terms of a telescope sum.

Due to the definition of the numerical method and the non-negativity of the *vⁱ* ,

$$
\begin{split} |v_{i}|_{L^{2}([0,T],H^{1}(\Omega))}^{2} &\lesssim \sum_{k=0}^{(T/h_{i})-1} \Biggl(\langle\!\!\!\!\langle \left(h_{i}\mathsf{E}_{i}^{\alpha}-\mathsf{Id} \right) v_{i}(s_{i}^{k+1},\cdot) + \left(h_{i}\right|_{i}^{\alpha}+\mathsf{Id} \right) v_{i}(s_{i}^{k},\cdot),\\ &v_{i}(s_{i}^{k},\cdot) \rangle\!\!\!\rangle \Biggr) + \frac{1}{2} \langle\!\!\!\langle v_{i}(T,\cdot),v_{i}(T,\cdot) \rangle\!\!\!\rangle + \|v_{i}(T,\cdot)\|_{H^{1}(\Omega)}^{2} \\ &\leq \sum_{k=0}^{(T/h_{i})-1} \langle\!\!\!\langle h_{i}\mathsf{C}_{i}^{\alpha},v_{i}(s_{i}^{k},\cdot) \rangle\!\!\!\rangle + \frac{1}{2} \langle\!\!\!\langle v_{i}(T,\cdot),v_{i}(T,\cdot) \rangle\!\!\!\rangle + \|v_{i}(T,\cdot)\|_{H^{1}(\Omega)}^{2} \\ &\lesssim T \|d_{i}^{\alpha}\|_{L^{1}(\Omega)} \|v_{i}\|_{L^{\infty}([0,T]\times\Omega)} + \|v_{i}(T,\cdot)\|_{H^{1}(\Omega)}^{2} .\end{split}
$$

Thus, with the *L*∞ control established in the previous section, it is apparent that the v_i are bounded in $L^2([0,T],H^1(\Omega))$ provided that $v_i(T, \cdot) = \mathcal{I}_i v_T$ are bounded in $H^1(\Omega)$; this condition holds if $v(T, \cdot) \in W^{1,\infty}(\Omega)$. The first convergence result for the gradient is therefore that, owing to the Banach-Alaoglu theorem, $v_i\to v$ weakly in $L^2([0,T],H^1(\Omega))$, using *L*[∞]((0,*T*) × Ω) convergence to pass from *L*²([0,*T*], *H*¹(Ω)) weak convergence of subsequences to *L*²([0,*T*], *H*¹(Ω)) weak convergence of the whole sequence.

The question arises under which circumstances the convergence in the gradient is also strong. We demonstrate this under the below Assumption 6. We note that supposing (29) points towards uniform ellipticity of L^a . Let Λ_0 be the level set $\{(t, x) \in (0, T) \times \Omega : v(t, x) = 0\}$. For a smooth *v* the boundary of Λ_0 is always a $d-1$ dimensional set if 0 is a regular value.

Assumption 6. *The value function v belongs to the space* $W = W^{1,\infty}((0,T) \times \Omega)$ *and the d-dimensional Lebesgue measure of the boundary of* Λ_0 *vanishes:* vol($\partial \Lambda_0$) = 0. The coefficients \bar{a}^α_i and \bar{a}^α_i belong to $W^{1,\infty}(\Omega)$ and (29) is satisfied.

Let us suppose momentarily that there are approximations $Q_i v \in W_i$ to v such that $Q_i v \leq v_i$ for all $i \in \mathbb{N}$ and

$$
\lim_{i \to \infty} ||v - Q_i v||_{L^2([0,T],H^1(\Omega))} = 0,
$$

and

$$
\lim_{i \to \infty} \sum_{k=0}^{(T/h_i)-1} \langle \! \langle (h_i \mathsf{E}_i^{\alpha} - \mathsf{Id}) Q_i v(s_i^{k+1}, \cdot) + (h_i \mathsf{I}_i^{\alpha} + \mathsf{Id}) Q_i v(s_i^k, \cdot), (v_i - Q_i v)(s_i^k, \cdot) \rangle \! \rangle \to 0. \tag{30}
$$

We will construct such $Q_i v$ below. With $\xi^k = v_i(s_i^k, \cdot) - Q_i v(s_i^k, \cdot)$,

$$
|v_{i} - Q_{i}v|_{L^{2}([0,T],H^{1}(\Omega))}^{2} \lesssim \sum_{k=0}^{(T/h_{i})-1} \langle\!\langle (h_{i}E_{i}^{\alpha} - \text{Id})\xi^{k+1} + (h_{i}|_{i}^{\alpha} + \text{Id})\xi^{k}, \xi^{k}\rangle\!\rangle
$$

\n
$$
= \sum_{k=0}^{(T/h_{i})-1} \langle\!\langle (h_{i}E_{i}^{\alpha} - \text{Id})v_{i}(s_{i}^{k+1}, \cdot) + (h_{i}|_{i}^{\alpha} + \text{Id})v_{i}(s_{i}^{k}, \cdot), \xi^{k}\rangle\!\rangle - \sum_{k=0}^{(T/h_{i})-1} \langle\!\langle (h_{i}E_{i}^{\alpha} - \text{Id})Q_{i}v(s_{i}^{k+1}, \cdot) + (h_{i}|_{i}^{\alpha} + \text{Id})Q_{i}v(s_{i}^{k}, \cdot), \xi^{k}\rangle\!\rangle
$$

\n
$$
\leq \sum_{k=0}^{(T/h_{i})-1} \langle\!\langle h_{i}C_{i}^{\alpha}, \xi^{k}\rangle\!\rangle - \sum_{k=0}^{(T/h_{i})-1} \langle\!\langle (h_{i}E_{i}^{\alpha} - \text{Id})Q_{i}v(s_{i}^{k+1}, \cdot) + (h_{i}|_{i}^{\alpha} + \text{Id})Q_{i}v(s_{i}^{k}, \cdot), \xi^{k}\rangle\!\rangle, \tag{31}
$$

using in ($*$) the numerical scheme, $\xi^{T/h_i} = 0$ and that, due to the assumptions on the Q_i , the sign of $v_i - Q_i v$ is known. Since

$$
\sum_{k=0}^{(T/h_i)-1} \langle\!\!\!\langle h_i C_i^{\alpha}, \xi^k \rangle\!\!\rangle \leq \|d_i^{\alpha}\|_{L^2(\Omega)} \sum_{k=0}^{(T/h_i)-1} h_i \Big(\|v_i(s_i^k, \cdot) - v(s_i^k, \cdot) \|_{L^2(\Omega)} + \|v(s_i^k, \cdot) - Q_i v(s_i^k, \cdot) \|_{L^2(\Omega)} \Big)
$$

$$
\lesssim \|d_i^{\alpha}\|_{L^2(\Omega)} \Big(\|v_i - v\|_{L^2((0,T)\times\Omega)} + \|v - Q_i v\|_{L^2((0,T)\times\Omega)} \Big),
$$

the first term in (31) vanishes as *i* → ∞. The second term vanishes due to (30). Hence $|v_i - v|_{L^2([0,T],H^1(\Omega))} \to 0$ as *i* → ∞.

Theorem 4. *If there is an α* ∈ *A such that Assumption 6 holds, then the numerical solutions converge to the exact solution strongly in* $L^2([0, T], H^1(\Omega))$ *.*

Proof. It remains to show that suitable *Qⁱ* can be constructed, given Assumption 6. Denoting the nodal interpolant on $[0, T] \times \overline{\Omega}$ by \mathcal{I}_i we define

$$
Q_i: W \to W_i, w \mapsto \mathcal{I}_i \max\{w - ||v - v_i||_{L^{\infty}((0,T) \times \Omega)}, 0\}.
$$
\n
$$
(32)
$$

Observe that the max operator in (32) switches between the first and second argument in the vicinity of $\partial Λ_0$ for *i* sufficiently large. Furthermore, $Q_i v \in W_i$ satisfies homogeneous boundary conditions and $Q_i v \leq v_i$ and, by the mean value theorem,

$$
||Q_i v||_{W^{1,\infty}((0,T)\times\Omega)} \le ||v||_{W^{1,\infty}((0,T)\times\Omega)}.
$$

Note also that for all nodes y_i^{ℓ} and time levels s_i^k

$$
0 \le (v_i - Q_i v) (s_i^k, y_i^\ell) = \min \{ (v_i - v) (s_i^k, y_i^\ell) + ||v_i - v||_{L^\infty((0, T) \times \Omega)}, v_i (s_i^k, y_i^\ell) \} \le 2 ||v_i - v||_{L^\infty((0, T) \times \Omega)}.
$$

Consider the set Γ*ⁱ* of points which is not 'affected by the cut-off below 0' in (32) in the sense that

$$
\Gamma_i := \big\{ (t, x) \in (0, T) \times \Omega : \inf_{j \ge i} Q_j v(t, x) > 0 \text{ or } (t, x) \in \Lambda_0 \big\}.
$$

The set Γ'_i contains the points which are at least one element's length away from the boundary of $\Gamma_i \setminus \partial \Lambda_0$:

$$
\Gamma'_i := \big\{ (t,x) \in \Gamma_i : \{(s,y) \in (0,T) \times \Omega : \|(t,x)-(s,y)\| < \sup_{j \geq i} h_j + (\Delta x)_j \} \subset \Gamma_i \setminus \partial \Lambda_0 \big\}.
$$

Notice that Γ_i and Γ'_i are hierarchical families. Since $\|v - v_i\|_{L^\infty((0,T)\times\Omega)} \to 0$ and $h_i + (\Delta x)_i \to 0$ as $i \to \infty$ it follows that

$$
\bigcup_{i\in\mathbb{N}}\Gamma'_i=\big((0,T)\times\Omega\big)\setminus\partial\Lambda_0.
$$

Crucially, $(\partial_t Q_j v)|_{\Gamma'_i} = (\partial_t \mathcal{I}_j v)|_{\Gamma'_i}$ and $(\nabla Q_j v)|_{\Gamma'_i} = (\nabla \mathcal{I}_j v)|_{\Gamma'_i}$ for $j \ge i$.

For every $\varepsilon > 0$ there are $i, j \in \mathbb{N}$ such that $\text{vol}(\Omega \setminus \Gamma'_i) \leq \varepsilon^2$ and $||Q_k v - v||_{H^1(\Gamma'_i)} \leq \varepsilon$ for all $k \geq j$. Therefore

$$
\|Q_k v - v\|_{H^1((0,T)\times\Omega)} \lesssim \|Q_k v - v\|_{H^1(\Gamma'_i)} + \sqrt{\mathrm{vol}(\Omega \setminus \Gamma'_i)} \|v\|_{W^{1,\infty}((0,T)\times\Omega)} \leq \varepsilon (1 + \|v\|_{W^{1,\infty}((0,T)\times\Omega)}),
$$

giving strong convergence in $H^1((0,T)\times \Omega)$, meaning convergence in the spatial gradient and the time derivative. The terms connected to the time derivative in (30) vanish in the limit as

$$
\sum_{k=0}^{(T/h_i)-1} \langle \! \langle Q_i v(s_i^{k+1}, \cdot) - Q_i v(s_i^k, \cdot), \xi^k \rangle \! \rangle = \sum_{k=0}^{(T/h_i)-1} h_i \langle \! \langle (\partial_t Q_i v) \vert_{(s_i^k, s_i^{k+1})}, \xi^k \rangle \! \rangle \lesssim \| \partial_t v \|_{L^2((0,T)\times\Omega)} \| \xi^k \|_{L^2((0,T)\times\Omega)}.
$$

Recall that

$$
\langle \langle [a_{i}^{\alpha} Q_{i} \nu(s_{i}^{k},\cdot),\xi^{k}\rangle \rangle = \sum_{\ell} (\nu_{i} - Q_{i} \nu)(s_{i}^{k},y_{i}^{\ell}) \Big(\bar{a}_{i}^{\alpha}(y_{i}^{\ell})\langle \nabla Q_{i} \nu(s_{i}^{k},\cdot),\nabla \phi_{i}^{\ell}\rangle + \langle \bar{b}_{i}^{\alpha} \cdot \nabla Q_{i} \nu(s_{i}^{k},\cdot) + \bar{c}_{i}^{\alpha} Q_{i} \nu(s_{i}^{k},\cdot),\phi_{i}^{\ell}\rangle \Big).
$$

The lower-order terms vanish due to the uniform convergence of $v_i - Q_i v$ to 0 and the bound

$$
\sup_i \|\bar{\bar{b}}_i^{\alpha}\cdot \nabla Q_i\, \nu(s_i^k,\cdot) + \bar{\bar{c}}_i^{\alpha}\, Q_i\, \nu(s_i^k,\cdot)\|_{L^{\infty}(\Omega)} < \infty.
$$

We note for the second-order term that

$$
\sum_\ell (\nu_i-Q_i\nu)(s_i^k,y_i^\ell)\bar{\bar{a}}_i^\alpha(y_i^\ell)\langle \nabla Q_i\nu(s_i^k,\cdot),\nabla\phi_i^\ell\rangle=\langle \nabla Q_i\nu(s_i^k,\cdot),\nabla \mathfrak{I}_i(\bar{\bar{a}}_i^\alpha(\nu_i-Q_i\nu))(s_i^k,\cdot)\rangle,
$$

so that in (30) the implicit part of the second-order term becomes

$$
\sum_{k=0}^{(T/h_i)-1} h_i \langle \nabla Q_i v(s_i^k, \cdot), \nabla \mathcal{I}_i (\bar{a}_i^{\alpha} (v_i - Q_i v))(s_i^k, \cdot) \rangle = \int_0^T \langle \mathcal{J}_i \nabla Q_i v, \mathcal{J}_i \nabla \mathcal{I}_i (\bar{a}_i^{\alpha} (v_i - Q_i v)) \rangle dt,
$$
\n(33)

where \mathcal{J}_i maps any $w:[0,T]\to L^2(\Omega;\mathbb{R}^d)$ onto the step function with $(\mathcal{J}_i w)|_{[s_i^k,s_i^{k+1})}\equiv w(s_i^k,\cdot)$. Note that $\mathcal{J}_i \nabla Q_i v$ converges strongly in $L^2((0, T) \times \Omega; \mathbb{R}^d)$. At a time $s_i^k \in [0, T)$ the bound

$$
\|\nabla \mathcal{I}_i(\bar{\bar{a}}_i^{\alpha}(\nu_i - Q_i \nu))\|_{L^2(\Omega;\mathbb{R}^d)} \lesssim \|\nabla \mathcal{I}_i(\bar{\bar{a}}_i^{\alpha} \nu_i)\|_{L^2(\Omega;\mathbb{R}^d)} + \|\bar{\bar{a}}_i^{\alpha} Q_i \nu\|_{W^{1,\infty}(\Omega)} \lesssim \|\bar{\bar{a}}_i^{\alpha}\|_{W^{1,\infty}(\Omega)} \cdot (\|v_i\|_{H^1(\Omega)} + \|v\|_{W^{1,\infty}(\Omega)})
$$

follows from an inverse estimate and

$$
\sum_{T} \|\nabla \mathcal{I}_{i}(\bar{a}_{i}^{\alpha} \nu_{i})\|_{L^{2}(T;\mathbb{R}^{d})}^{2} \lesssim \sum_{T} (\Delta x)_{T}^{d} \|\nabla \mathcal{I}_{i}(\bar{a}_{i}^{\alpha} \nu_{i})\|_{L^{\infty}(T;\mathbb{R}^{d})}^{2} \lesssim \sum_{T} \|\bar{a}_{i}^{\alpha}\|_{W^{1,\infty}(T)}^{2} \left((\Delta x)_{T}^{d} \|\nu_{i}\|_{W^{1,\infty}(T)}^{2}\right).
$$

The convergence

$$
\lim_{i \to \infty} \int_0^T \langle w, \mathcal{J}_i \nabla \mathcal{I}_i (\bar{a}_i^{\alpha}(v_i - Q_i v)) \rangle dt = -\lim_{i \to \infty} \int_0^T \langle \nabla \cdot w, \mathcal{J}_I \mathcal{I}_i (\bar{a}_i^{\alpha}(v_i - Q_i v)) \rangle dt = 0
$$

with test functions w in the dense subset $C^1((0,T)\times \Omega;\mathbb{R}^d)$ gives weak convergence of $\nabla \mathcal{I}_i(\bar{a}_i^{\alpha}(\nu_i - Q_i \nu))$ in $L^2((0,T)\times \Omega)$ $\Omega; \R^d)$, see [28, p. 121]. Combing weak and strong convergence [29, Prop. 21.23], it is ensured that (33) converges to 0 as $i \to \infty$. A similar argument guarantees that $\sum_k h_i \langle \mathsf{E}_i^{\alpha} Q_i v(s_i^{k+1}, \cdot), \xi^k \rangle$ vanishes in the limit. \Box

The regularity of the exact value function ν is, for instance, discussed in Section IV.8 and IV.9 of [17]. Another item of Assumption 6, namely the justification of (29), is examined in the following example:

Example 3. *a) Suppose that a^α is positive and constant and, for all smooth w,*

$$
L^{\alpha} w = I^{\alpha} w = -a^{\alpha} \Delta w + b^{\alpha} \cdot \nabla w + c^{\alpha} w, \qquad E^{\alpha} w = 0,
$$

and, to obtain semi-definiteness in the lower-order terms, $c^{\alpha} - \frac{1}{2} \nabla \cdot b^{\alpha} \ge 0$ *. Then, for* $w \in W_i$ *,*

$$
a^{\alpha} |w|_{L^{2}([0,T],H^{1}(\Omega))}^{2} \lesssim a^{\alpha} \sum_{k=0}^{(T/h_{i})-1} h_{i} \langle \nabla w(s_{i}^{k},\cdot),\nabla w(s_{i}^{k},\cdot) \rangle + ||w(T,\cdot)||_{H^{1}(\Omega)}^{2} = \sum_{k=0}^{(T/h_{i})-1} h_{i} \langle \langle |_{i}^{\alpha} w(s_{i}^{k},\cdot), w(s_{i}^{k},\cdot) \rangle + ||w(T,\cdot)||_{H^{1}(\Omega)}^{2}.
$$

b) Suppose that $a^{\alpha} \in W^{2,\infty}(\Omega)$ is non-constant, positive, uniformly bounded from below and that $c^{\alpha} - \frac{1}{2}(\nabla \cdot b^{\alpha} + \Delta a^{\alpha}) \ge 0$, *noting for smooth w:*

$$
\langle L^{\alpha} w, w \rangle = \langle a^{\alpha} \nabla w, \nabla w \rangle + \langle (c^{\alpha} - \frac{1}{2} (\nabla \cdot b^{\alpha} + \Delta a^{\alpha})) w, w \rangle.
$$

 A gain choosing a fully implicit scheme with $L^a=I^a$, the highest order term in $\langle\!\langle\mathbf{l}_i^a w,w\rangle\!\rangle$ is at time s_i^k :

$$
\sum_{\ell} w(s_i^k, y_i^{\ell}) a^{\alpha}(s_i^k, y_i^{\ell}) \langle \nabla w(s_i^k, \cdot), \nabla \phi_i^{\ell} \rangle = \langle \nabla w(s_i^k, \cdot), \nabla J_i(a^{\alpha}(s_i^k, \cdot) w(s_i^k, \cdot)) \rangle.
$$

According to Theorem 2.1 in [11] there is a constant C = $C(\|a^\alpha\|_{W^{2,\infty}(\Omega)})$ such that for i sufficiently large

$$
\langle \nabla w, \nabla \mathcal{I}_i(a^\alpha w) \rangle - \langle \nabla w, \nabla a^\alpha w \rangle \le ||\nabla w||_{L^2(\Omega;\mathbb{R}^d)} \cdot ||\mathcal{I}_i(a^\alpha w) - a^\alpha w||_{H^1(\Omega)} \le C (\Delta x)_i ||w||_{H^1(\Omega)}^2,
$$

using that the η appearing in the proof in [11] is defined in terms of nodal interpolation. It then follows from Poincaré's inequality that there is some C such that for $C(\Delta x)_i < \frac{1}{2} \inf_\Omega a^\alpha$ that $|w|^2_{H^1(\Omega)} \lesssim \langle \langle |_i^\alpha w, w \rangle \rangle$ for $w \in V_i^0$, implying (29).

VIII Example: the method of artificial diffusion

The purpose of this section is to provide a way of constructing the operators E_i^{α} and I_i^{α} in order to satisfy Assumptions 1 and 2. This approach, called the method of artificial diffusion, is based on the fact that for strictly acute meshes, the discrete Laplacian is monotone. Further details on the method of artificial diffusion and monotone finite element schemes may, for example, be found in [5], [9] and [27].

Let \mathfrak{T}_i be the mesh corresponding to the finite element space $V_i.$ Given a function $f:\Omega\to\mathbb{R}^d$ we denote

$$
|f|_T := \left(\sum_{j=1}^d \|f_j\|_{L^\infty(T)}^2\right)^{\frac{1}{2}}, \qquad T \in \mathfrak{T}_i, \ i \in \mathbb{N}.
$$

If *f* is elementwise constant then $|f|_T$ is simply the Euclidean norm of *f* on *T*. Let $(\Delta x)_T$ denote the diameter of *T*. We assume that the meshes \mathcal{T}_i are strictly acute [5] in the sense that there exists $\theta \in (0, \pi/2)$ such that

$$
\nabla \phi_i^{\ell} \cdot \nabla \phi_i^l|_{T} \le -\sin(\theta) |\nabla \phi_i^{\ell}|_{T} |\nabla \phi_i^l|_{T} \qquad \forall \ell, l \le N \,\forall i \in \mathbb{N}.
$$

We choose a splitting of the form $a^{\alpha} = \tilde{a}^{\alpha}_{i} + \tilde{a}^{\alpha}_{i}$, $b^{\alpha} = \bar{b}^{\alpha}_{i} + \bar{b}^{\alpha}_{i}$, $c^{\alpha} = \bar{c}^{\alpha}_{i} + \bar{c}^{\alpha}_{i}$ and $d^{\alpha} = d^{\alpha}_{i}$, where all terms are in $C(\overline{\Omega})$, \tilde{a}^{α}_{i} and $\tilde{\tilde{a}}_i^\alpha$ are non-negative and all \bar{c}_i^α and $\bar{\tilde{c}}_i^\alpha$ are non-negative and satisfy inequality (6). Choose non-negative $\tilde{v}_i^{\alpha,\ell}$ $\frac{\alpha}{i}$ and $\bar{\bar{\nu}}_i^{\alpha,\ell}$ *i* such that for all *T* which have y_i^{ℓ} as vertex:

$$
\left(\left| \bar{b}_i^{\alpha} \right|_T + (\Delta x)_T \left\| \bar{c}_i^{\alpha} \right\|_{L^{\infty}(T)} \right) \le \bar{v}_i^{\alpha, \ell} \sin(\vartheta) \left| \nabla \hat{\phi}_i^{\ell} \right|_T \text{vol}(T),\tag{35a}
$$

$$
\left(\|\bar{\bar{b}}_i^{\alpha}\|_{T} + (\Delta x)_{T} \|\bar{c}_i^{\alpha}\|_{L^{\infty}(T)} \right) \le \bar{\bar{v}}_i^{\alpha,\ell} \sin(\vartheta) \|\nabla \hat{\phi}_i^{\ell}\|_{T} \operatorname{vol}(T). \tag{35b}
$$

Choose \bar{a}_i^{α} and \bar{a}_i^{α} both in $C(\overline{\Omega})$ such that $\bar{a}_i^{\alpha}(y_i^{\ell}) \ge \max\{\tilde{a}_i^{\alpha}(y_i^{\ell}), \tilde{v}_i^{\alpha,\ell}\}$ $\{\vec{a}_i^{\alpha}(y_i^{\ell}) \ge \max\{\tilde{a}_i^{\alpha}(y_i^{\ell}), \bar{v}_i^{\alpha,\ell}\}\}$ $\binom{\alpha,\ell}{i}$. Now suppose that $w \in V_i$ has a non-positive minimum at an interior node y_i^{ℓ} . By extending the arguments of [5], we show that

$$
(\mathsf{E}_i^{\alpha} w)_{\ell} \le 0, \qquad (\mathsf{I}_i^{\alpha} w)_{\ell} \le 0. \tag{36}
$$

We illustrate the proof of (36) for the implicit term. From the strict acuteness condition on the mesh, it can be shown that on the restriction to *T* [5, Lemma 3.1]

$$
\nabla w \cdot \nabla \phi_i^{\ell} = \cos \left(\angle (\nabla w, \nabla \phi_i^{\ell}) \right) |\nabla w|_T |\nabla \phi_i^{\ell}|_T \leq -\sin(\vartheta) |\nabla w|_T |\nabla \phi_i^{\ell}|_T.
$$

Using $\bar{c}_i^{\alpha} \geq 0$, $w(y_i^{\ell}) \leq 0$ and $\|\hat{\phi}_i^{\ell}\|_{L^1(\Omega)} = 1$,

$$
\langle \bar{c}_i^{\alpha} w, \hat{\phi}_i^{\ell} \rangle = \int_{\Omega} \bar{c}_i^{\alpha}(x) \left(w(y_i^{\ell}) + \nabla w(x) \cdot (x - y_i^{\ell}) \right) \hat{\phi}_i^{\ell}(x) dx
$$

$$
\leq \int_{\Omega} \bar{c}_i^{\alpha}(x) \nabla w(x) \cdot (x - y_i^{\ell}) \hat{\phi}_i^{\ell}(x) dx \leq \sum_{T} ||\bar{c}_i^{\alpha}||_{L^{\infty}(T)} ||\nabla w||_T (\Delta x)_T.
$$

Consequently,

$$
\begin{split} (\mathsf{I}_{i}^{\alpha}w)_{\ell}&=\bar{\bar{a}}_{i}^{\alpha}(y_{i}^{\ell})\langle\nabla w,\nabla \hat{\phi}_{i}^{\ell}\rangle+\langle\bar{\bar{b}}_{i}^{\alpha}\cdot\nabla w+\bar{\bar{c}}_{i}^{\alpha}\,w,\hat{\phi}_{i}^{\ell}\rangle\\ &\leq\sum_{T}-\bar{\bar{a}}_{i}^{\alpha}(y_{i}^{\ell})\sin(\vartheta)|\nabla w|_{T}|\nabla \hat{\phi}_{i}^{\ell}|_{T}\,\mathrm{vol}(T)+|\bar{\bar{b}}_{i}^{\alpha}|_{T}|\nabla w|_{T}+\|\bar{\bar{c}}_{i}^{\alpha}\|_{L^{\infty}(T)}|\nabla w|_{T}\,(\Delta x)_{T}\\ &\leq\sum_{T}|\nabla w|_{T}\big(\big(|\bar{\bar{b}}_{i}^{\alpha}|_{T}+(\Delta x)_{T}\|\bar{\bar{c}}_{i}^{\alpha}\|_{L^{\infty}(T)}\big)-\bar{\bar{v}}_{i}^{\alpha,\ell}\,\sin(\vartheta)\,|\nabla \hat{\phi}_{i}^{\ell}|_{T}\,\mathrm{vol}(T)\big)\leq0.\end{split}
$$

The proof of $(E_i^a w)_\ell \le 0$ is analogous. As hat functions ϕ_i^ℓ attain a non-positive minimum at all y_i^j i_j where $j \neq \ell$, all offdiagonal entries of E_i^{α} are non-positive. Hence with a suitable time step restriction the $h_i E_i^{\alpha}$ – Id are monotone, which ensures that Assumption 2 is satisfied.

The scaling of the terms in (35) with respect to (∆*x*)*^T* leads to Assumption 1. Due to shape-regularity all elements *T* on a patch are of comparable size; giving $\|\phi_i^{\ell}\|_{L^1(\Omega)} \le C$ vol (T) for all T \subset supp ϕ_i^{ℓ} with a constant C which is independent of *h* and *ℓ*. Hence in (35), we see that

$$
\text{vol}(T) \, |\nabla \hat{\phi}_i^\ell|_T \ge \frac{\text{vol}(T)}{(\Delta x)_T \, \|\phi_i^\ell\|_{L^1(\Omega)}} \ge \frac{1}{C(\Delta x)_T}.
$$

Thus, if $\bar{v}^{\alpha,\ell}_i$ $\frac{\alpha}{i}$ and $\bar{\bar{\nu}}_i^{\alpha,\ell}$ a, l are chosen optimally then for $T \subset \text{supp } \phi_i^l$

$$
\bar{v}_i^{\alpha,\ell} = O\left(\sup_T \left\{ |\bar{b}_i^{\alpha}|_T (\Delta x)_T + ||\bar{c}_i^{\alpha}|_{L^{\infty}(T)} (\Delta x)_T^2 \right\} \right), \qquad \bar{v}_i^{\alpha,\ell} = O\left(\sup_T \left\{ |\bar{b}_i^{\alpha}|_T (\Delta x)_T + ||\bar{c}_i^{\alpha}|_{L^{\infty}(T)} (\Delta x)_T^2 \right\} \right).
$$
\n(37)

With (37) in mind we return to the time step restriction for semi-implicit and explicit methods. The non-positivity of the diagonal terms of $h_i \mathsf{E}_i^{\alpha}$ – ld expands to

$$
1 \ge h_i \Big(\bar{a}_i^{\alpha} (y_i^{\ell}) \langle \nabla \phi_i^{\ell}, \nabla \hat{\phi}_i^{\ell} \rangle + \langle \bar{b}_i^{\alpha} \cdot \nabla \phi_i^{\ell} + \bar{c}_i^{\alpha} \phi_i^{\ell}, \hat{\phi}_i^{\ell} \rangle \Big) = h_i \Big(O \Big(\bar{a}_i^{\alpha} (\Delta x)^{-2} \Big) + O \Big(|\bar{b}_i^{\alpha}|_T (\Delta x)^{-1} \Big) + O \Big(\bar{c}_i^{\alpha} \Big) \Big).
$$

Therefore the time step restriction imposed by L^α is $h_i\lesssim \sup_T((\Delta x)_T^2/\bar a_i^\alpha(y_i^\ell)),$ $y_i^\ell\in\overline{T}$, if there is a non-zero $\tilde a_i^\alpha$ and i is large. It is $h_i \lesssim \sup_T((\Delta x)_T/\vert \bar b_i^\alpha(y_i^\ell)\vert_T)$ if all $\bar a_i^\alpha=0$, $i\in\mathbb N$, and there are non-zero $\bar b_i^\alpha$, and is O(1) if all $\bar a_i^\alpha$ and $\bar b_i^\alpha$ vanish. There is no restriction if also all \bar{c}^{α}_i are zero.

References

- [1] G. Barles and E.R. Jakobsen, *Error bounds for monotone approximation schemes for parabolic Hamilton-Jacobi-Bellman equations*, Math. Comp. 76(240):1861–1893, 2007.
- [2] G. Barles and B. Perthame, *Exit time problems in optimal control and vanishing viscosity method*, SIAM J. Control Optim., 26(5):1133–1148, 1988.
- [3] G. Barles and P.E. Souganidis, *Convergence of approximation schemes for fully nonlinear second order equations*, J. Asymptotic Analysis, 4:271–283, 1991.
- [4] O. Bokanowski and S. Maroso and H. Zidani, *Some convergence results for Howard's algorithm*, SIAM J. Num. Anal., 47(4):3001– 3026, 2009.
- [5] E. Burman and A. Ern, *Nonlinear diffusion and discrete maximum principle for stabilized Galerkin approximations of the convection-diffusion-reaction equation*, Comput. Methods Appl. Mech. Engrg., 191(35):3833–3855, 2002.
- [6] E. Burman and A. Ern, *Stabilized Galerkin approximation of convection-diffusion-reaction equations: discrete maximum principle and convergence*, Math. Comp., 74(252):1637–1652, 2005.
- [7] F. Camilli and M. Falcone, *An approximation scheme for the optimal control of diffusion processes*, RAIRO Modél. Math. Anal. Numér. 29(1):97–122, 1995.
- [8] F. Camilli and E.R. Jakobsen, *A finite element like scheme for integro-partial differential Hamilton-Jacobi-Bellman equations*, SIAM J. Numer. Anal. 47(4):2407–2431, 2009.
- [9] R. Codina, *A discontinuity capturing crosswind dissipation for the finite element solution of the convection-diffusion equation*, Comput. Meth. Appl. Mech. Engrg. 110(3–4):325–342, 1993.
- [10] M.G. Crandall and H. Ishii and P.-L. Lions, *User's guide to viscosity solutions of second order partial differential equations*, Bull. Amer. Math. Soc. 27(1):1–67, 1992.
- [11] A. Demlow, J. Guzmán, A.H. Schatz, *Local energy estimates for the finite element method on sharply varying grrids*, Math. Comp. 80(273):1–9, 2011.
- [12] A. Demlow, D. Leykekhman, A.H. Schatz, and L.B. Wahlbin, *Best approximation property in the W* $_{\infty}^1$ norm on graded meshes, Math. Comp., to appear.
- [13] H. Dong and N.V. Krylov, *The rate of convergence of finite-difference approximations for parabolic Bellman equations with Lipschitz coefficients in cylindrical domains*, Appl. Math. Optim. 56(1):37–66, 2007.
- [14] A. Ern and J-L. Guermond, *Theory and practice of finite elements*, Springer, 2004.
- [15] X. Feng and R. Glowinsky and M. Neilan, *Recent developments in numerical methods for fully nonlinear second order partial differential equations*, submitted to SIAM Review.
- [16] X. Feng and M. Neilan, *The vanishing moment method for fully nonlinear second order partial differential equations: formulation, theory, and numerical analysis*, arXiv:1109.1183, 2011.
- [17] W. Fleming and H. Soner, *Controlled Markov processes and viscosity solutions*, Stochastic Modelling and Applied Probability, vol. 25, 2nd ed., Springer, New-York, 2006.
- [18] R. Horn and C. Johnson, *Topics in matrix analysis*, Cambridge University Press, 1991.
- [19] R. Howard, *Dynamic Programming and Markov Processes*, The Technology Press of M.I.T., Cambridge, Mass. and John Wiley & Sons, Inc., New York-London, 1960.
- [20] N.V. Krylov, *On the rate of convergence of finite–difference approximations for Bellman's equations*, Algebra i Analiz, 9(3):245–256, 1997; translation in St. Petersburg Math. J. 9(3):639–650.
- [21] N.V. Krylov, *On the rate of convergence of finite–difference approximations for Bellman's equations with variable coefficients*, Probab. Theory Related Fields 117(1):1–16, 2000.
- [22] H.J. Kushner, *Numerical methods for stochastic control problems in continuous time*, SIAM J. Control and Optimization 28(5):999– 1048, 1990.
- [23] H.J. Kushner and P. Dupuis, *Numerical methods for stochastic control problems in continuous time*, Applications of Mathematics 24, Springer-Verlag, 2001.
- [24] P.-L. Lions and B. Mercier, *Approximation numérique des équations de Hamilton-Jacobi-Bellman*, RAIRO Anal. Numér., 14(4):369– 393, 1980.
- [25] O.A. Oleinik and E.V. Radkevič, *Second order equations with nonnegative characteristic form*, AMS, 1973.
- [26] J.-P. Quadrat, *Sur l'identification et le contrôle de systèmes dynamiques stochastiques*, Thèse d'Etat, Paris 9, 1981.
- [27] J. Xu and L. Zikatanov, *A monotone finite element scheme for convection-diffusion equations*, Mathematics of Computation, 68(228):1429–1446, 1999.
- [28] K. Yosida, *Functional Analysis*, 6th ed., Springer, New York, 1980.
- [29] E. Zeidler, *Nonlinear functional analysis and its applications II*, Springer, New York, 1990.