

Sussex Research

Interpreting high [O III]/H ß ratios with maturing starbursts

Elizabeth R Stanway, John J Eldridge, Stephanie M L Greis, Luke J M Davies, Stephen Wilkins, Malcolm N Bremer

Publication date

11-11-2014

Licence

This work is made available under the Copyright not [evaluated](https://rightsstatements.org/page/CNE/1.0/?language=en) licence and should only be used in accordance with that licence. For more information on the specific terms, consult the repository record for this item.

Citation for this work (American Psychological Association 7th edition)

Stanway, E. R., Eldridge, J. J., Greis, S. M. L., Davies, L. J. M., Wilkins, S., & Bremer, M. N. (2014). Interpreting high [O III]/H ß ratios with maturing starbursts (Version 1). University of Sussex. https://hdl.handle.net/10779/uos.23423273.v1

Published in

Monthly Notices of the Royal Astronomical Society

Link to external publisher version

<https://doi.org/10.1093/mnras/stu1682>

Copyright and reuse:

This work was downloaded from Sussex Research Open (SRO). This document is made available in line with publisher policy and may differ from the published version. Please cite the published version where possible. Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners unless otherwise stated. For more information on this work, SRO or to report an issue, you can contact the repository administrators at [sro@sussex.ac.uk.](mailto:sro@sussex.ac.uk) Discover more of the University's research at <https://sussex.figshare.com/>

Sussex Research Online

Interpreting high [O III]/H β ratios with maturing starbursts

Article (Unspecified)

Stanway, Elizabeth R, Eldridge, John J, Greis, Stephanie M L, Davies, Luke J M, Wilkins, Stephen M and Bremer, Malcolm N (2014) Interpreting high [O III]/H β ratios with maturing starbursts. Monthly Notices of the Royal Astronomical Society, 444 (4). pp. 3466-3472. ISSN 0035-8711

This version is available from Sussex Research Online: http://sro.sussex.ac.uk/id/eprint/57540/

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Interpreting high [O III]/H*β* **ratios with maturing starbursts**

Elizabeth R. Stanway, \rightarrow John J. Eldridge, ² Stephanie M. L. Greis, ^{[1](#page-2-0)} Luke J. M. Davies, 3 Stephen M. Wilkins⁴ and Malcolm N. Bremer⁵

¹*Department of Physics, University of Warwick, Gibbet Hill Road, Coventry CV4 7AL, UK*

²*Department of Physics, University of Auckland, Private Bag 92019, Auckland, New Zealand*

³*ICRAR, The University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia*

⁴*Astronomy Centre, Department of Physics and Astronomy, University of Sussex, Brighton BN1 9QH, UK*

⁵*H. H. Wills Physics Laboratory, University of Bristol, Tyndall Avenue, Bristol BS8 1TL, UK*

Accepted 2014 August 15. Received 2014 August 5; in original form 2014 May 28

ABSTRACT

Star-forming galaxies at high redshift show ubiquitously high-ionization parameters, as measured by the ratio of optical emission lines. We demonstrate that local $(z < 0.2)$ sources selected as Lyman break analogues also manifest high line ratios with a typical [O $\text{III}/\text{H}\beta = 3.36^{+0.14}_{-0.04}$ – comparable to all but the highest ratios seen in star-forming galaxies at $z \sim 2-4$. We argue that the stellar population synthesis code BPASS can explain the high-ionization parameters required through the ageing of rapidly formed star populations, without invoking any AGN contribution. Binary stellar evolution pathways prolong the age interval over which a starburst is likely to show elevated line ratios, relative to those predicted by single stellar evolution codes. As a result, model galaxies at near-solar metallicities and with ages of up to ∼100 Myr after a starburst typically have a line ratio [O $\text{III}/\text{H}\beta \sim 3$, consistent with those seen in Lyman break galaxies and local sources with similar star formation densities. This emphasises the importance of including binary evolution pathways when simulating the nebular line emission of young or bursty stellar populations.

Key words: galaxies: evolution – galaxies: high-redshift – galaxies: star formation.

1 INTRODUCTION

Understanding the sites of star formation in the distant Universe is key to developing our picture of the early stages of galaxy evolution. The low-mass, low-metallicity, intensely star-forming galaxies observed in deep field surveys are the building blocks which form the more massive systems we currently inhabit and observe evolve. They are also the most likely source of the energetic photons that ionized the intergalactic medium (IGM) at early times (e.g. Bunker et al. [2010\)](#page-8-0), creating the conditions which persist to the current day. The source and spectrum of those ionizing photons are key parameters in cosmological simulations, affecting the process by which small regions of ionized hydrogen surrounding the first galaxies grow and eventually overlap.

However, spectroscopy of such distant sources pushes the technical limits of existing spectrographs, and is often impossible. Only the most highly lensed galaxies, or extreme examples such as submillimetre galaxies or quasar hosts, are sufficiently luminous to measure optical emission lines at $z > 5$, where the rest-UV and optical wavelength ranges have been shifted into the near-infrared. None the less, the fitting of spectral energy distributions (SEDs)

E-mail: e.r.stanway@warwick.ac.uk

across a broad redshift range ($z \sim 3-8$) has strongly suggested that the majority of high-redshift star-forming galaxies may show strong optical line emission, which contributes significantly to their observed flux at 3–5 µm (González et al. [2012;](#page-8-1) Stark et al. [2013;](#page-8-2) de Barros, Schaerer & Stark [2014\)](#page-8-3). Local galaxies selected to have similar ultraviolet emission densities are also confirmed to have very prominent emission lines (Heckman et al. [2005;](#page-8-4) Stanway & Davies [2014\)](#page-8-5).

At slightly lower redshifts, $z \sim 2-4$, direct measurements of the rest-optical emission spectrum become possible. Work by Holden et al. [\(2014\)](#page-8-6) compiled *K*-band spectroscopy on a sample of 67 $z \sim 3.5$ galaxies, determining that high [O III 5007 Å]/H β ratios are ubiquitous in the high-redshift population. More recently still, Steidel et al. [\(2014\)](#page-8-7) compiled a large sample of 179 2.0 $< z < 2.6$ objects with near-infrared spectroscopy and confirmed that these high-redshift, star-forming galaxies occupy a distinct locus with higher line ratios than seen in typical local galaxies. However, efforts to model these high ratios are problematic, requiring ionization parameters orders of magnitudes higher than those typically seen in local galaxies (Dopita et al. [2000;](#page-8-8) Kewley et al. [2001,](#page-8-9) [2013\)](#page-8-10) or invoking an otherwise unrealistic high oxygen abundance (Contini [2014\)](#page-8-11).

In this pape[r,](#page-3-0) we present measurements of the line ratios determined in a sample of local galaxies which are selected to match the

Figure 1. The distribution of O III/Hβ ratios seen in galaxies observed by the SDSS (Brinchmann et al. [2004,](#page-8-12) grey-scale). Overplotted on the left are the ratios seen at z ∼ 2–4 by Holden et al. [\(2014,](#page-8-6) asterisks), and Schenker et al. [\(2013,](#page-8-13) crosses). The ratios measured in the Lyman break analogue sample of Stanway & Davies [\(2014\)](#page-8-5), and presented here for the first time, are shown on the right. The measurement error on the spectra of these bright sources is smaller than the plotting symbols. The mass uncertainty in any SED fitting analysis is at least ∼0.3 dex. As both high- and low-redshift, UV-selected galaxies exceed the typical line ratios for their mass. Note that the high-mass, high line ratio sources in the local sample are typically AGN dominated.

distant population in photometric properties (Section 2), and consider possible interpretations in the light of models from the Binary Population and Spectral Synthesis (BPASS) models (Section 3). In Section 4, we discuss implications for our understanding of stellar populations in the distant Universe, before presenting our conclusions.

2 OBSERVED OPTICAL EMISSION LINE RATIOS

2.1 Evidence for high line ratios at $z > 1$

The ratios of optical spectral lines, arising from nebular emission, are dependent on the combination of ionizing ultraviolet flux incident on the ISM and its density. Amongst them, one of the most useful in the distant Universe is the ratio of the $[O \text{III}]$ 5007 Å line to the H β 4861 Å Balmer line. The small wavelength interval between these lines minimizes the impact of dust absorption uncertainties on their ratio, and they are more accessible at high redshifts $(z > 1)$ than the redder H α line region. Samples of distant star-forming galaxies with rest-frame optical spectroscopy have remained small nevertheless, largely due to the challenging nature of the observations which have required observations of individual targets from the ground or low-resolution grism spectra from the *Hubble Space Telescope* (e.g. Xia et al. [2012\)](#page-8-14). Lensed targets have been the most studied, for example in Hainline et al. [\(2009\)](#page-8-15) who used [O $\text{III}/\text{H}\beta$ and other line ratios to measure high-ionization parameters in three lensed galaxies at $z \sim 2-2.5$.

As multi-object near-infrared spectrographs become available, analyses of larger samples are possible. Nakajima et al. [\(2013\)](#page-8-16) found high-ionization parameters in $z = 2.2$ star-forming galaxies selected as Lyman α emitters, while Masters et al. [\(2014\)](#page-8-17) measured an [O III]/H β ratio of 4.12 \pm 0.01 in a composite spectrum of 24 z ∼ 2 galaxies, exceeding the predictions of photoionization models. Recent work by Holden et al. [\(2014\)](#page-8-6) compiled *K*-band spectroscopy on a sample of 67 z ∼ 3 ultraviolet-selected 'Lyman break' galaxies (LBGs, including 20 observed by Schenker et al. [2013\)](#page-8-13), determining a median of $[O \text{ III}]/\text{H}\beta = 4.8^{+0.8}_{-1.7}$ for the sample. In Fig. [1,](#page-3-0) we replot the high $[O \text{ III}]/H\beta$ line ratios measured by Holden et al. as a function of mass, together with the distribution of ratios observed in low-redshift galaxies (Brinchmann et al. [2004\)](#page-8-12). Distant galaxies lie well above the line ratios seen in local star-forming galaxies at the same mass. The hard ionizing radiation field required to explain these observations is very difficult to reproduce with normal stellar populations (Brinchmann, Pettini & Charlot [2008;](#page-8-18) Kewley et al. [2013\)](#page-8-10) at the metallicities seen at $z \sim 2-5$ (typically 0.1– $1.5 Z_O$, e.g. Pettini et al. 2001 ; Erb et al. 2006 ; Hainline et al. 2009 ; Richard et al. [2011\)](#page-8-21).

2.2 Analogues in the local universe

While the vast majority of galaxies in the local Universe differ in size, star formation rate and character from those at high redshift, it is none the less possible to identify local sources which appear to match the distant population in their continuum properties. In our recent paper (Stanway & Davies [2014\)](#page-8-5), we identified a pilot sample of such analogues, lying at $z \sim 0.05-0.25$. Galaxies were identified based on their ultraviolet luminosity and colour, and to ensure that the resulting sample matched the high specific star formation densities observed at $z \sim 5$. Given that high line ratios appear ubiquitous at $z > 1$, we would expect these to share that characteristic.

Our samples were also selected to have spectroscopy from the Sloan Digital Sky Survey (SDSS), precluding the presence of strong AGN activity and confirming their redshift. In Fig. [1,](#page-3-0) we present the $[O \text{ III}]/H\beta$ line ratios measured in our local analogue sample, as a function of stellar mass. Line ratios are measured at high signal to noise in SDSS spectroscopy for these relatively bright sources – we use line fluxes calculated by the SDSS pipeline, but have checked that these agree to within 0.02 dex with the independent MPA-JHU DR7 analysis.¹ Stellar masses are also taken from the MPA-JHU data base. These are derived from simultaneous fitting of age, dust, star formation history and stellar mass to the galaxies' SED. Thus they are uncertain at the ∼0.3 dex level, due to reliance on the synthesis model templates used for fitting and degeneracies in the resulting colours. The uncertainties in mass fitting of these sources will be explored in more detail in Greis et al. (in preparation).

As Fig. [1](#page-3-0) demonstrates, our analogue sample also lies well above the norm for line ratios in star-forming galaxies of the same mass in the local Universe (Brinchmann et al. [2004\)](#page-8-12), despite being drawn from the same underlying data set. While the analogue sample

¹ see www.mpa-garching.mpg.de/SDSS/DR7 (Brinchmann et al. [2004\)](#page-8-12).

Figure 2. The distribution of [O III]/Hβ ratios predicted by BPASS models as a function of stellar population age for single star (left) and binary (right) population stellar evolution pathways. Upper panels show the ageing of an instantaneous burst. Lower panels show the near-constant ratios seen in populations with stable star formation rates. Tracks are shown at three different metallicities.

does not match the most extreme high redshift examples, the ultraviolet continuum selection of our Lyman break analogue sample appears to identify galaxies with very comparable spectral properties to those seen in the distant Universe. We find that the lowredshift analogue sample presented here has a median of $[O \text{ III}]/H\beta$ $= 3.36^{+0.143}_{-0.04}$, where the quoted uncertainty gives the interquartile range, consistent with that quoted by Holden et al. [\(2014\)](#page-8-6) for their $z \sim 2-3$ sample.

3 MODELLING HIGH RATIOS WITH BPASS

3.1 BPASS

The $BPASS models²$ are a set of galaxy population synthesis models which were developed to address the effects of massive stars on the SEDs of galaxies (Eldridge, Izzard & Tout [2008;](#page-8-22) Eldridge & Stanway [2012,](#page-8-23) [2009\)](#page-8-24). Given a young stellar population, for example in the aftermath of a major star formation episode, the optical spectrum of galaxies is dominated by hot and massive stars which have not yet reached the ends of their lifespan. However, as the population ages, the population averaged colour, temperature and SED are all strongly influenced by the evolutionary state of the remaining massive stars. The processes of angular momentum transfer, mass-loss or mass gain due to a binary companion all modify this evolutionary state, allowing evolved secondary stars to extend the highly luminous phase, and boosting the population of rapidly rotating, hydrogen-depleted, Wolf–Rayet stars.

The BPASS code tracks the evolution of stellar populations, sampled from an initial mass function (IMF) and range of binary system properties, and creates a composite stellar spectrum at a given age. Binary evolution is treated explicitly for initial stellar masses \geq 5 M_{\odot}, and empirical terms are included for the evolution of rotationally mixed, quasi-homogeneous stars. For comparison, an equivalent population of single stars is also permitted to evolve. The radiative transfer of the stellar emission from both populations,

through a dust and gas screen within the source galaxy, is then modelled using the radiative transfer code CLOUDY (Ferland et al. [1998\)](#page-8-25) to assess the contribution of nebular continuum and line emission. The ionization parameter of the local radiation field is defined as a ratio of the number of ionizing photons to the local gas density. It is thus not a directly tunable parameter, but rather constructed through the combination of appropriate stellar atmosphere models and a choice of gas distribution. The assumed total hydrogen gas density of our baseline model set is 10^2 cm⁻³, distributed in a sphere around the stellar population. This is a fairly typical gas density for extragalactic star-forming H_{II} regions, although we note that these range over several orders of magnitude in density (see e.g. Hunt & Hirashita [2009\)](#page-8-26). We explore effects of varying the gas density relative to our baseline models in Section 3.3. The evolution of an instantaneous, rapid burst of star formation and of a continuous moderate (1 M_{\odot} yr⁻¹) star formation rate are modelled separately. In this paper, we use nebular emission line flux predictions determined as part of the current (v1.0) BPASS model data release.

3.2 Varying [O III]/H*β* **with age and metallicity**

The time evolution of the [O $\text{III}/\text{H}\beta$ ratio, at a fixed hydrogen gas density, is significantly affected by the introduction of binary stellar evolution pathways, as shown in Fig. [2.](#page-4-0) In both single and binary population models, a continuous star formation rate leads to a very stable $[O \text{ III}]/H\beta$ ratio, which does not vary significantly after the initial few Myr, since the line flux is always dominated by the youngest stars. This ratio can reproduce the observed high redshift (and local analogue data) at sub-solar metallicities, for either single or binary populations, but at solar metallicity, the predicted line ratio falls well below the measured values. Continuous star formation, observed at late times, may also be consistent with the bulk of observed H β emission line equivalent widths (as shown in Fig. [3\)](#page-5-0), but struggles to reproduce some of the lower observed values.

By contrast, the aging of a rapid burst leads to line ratios that are high at early ages, as the massive stars formed in the initial burst evolve off the main sequence and eventually undergo supernovae. In single star population models, these intervals of high line ratios

² See [http://bpass.org.uk/.](http://bpass.org.uk/)

Figure 3. The distribution of [O III]/H β ratios predicted by BPASS models as a function of rest-frame H β recombination line equivalent width for single star (left) and binary (right) population stellar evolution pathways. Labels and linestyles are as in Fig. [2.](#page-4-0) Age intervals of $log(age)=0.1$ are marked, and age increases to the left on each track in this parameter space. Pale grey points indicate high-redshift data from Schenker et al. [\(2013\)](#page-8-13) and Holden et al. [\(2014\)](#page-8-6). Red crosses indicate our local analogue sample, the error bars for which are smaller than the points.

([O III]/H β > 1) are brief, lasting no more than ∼10 Myr and occurring primarily at low metallities (5–40 per cent of solar). Such young stars are likely dust embedded and heavily extincted, making this an unattractive interpretation.

However, the evolution of a binary stellar population is very different. While the initial phase of \sim 1:1 [O III]/H β ratios is still seen, it is extended over a much longer epoch by the formation first of lower mass Wolf–Rayet stars than are possible in a single star population, and then by longer lived, hot, helium stars. A binary stellar population will exceed a 1:1 line ratio for up to 100 Myr after its initial formation epoch (roughly the lifetime of the minimum mass considered for binary stars in BPASS), and at much higher metallicities (∼40–100 per cent solar) than those seen in the single star case. At 40 per cent Z_{\odot} , the binary population reaches [O III]/H β line ratios >3 for much of its first 100 Myr, improving the probability that these ratios will be observed in conditions where a bursty, binary-rich population may dominate the emission.

During this interval, as Fig. [3](#page-5-0) shows, a binary population model also generates Balmer line luminosities and equivalent widths consistent with those seen in both the high-redshift sample and the low-redshift analogue population presented here. The single star instantaneous burst model, by contrast, cannot reproduce the line ratios without overproducing $H\beta$ line flux. Similarly, as noted above, the continuous star formation models (with or without binaries) struggle to reproduce the weakest $H\beta$ lines while simultaneously maintaining the strong line ratios. For both instantaneous and continuous star-forming models, the lowest line equivalent widths (and luminosities) are generated at late times, as the continuum contribution from older underlying stars builds up.

3.3 Varying [O III]/H*β* **with gas density**

While the emission line strengths from star-forming regions are a strong function of irradiating spectrum, they also depend on the geometry and density of the emitting gas. Self-shielding and collisional excitation or de-excitation of ions can alter the transition probabilities in either very sparse or very dense gas. As mentioned in Section 3.1, the baseline BPASS model set uses a total hydrogen gas density of 10^2 cm⁻³, distributed in a sphere around the stellar population to model the radiative transfer and nebular emission. While this is a reasonable gas density for extragalactic star-forming H II regions (see, for example, Hunt & Hirashita [2009\)](#page-8-26), it is possible that a difference in the typical gas density, rather than in the dominant irradiating spectrum, could be responsible for the line properties of the high-redshift starburst population and their local analogues.

In Fig. [4,](#page-6-0) we consider the effect of total hydrogen gas density on the line ratios as a function of age, for the instantaneous, $Z = 0.08$ starburst models that best fit the data at our baseline gas density (see Section 3.2).

The effect of changing the assumed density of the illuminated gas has a negligible effect on the lifetime of regions with high line ratios for single star populations – at all densities, the high line ratio epoch ends within 10 Myr of the onset of star formation. The effect of gas density is, however, seen in the strength of the initial [O $\text{III}/\text{H}\beta$ line ratios measured. The line ratio at a given age decreases systematically with decreasing gas density, except at the highest densities considered, 10^4 cm⁻³ – equivalent to the upper end of the H_{II} region distribution.

By contrast, when binary stellar populations are considered, both the strength of the ratio and the duration over which it remains elevated are functions of the assumed hydrogen gas density. At low densities, \langle 1 cm⁻³, the [O III]/H β ratio reached by the population increases with gas density, as does the lifetime over which it remains at the levels seen in the distant population. At higher densities, >1 cm⁻³, the [O III]/H β ratio remains very nearly constant at [O III]/H $\beta \sim 5$, but the epoch over which this level is reached becomes shorter with increasing gas density. At a density of 10^4 cm^{-3} , the lifetime of strong line emission is comparable to that seen in the single star populations.

Comparison with Fig. [1](#page-3-0) suggests that moderate gas densities, 10^{-2} −10⁴ cm^{−3} are required to reproduce the high line ratios seen in the distant population and their local analogues. The extended lifetime of enhanced [O $\text{III}/\text{H}\beta$ emission in binary populations at gas densities \sim 1−100 cm⁻³ and at moderate metallicities suggest

Figure 4. The distribution of $[O \text{ III}]/H\beta$ ratios predicted by BPASS models for single star (left) and binary (right) population stellar evolution pathways at $Z = 0.08$ (40 per cent solar), as a function of gas density.

that these properties may be consistent with the distant population. We note that this is not necessarily the scenario with the highest ionization parameter (which would correspond to the lowest density for a given stellar input spectrum) but is comparable to the densities seen in star-forming regions.

4 DISCUSSION

The effect of ionizing spectra (as manifest in the $[O \text{ III}]/H\beta$ ratio) has traditionally been attributed to an AGN contribution and is commonly seen in local Seyfert-type galaxies (Ferland & Netzer [1983\)](#page-8-27). An alternate explanation could conceivably be a very high oxygen abundance relative to hydrogen. To rule out this later possibility, we plot both observed examples of $[O \text{ III}] / H \beta$ ratios and predictions for binary stellar populations in the BPASS models against the largely abundance-independent [O II]/[O III] ratio (see e.g. Ferland & Netzer [1983\)](#page-8-27) in Figs [5](#page-6-1) and [6.](#page-6-2) A binary star formation model at $Z = 0.08$ and a gas density of 10² cm⁻³ (our baseline model) provides a remarkably good fit to the distribution of line ratios seen in both high- and low-redshift starburst galaxies, over an extended

Figure 5. The [O II]/[O III] ratios seen in the data, and predicted by BPASS models for binary populations at ages $10^{7}-10^{8.3}$ yr following an instantaneous burst. A smooth third-order polynomial has been fitted through the predictions at discrete model timesteps at each metallicity. Grey points are $z = 2-3$ LBGs (Pettini et al. [2001;](#page-8-19) Masters et al. [2014\)](#page-8-17), red points are our Lyman break analogue sample.

Figure 6. As Fig. [5,](#page-6-1) but now varying assumed gas density at a fixed metallicity $(Z = 0.08)$. Single star models are plotted as dashed lines, fitted at ages of 10^6 – $10^{7.2}$ yr after an instantaneous burst. For binary populations (solid lines), gas density has only slight effect on the line ratios, such that the models occupy a narrow locus in parameter space. The single star models span a broader range, and show a more pronounced evolution in line ratios with gas density.

period of the order of 100 Myr. In fact, the line ratios generated by binary stellar populations of ages <100 Myr shows only mild dependence on the surrounding gas density. The ratios do however show a strong dependence on binary as opposed to single star evolution pathways, with the latter typically producing lower line ratios and showing a stronger gas density dependence.

We find that high $[O \text{ III}]/H\beta$ ratios fall naturally out of a selfconsistent treatment of binary evolution in an ageing starburst stellar population, without invoking AGN emission or an unusual IMF (Masters et al. [2014\)](#page-8-17). The modest star formation rates and metallicities required to create such spectra do not require fine tuning of the conditions, but are well matched to the typical properties of both $z \sim 2-4$ LBGs (e.g. de Barros et al. [2014\)](#page-8-3) and lower redshift UV-selected Lyman break analogues (e.g. Stanway & Davies [2014,](#page-8-5) Greis et al.; in preparation), while being atypical of the $z = 0$ galaxy population. Given the discrete sampling in metallicity and age of the BPASS models, and the simple prescription for the dust and gas screen, it is uncertain whether the effect of binaries is sufficient to reproduce the highest observed line ratios, [O III]/H $\beta \sim 10$ in the

 $z = 2-4$ sample. However, the median object is well matched to post-starburst BPASS models at ∼0.4 Z_∩.

Holden et al. [\(2014\)](#page-8-6) argued against a bursty starburst model for $z = 2-4$ galaxies on the basis of consistency between star formation rates derived from $H\beta$, dominated by the youngest stellar population, and the rest-frame ultraviolet continuum. However, the time-scale for establishment of ultraviolet emission is of the order of ∼10−30 Myr. This would be problematic for a single stellar population, but is significantly shorter than the time-scales for elevated [O $\text{III}/\text{H}\beta$ ratios in binary populations, which would show an established UV continuum. As noted above, we have studied the BPASS prediction for Balmer line luminosity and equivalent width and find it consistent with the observed high-redshift data.

So is young, bursty star formation a good model for distant galaxies with high specific star formation rates? To some extent, any answer depends on definitions. Degeneracies in the possible interpretation of photometric data make the 'age' of a galaxy difficult to constrain. Given the ultraviolet selection of LBGs (and analogues), they inevitably have a component of relatively young $(< 1$ Gyr) stars, but many also show evidence for an older, underlying stellar population (e.g. Eyles et al. [2005,](#page-8-28) [2007;](#page-8-29) Shapley et al. [2005;](#page-8-30) de Barros et al. [2014\)](#page-8-3). Since stellar ages are typically derived from continuum photometry, stretching into the rest-frame near-infrared, they reflect the population dominating the galaxy mass rather than the young stars (which drive nebular emission lines).

None the less, SED fitting of multiwavelength data for LBGs at z ∼ 3–6 produces typical ages of a few tens or hundreds of Myr, depending on the assumed star formation history (de Barros et al. [2014\)](#page-8-3). Shapley et al. suggest that most $z = 3$ galaxies may have undergone a very rapid early burst on time-scales of 50–100 Myr, before continuing to form stars at a significantly lower rate. In such a paradigm, the evolution of that early burst may dominate the nebular continuum at ages >300 Myr. While the SEDs of LBGs at $1.4 < z < 2.6$ (Reddy et al. [2012\)](#page-8-31) yield median stellar population ages of around 500 Myr – somewhat older than the populations discussed in Section 3 – near-infrared photometry of $z = 3$ sources yields a younger median age (∼320 Myr; Shapley et al. [2001\)](#page-8-32).

A recent study at intermediate redshifts, $2.0 < z < 2.6$, by Steidel et al. [\(2014\)](#page-8-7) has also concluded that both hard stellar radiation fields and a high-ionization parameter (i.e. low gas density with relative to number of ionizing photons) are required to reproduce the distribution of line ratios in their data. As Steidel et al. comment, our BPASS models generate a similar ionizing spectral continuum to their assumed blackbody source (with effective temperature $T_{\text{eff}} \sim 42000 \text{ K}$) and the inclusion of binary evolution pathways and stellar rotation is necessary to generate plausible ionizing spectra. We note that Steidel et al. also identified the 'extreme green pea' sample of Jaskot & Oey [\(2013\)](#page-8-33) as having comparable line ratios to their $z \sim 2.5$ sample, a property which appears to be shared by the galaxies we have selected as analogues to $z \sim 5$ star-forming galaxies.

It is likely that the hot, low-metallicity starburst with associated binary evolution presented here is not a unique explanation. As discussed in Section 3.2, models derived for continuous star formation at metallicity also recover high emission line ratios for a large fraction of their lifetime, although they show some tendency to overpredict the H β equivalent width. Some fraction of ongoing star formation, following an initial starburst would also likely recover similar line ratios. Any stellar population synthesis code necessarily explores a limited parameter set, exploring discrete stellar metallicities, interstellar gas densities, geometries and ages. By contrast real galaxies are a composite of many star-forming regions, each of different age and with different physical conditions. At best, a theoretical model can only be an approximate match.

The high line ratios in post-starburst conditions may well be diminished by combination with other stellar populations. It is also challenging to entirely rule out a weak AGN contribution to the emission lines, although we note that X-ray stacking analyses have constrained the moderate AGN fraction in the distant population to be <3 per cent at $z \sim 3$ (Laird et al. [2006\)](#page-8-34), inconsistent with the ubiquitous high line ratios. None the less, the ability of binary stellar population synthesis models to match the properties of distant galaxies over a reasonable, few 100 Myr, period is encouraging, and suggests that the hard ionizing spectra of these populations may plausibly play a significant role in the evolution of the ISM (and potentially IGM) at early times.

5 CONCLUSIONS

Our main conclusions can be summarized as follows.

(i) We measure the ionization-sensitive $[O \text{ III}]/H\beta$ ratio in ultraviolet-luminous local galaxies selected as z ∼ 5 Lyman break analogues. We find they lie well above the local average ratios for their SED-derived masses, with a median $[O \text{ III}]/H\beta = 3.36^{+0.14}_{-0.04}$, similar to those seen in high-redshift galaxy populations.

(ii) We consider the line ratios derived from the BPASS models. We determine that they can reproduce high [O III]/H $\beta \sim 3$ ratios for an extended period, at ages ∼50–300 Myr, at modest (0.2– $1.0 Z_{\odot}$) metallicities. They also accurately predict the behaviour of the [O II]/[O III] line ratio.

(iii) The density of the illuminated nebular gas appears to have only small effects on the predicted line ratios in binary stellar evolution models at moderate metallicities (0.4 Z_{\odot}) , and models can reproduce the data at densities seen in extragalactic star-forming regions. Single stellar models are rather more sensitive to gas density, with the predicted line ratios decreasing strongly with density at a given age.

(iv) While continuous star formation can generate similarly high line ratios at moderate metallicities, they struggle to reproduce the measured $H\beta$ luminosities and equivalent widths.

(v) We conclude that including binary population effects may be important when modelling stellar populations at <500 Myr, where line ratios depend sensitively on the evolution of the most massive stars.

Ideally, the comparison of multiple emission lines, or better still modelling of the full near-ultraviolet/optical spectrum, will be required to further constrain the star formation history and properties of distant galaxies. We plan to explore those of the local analogue population further in a forthcoming paper, in the hopes of gaining further insights into possible explanations of their ionizing spectra.

ACKNOWLEDGEMENTS

ERS acknowledges partial funding under STFC grant ST/L000733/1.

Based in part on public data from the SDSS DR7. Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the US Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS website is [http://www.sdss.org/.](http://www.sdss.org/) The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions.

Calculations were performed with version 8.0 of CLOUDY, last described by Ferland et al. [\(2013\)](#page-8-35).

REFERENCES

- Brinchmann J., Charlot S., White S. D. M., Tremonti C., Kauffmann G., Heckman T., Brinkmann J., 2004, MNRAS, 351, 1151
- Brinchmann J., Pettini M., Charlot S., 2008, MNRAS, 385, 769
- Bunker A. J. et al., 2010, MNRAS, 409, 855
- Contini M., 2014, A&A, 564, A19
- de Barros S., Schaerer D., Stark D. P., 2014, A&A, 563, A81
- Dopita M. A., Kewley L. J., Heisler C. A., Sutherland R. S., 2000, ApJ, 542, 224
- Eldridge J. J., Stanway E. R., 2009, MNRAS, 400, 1019
- Eldridge J. J., Stanway E. R., 2012, MNRAS, 419, 479
- Eldridge J. J., Izzard R. G., Tout C. A., 2008, MNRAS, 384, 1109
- Erb D. K., Shapley A. E., Pettini M., Steidel C. C., Reddy N. A., Adelberger K. L., 2006, ApJ, 644, 813
- Eyles L. P., Bunker A. J., Stanway E. R., Lacy M., Ellis R. S., Doherty M., 2005, MNRAS, 364, 443
- Eyles L. P., Bunker A. J., Ellis R. S., Lacy M., Stanway E. R., Stark D. P., Chiu K., 2007, MNRAS, 374, 910
- Ferland G. J., Netzer H., 1983, ApJ, 264, 105
- Ferland G. J., Korista K. T., Verner D. A., Ferguson J. W., Kingdon J. B., Verner E. M., 1998, PASP, 110, 761
- Ferland G. J. et al., 2013, Rev. Mex. Astron. Astrofis., 49, 137
- González V., Bouwens R. J., Labbé I., Illingworth G., Oesch P., Franx M., Magee D., 2012, ApJ, 755, 148
- Hainline K. N., Shapley A. E., Kornei K. A., Pettini M., Buckley-Geer E., Allam S. S., Tucker D. L., 2009, ApJ, 701, 52
- Heckman T. M. et al., 2005, ApJ, 619, L35
- Holden B. P. et al., 2014, preprint [\(arXiv:1401.5490\)](http://arxiv.org/abs/1401.5490)
- Hunt L. K., Hirashita H., 2009, A&A, 507, 1327
- Jaskot A. E., Oey M. S., 2013, ApJ, 766, 91
- Kewley L. J., Dopita M. A., Sutherland R. S., Heisler C. A., Trevena J., 2001, ApJ, 556, 121
- Kewley L. J., Dopita M. A., Leitherer C., Dave R., Yuan T., Allen M., Groves ´ B., Sutherland R., 2013, ApJ, 774, 100
- Laird E. S., Nandra K., Hobbs A., Steidel C. C., 2006, MNRAS, 373, 217
- Masters D. et al., 2014, ApJ, 785, 153
- Nakajima K., Ouchi M., Shimasaku K., Hashimoto T., Ono Y., Lee J. C., 2013, ApJ, 769, 3
- Pettini M., Shapley A. E., Steidel C. C., Cuby J.-G., Dickinson M., Moorwood A. F. M., Adelberger K. L., Giavalisco M., 2001, ApJ, 554, 981
- Reddy N. A., Pettini M., Steidel C. C., Shapley A. E., Erb D. K., Law D. R., 2012, ApJ, 754, 25
- Richard J., Jones T., Ellis R., Stark D. P., Livermore R., Swinbank M., 2011, MNRAS, 413, 643
- Schenker M. A., Ellis R. S., Konidaris N. P., Stark D. P., 2013, ApJ, 777, 67
- Shapley A. E., Steidel C. C., Adelberger K. L., Dickinson M., Giavalisco M., Pettini M., 2001, ApJ, 562, 95
- Shapley A. E., Steidel C. C., Erb D. K., Reddy N. A., Adelberger K. L., Pettini M., Barmby P., Huang J., 2005, ApJ, 626, 698
- Stanway E. R., Davies L. J. M., 2014, MNRAS, 439, 2474
- Stark D. P., Schenker M. A., Ellis R., Robertson B., McLure R., Dunlop J., 2013, ApJ, 763, 129
- Steidel C. C. et al., 2014, preprint [\(arXiv:1405.5473\)](http://arxiv.org/abs/1405.5473)
- Xia L. et al., 2012, AJ, 144, 28

This paper has been typeset from a TEX/LATEX file prepared by the author.

MNRAS **444,** 3466–3472 (2014)