

Sussex Research

Development and validation of the patient roles and responsibilities scale in cancer patients

Valerie Shilling, Rachel Starkings, Valerie Jenkins, David Cella, Lesley Fallowfield

Publication date

01-11-2018

Licence

This work is made available under the CC BY 4.0 licence and should only be used in accordance with that licence. For more information on the specific terms, consult the repository record for this item.

Document Version

Published version

Citation for this work (American Psychological Association 7th edition)

Shilling, V., Starkings, R., Jenkins, V., Cella, D., & Fallowfield, L. (2018). *Development and validation of the patient roles and responsibilities scale in cancer patients* (Version 1). University of Sussex. https://hdl.handle.net/10779/uos.23460026.v1

Published in

Quality of Life Research

Link to external publisher version

https://doi.org/10.1007/s11136-018-1940-2

Copyright and reuse:

This work was downloaded from Sussex Research Open (SRO). This document is made available in line with publisher policy and may differ from the published version. Please cite the published version where possible. Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners unless otherwise stated. For more information on this work, SRO or to report an issue, you can contact the repository administrators at sro@sussex.ac.uk. Discover more of the University's research at https://sussex.figshare.com/



Development and validation of the patient roles and responsibilities scale in cancer patients

Valerie Shilling¹ · Rachel Starkings¹ · Valerie Jenkins¹ · David Cella² · Lesley Fallowfield¹

Accepted: 16 July 2018 / Published online: 26 July 2018 $\ensuremath{\textcircled{O}}$ The Author(s) 2018

Abstract

Purpose The Patient Roles and Responsibilities Scale (PRRS) was developed to enable a broader evaluation of the impact of cancer and cancer treatment, measuring 'real world' roles and responsibilities such as caring for others and financial and employment responsibilities. Here, we report the development and initial validation.

Methods The 29-item PRRS was developed from the thematic analysis of two interview studies with cancer patients and caregivers. In the evaluation study, participants completed the PRRS alongside the Social Difficulties Inventory (SDI), the main criterion measure for concurrent validity, and the Functional Assessment of Cancer Therapy - General and WHO Quality of Life-BREF (WHOQOL-BREF) for additional convergent validity data. Questionnaires were completed at baseline, 7-days (PRRS only) and 2 months. Demographic data and patient characteristics were collected at baseline.

Results One hundred and thirty-five patients with stage III/IV breast, lung or gynaecological cancer or melanoma completed the PRRS at least once. Five items performed poorly and were removed from the scale. The final 16 core items selected comprised 3 dimensions: family well-being, responsibilities and social life, and financial well-being, identified in principal component analysis, accounting for 61.5% of total variance. Missing data (0.6%) and floor/ceiling effects were low (0%/1.5%). Cronbach's alpha was 0.9 for the PRRS-16; 0.79–0.87 for the subscales. PRRS showed good test–retest reliability (ICC-0.86), sensitivity to change and the predicted pattern of correlation with validation measures r = |0.65-0.77|. The standalone 7-item jobs and careers subscale requires further validation.

Conclusions Initial evaluation shows that the PRRS is psychometrically robust with potential to inform the evaluation of new treatments in clinical trials and real-world studies.

Keywords Cancer · Outcome measures · Psychometric performance · Validation · Questionnaire development

Background

More patients with cancer are living longer due in part to advances in molecular biology and the development of novel-targeted therapies [1]. There is growing recognition that assessment of the quality of this extended survival is

Electronic supplementary material The online version of this article (https://doi.org/10.1007/s11136-018-1940-2) contains supplementary material, which is available to authorized users.

important, namely how broader aspects of patients' lives are managed and affected. Quality of survival (QoS) looks beyond traditional endpoints of survival times, response and adverse events towards an understanding of the longer term patient experience. Topics include emotional concerns, financial burdens and impacts on role functioning [2]. Despite the extended survival benefits, associated side-effects from novel treatments can impact on a patient's ability to live a 'normal' life. With maintenance therapy, patients may face a trade-off between the putative extension and quality of that survival [3]. Uncertainties about likely survival benefit can leave patients and their families unable, or unwilling to make plans for the future [4].

Well-validated measures of financial and social distress exist (Comprehensive Score for Financial Toxicity [5, 6] and Social Difficulties Inventory [7, 8]), but there are currently no such measures that capture fully all aspects of the

Valerie Shilling v.m.shilling@sussex.ac.uk

¹ Sussex Health Outcomes Research and Education in Cancer (SHORE-C), Brighton and Sussex Medical School, University of Sussex, Falmer, Brighton, UK

² Department of Medical Social Sciences, Northwestern University Feinberg School of Medicine, Chicago, IL, USA

concept of QoS. The Quality of Life in Adult Cancer Survivors (QLACS) measure covers a number of these areas e.g. financial problems, family-related distress, but was developed specifically for use with long-term cancer survivors, defined by the authors as greater than five years from diagnosis [9]. Although the European Organisation for Research and Treatment of Cancer (EORTC) and the National Institute of Health (NIH) funded Patient Reported Outcomes Measurement Information System (PROMIS) initiatives have developed role functioning item banks for use in computer adaptive testing [10, 11], the impact of disease and treatment on roles and responsibilities is generally underrepresented in health-related patient reported outcome measures (PROMs). This is a serious oversight as many patients have caregiving responsibilities, for children, grandchildren, parents, partners or even pets. Qualitative studies demonstrate that these types of roles and responsibilities outside of work are a significant part of patients' lives, which cancer and its treatment can seriously impinge upon [12-16]. As role functioning issues are somewhat poorly described, many patients feel that information on how to cope at home is inadequate [17].

Our recent systematic reviews of PROMs currently in use identified areas of concern that are insufficiently researched, namely family, financial and occupational circumstances [18, 19]. Furthermore, QoS is not a static concept; it includes recognition that the relative importance of different dimensions will change across the disease trajectory. As such we designed the PROACT programme of work (Patient Reported Outcomes impact of Age and Carer role demands associated with Treatment) recognising that patients and their families are continually adjusting to a fluid situation whilst trying to maintain their 'real world' roles and responsibilities beyond cancer, such as caregiving responsibilities for a spouse or children/grandchildren, jobs and financial responsibilities [4].

Overview

PROACT is a multi-phase project, informed by the QoS conceptual framework, with the primary aim to develop and evaluate two self-report measures; one for patients and another for informal caregivers to assess comprehensively broader impacts of disease and treatment. The measures have been developed to be compatible with the Functional Assessment of Chronic Illness Therapy (FACIT) measurement system and are intended to benefit future patients and caregivers by capturing these wider impacts of cancer and its treatment during the evaluation of new treatments in clinical trials and could be used to aid clinical conversations around treatment and intervention.

To develop the PROACT measures, we conducted two systematic reviews [18, 19] to inform our understanding of the

content and psychometric properties of measures currently being used, and to identify gaps in relevant domains. Development of the two PROACT measures then followed three phases of scale development: item generation (study 1), item-reduction and scale construction (study 2) and initial scale evaluation and validation (study 3). Because the topics covered in the scale are not related to specific issues such as treatment side effects, rather to the broader impact such as time off work, it is intended that it will be useful for all groups of cancer patients. The development studies were conducted with patients with stage III/IV cancer and the person they nominated as their main source of support. However, the measures are intended to be suitable for all stages of cancer and are currently being validated in early stage disease. This paper reports the development and evaluation of the multi-dimensional Patient Roles and Responsibilities Scale (PRRS). The Caregiver Roles and Responsibilities Scale (CRRS) was developed concurrently and is reported elsewhere.

Ethics statement

Studies received ethics approval from London Queen Square Research Ethics Committee (ref: 15/LO/1323 Studies 1 and 2; ref: 16/LO/1125 Study 3). Signed informed consent was obtained from all participants.

Eligibility criteria

Inclusion/exclusion criteria were similar across all three studies. Eligible patients had advanced (stage III/IV) melanoma, lung or gynaecological cancer, with a requirement that they could nominate an informal caregiver also willing to take part. Both patients and caregivers were required to be over 18 years of age, have capacity to give informed consent and be able to read and speak English. Those who were currently inpatients or acutely distressed for any reason were excluded. In studies 2 and 3, we expanded the patient population to include women with breast cancer. The cancer types were selected to represent a range of potential cancer experiences in order that the measure might be sufficiently generic for other tumour sites. Study 3 did not require that both patient and caregiver consent to the study; either party could participate alone.

Scale development: methods and results

Study 1: item generation

We conducted in-depth qualitative interviews with 24 patients with advanced cancer and separate interviews with their nominated informal caregivers (N=23) about the impact of extended cancer survival on broader aspects

of life and well-being (see Supplementary File S1 for participant demographics). Topic guides were informed by our earlier systematic reviews [18, 19] and through discussion with advisors with a lived experience of cancer or who were supporting someone with cancer. Participants were first invited to generate and speak freely about topics important to their well-being and if and how these areas of life had been impacted by their/their loved one's cancer diagnosis and treatment. Further discussion was prompted (where necessary) around topics such as family life, relationships, leisure and social activities, employment and finances, physical and emotional health. Interviews were recorded (38.5 h of recorded data) and transcribed verbatim.

A thematic framework was developed from an initial process of open coding and tested iteratively as new data were collected. Thematic analysis identified 20 themes and 33 sub-themes from which a long list of 179 potential items was generated for the patient scale. These were reviewed for relevance and redundancy by the authors before 44 potential items were reviewed by our panel of advisors with lived experience. Consensus decisions on whether an item was retained or not were made through discussion informed by the thematic analysis of the interviews. Items were evaluated based on the relative importance the content appeared to have across interviews both in terms of the frequency with which the topic was discussed by participants and the significance attached to it.

Thirty items were retained for evaluation in Study 2. Where appropriate items already existed in the FACIT system, these validated items were used, rather than generating new ones. This included three items from the FACIT Comprehensive Score for Finacial Toxicity (COST) measure [5].

Interviews and thematic analysis were conducted by VS/ RS. The long list of potential items was generated by VS/ RS/VJ/LF and reviewed and reduced through discussion between VS/RS/VJ/LF and our panel of five advisors with lived experience.

Study 2: item reduction and scale construction

Cognitive interview techniques were employed to 'test' the potential questionnaire items. The processes participants applied when answering were explored, enabling us to check that the items were being understood as intended and consistently by different people, for example [20]. Cognitive interviews were conducted with a new cohort of 20 patients and informal caregivers using a mixture of the 'think aloud' technique (where participants verbalise the thought processes they go through in answering a questionnaire item) and specific probes around comprehension, retrieval, judgement and response options to assess each of the potential scale items. Example probes included: *What does that question mean to you, in your own words*? *What factors did you*

consider when you were deciding on your answer? Were you able to match the answer you wanted to give to one of the choices we've given you? Items were also discussed in terms of acceptability, relevance, redundancy and importance. Participants were also asked to identify missing topics. See Supplementary File S1 for participant demographics.

Scale items were revised, added and removed in an iterative fashion through the course of the study. Through this process 51 changes were made to the scale including 44 wording revisions, 4 deletions and 3 additions, resulting in a 29-item scale grouped conceptually into items around family and home life (N=13), financial well-being (N=8) and jobs and careers (N=8).

As with study 1, interviews and initial analysis were conducted by VS/RS with any changes to the scale in the course of the study agreed by consensus between VS/RS/ VJ/LJ after discussion of the interview content. Final scales were reviewed and agreed by VS/RS/VJ/LF and our panel of five advisors with lived experience.

Initial evaluation and validation: study 3

Methods

Population and procedure Participants were recruited from 11 sites in the United Kingdom, stratified by age group (\leq 50, 51–65, \geq 66 years) and tumour site (breast, gynaecological, lung, melanoma). All participants had stage III/IV cancer. Participants completed questionnaires at home either on paper or online, whichever their preference. Demographics and the full validation pack were completed at baseline, the PRRS was completed alone after 7 days (for test–retest reliability) and the full battery completed again after 2 months (for sensitivity to change).

Measures Participants completed the PRRS alongside the FACT-G, the SDI and the WHOQOL-BREF and provided basic demographic information such as age, employment status, level of education, relationship status and caregiving responsibilities along with information regarding diagnosis and treatment.

The PRRS The PRRS as completed in the validation study comprised 29 items: 21 core items formatted for the FACIT measurement system (responses as item applies to the past 7 days, 5 response options ranging from not at all—very much) and 1 binary response item on whether the participant has stopped work due to illness. A standalone scale, jobs and careers comprising 7 items is completed only by participants currently in paid employment (including those on sick leave). Negatively worded items were reverse scored so that a higher score corresponds to better quality of life. Where missing data occurred, total scores were prorated as long as more than half of the scale items were completed. The SDI [7] The SDI is a 21-item questionnaire developed for use in oncology practice. It covers a range of difficulties that might be experienced by patients in their everyday life and includes an everyday living subscale; money matters subscale and self and others subscale. There are five standalone items about difficulties around: sex, plans to have a family, living conditions, holidays and 'other'. Items are rated on a 4-point scale of no difficulty—very much (scored 0–3). For this analysis, we use the SD-16 scale which excludes the standalone items. The SD-16 generates a total score with range 0–44 (4 items are scored 0–2) with higher scores indicating greater social difficulty and also provides an established cut-off point to categorise patients as distressed/not distressed with the threshold set at SD-16 ≥ 10 [21].

The FACT-G [22] The FACT-G is a well-validated measure of health-related quality of life with 27 items comprised of four subscales: physical well-being (7 items), social well-being (7 items), emotional well-being (6 items) and functional well-being (7 items). Items are rated on a 5-point scale of not at all—very much (scored 0–4). Total score range is 0–108 with higher scores indicating higher quality of life.

Item GF7 "I am content with the quality of life right now" is used as the anchor variable to assess PRRS sensitivity to change.

The WHOQOL-BREF [23] The WHOQOL-BREF has 26-items producing a quality of life profile with four domain scores (physical health, psychological health, social relationships and environment) and two individually scored items about an individual's overall perception of quality of life and health. Each item is rated on a 5-point scale (scored 1–5). The four domain scores are scaled in a positive direction with higher scores indicating a higher quality of life.

Data analysis

Analyses were conducted using the Statistical Package for Social Sciences (IBM SPSS; version 23). Missing data from PRRS were managed by pro-rating total and were appropriate subscale scores. Missing data from other questionnaires were managed as per the specific instrument guidance. We used guidelines from the International Society for Pharmacoeconomics and Outcomes Research (ISPOR) [24, 25] and COSMIN (Consensus-based Standards for the selection of health Measurement Instruments) [26] in the development and evaluation of this measure.

Content validity

The PRRS was developed through extensive qualitative interviews in studies 1 and 2 and through collaboration with advisors with lived experience of cancer or caring for someone with cancer; no further content validation was undertaken.

Acceptability and precision

Acceptability was assessed by consideration of the amount of missing data overall and per item and the time to complete the scale. 15% missing data were set as the acceptable threshold for individual items.

Precision was assessed on a number of dimensions. Individual items with floor/ceiling effects > 70% were considered for removal as were those with a z-score of skewness >|4|. Individual items were also evaluated in terms of the pattern of correlation (items failing to show a pattern of correlation above 0.3 with multiple items or correlation with another item at \geq 0.8 were reviewed) and of corrected item-total correlation (items with CITC < 0.3 were reviewed). Items that performed poorly on any of these criteria were considered for removal from the scale.

Exploratory factor analysis

Principal components analysis with oblique rotation was performed to identify underlying factors. Bartlett's test of sphericity and the Kaiser–Meyer–Olkin (KMO) value were checked to confirm that the data were suitable for factor analysis. We applied the criterion of Eigenvalue > 1.0 and also examined the scree plot to determine the number of factors retained. Items were included in the factor on which they loaded highest (minimum accepted 0.4). The unidimensionality of subscales was assessed with corrected item-total correlations ($r \ge 0.3$ considered acceptable) and reliability with Cronbach's alpha. Missing values were managed using listwise deletion. Finally, we assessed whether the solution made conceptual and practical sense.

Internal consistency

Internal consistency was assessed using Cronbach's alpha (for good internal consistency we would expect $\alpha \ge 0.70$).

Criterion and convergent validity

Criterion validity was assessed by correlation with a commonly used legacy measure of the same concept, the SDI. We predicted a strong, negative correlation (r > -0.7). Convergent validity was further evaluated by correlating with other similar measures. We predicted a moderate to strong positive correlation (0.5 < r < 0.7) between PRRS scores and the FACT-G and WHOQOL-BREF.

Test-retest reliability

Test–retest reliability was assessed by comparing responses at baseline and at 7 days. Intraclass Correlations (ICCs; two-way random, absolute agreement) were calculated for total PRRS scores and sub-scale totals. An ICC > 0.7 is considered adequate, > 0.9, excellent. Weighted kappa scores (using linear weighting) were calculated for individual items; a score of $\kappa \ge 0.4$ was considered acceptable, $\kappa \ge 0.6$, good.

Sensitivity to change

PRRS change scores were calculated (T1–T3). Patients were categorised as improved, worsened or unchanged based on responses to the anchor question 'I am content with the quality of my life right now'. Paired *t* tests were used to determine if PRRS change within a group was significantly different from zero. Spearman correlation coefficients were calculated to explore the relationship between changes on the PRRS, anchor variable and other validation measures. We would expect moderate positive correlation (0.3 < r < 0.5) between change on the PRRS and the FACT-G and WHOQOL-BREF and moderate negative correlation between change on the PRRS and SDI (-0.3 < r < -0.5).

Results

Participants

One hundred and forty-three patients consented to take part, 135 completed baseline (37 breast, 35 gynaecological, 33 lung, 30 melanoma), 128 completed T2 and 118 completed T3. At baseline age ranged from 33 to 85 (median 61 years), 77% were female. See Fig. 1 for full details of patients approached, consented and completed questionnaires at each time point, along with reasons for decline/ drop out and Table 1 for key participant characteristics.

Acceptability of PRRS

Missing data rate for the PRRS was extremely low, 0.6% at baseline, 0.5% at Time 2 and 0.6% at Time 3. Missing data were distributed across 10 of the 28 Likert-scale items at baseline, 7 of the 28 items at Time 2 and 10 at Time 3 with no single item having more than 4 missing responses. The highest rate of missing data for a single item at any time point was 7% (an employment question with 26/28 possible responses); for core questions with the larger sample, the highest rate of missing data for a single item at any time point was 3% (threshold for investigation was >15% missing data).

At baseline, 60 participants completed questionnaires online, 75 completed on paper but only 61 recorded the time taken to complete the PRRS. Time to complete online ranged from 1.87 to 14.05 min with 1 extreme outlier at 34.78 min (mean excluding outlier=4.71 min, SD=2.21 min). On paper, time to complete ranged from 2 to 20 min with one extreme outlier at 30 min (mean excluding outlier=8.52 min, SD=4.22 min).

Remaining analyses and results are presented separately for the PRRS core items and the jobs and career subscale, which is only completed by participants currently in paid employment.

PRRS core items

Precision and item reduction

Five items had an unacceptably low level of correlation with other items and low (<0.3) corrected item-total correlations in reliability analysis. In addition, one of these did not meet the measure of sampling adequacy (set at 0.7). These five underperforming items were removed from further analysis. Four of the retained items had z-scores of skewness >|4|; however, their performance in terms of relationship to other items corrected item-total correlation and/or conceptual significance to the measure ensured their retention at this stage. No item correlated r > 0.8 with any other item.

None of the remaining individual items had missing data > 15% or floor effects exceeding 70% (range 1.5–41.5%, on 12/16 items less than 20% of responses were the minimum '0' response option). One item showed above threshold ceiling effects (74% 'my family have to help me financially', note this item is reverse scored; a response of 'not at all' receives a score of '4'). Scores of maximum '4' ranged from 7.4 to 74%; 4/16 items had less than 20% of responses at maximum '4'.

Total scores on the modified PRRS Core-16 (N=135) ranged from 10 to 64 (possible range 0–64), with mean, 39.20, SD 13.04, with skewness = -0.08 (SE = 0.204). No participants had minimum scores. 2 participants (1.5%) had maximum scores (threshold for ceiling effects set at 15%).

Exploratory factor analysis

Following listwise deletion based on all variables, N=130. Bartlett's test of sphericity was significant at <0.0001 and the KMO value was 0.850 confirming that the data were suitable for factor analysis. MSA for all variables was >0.7 suggesting adequate communality with other variables. 3 Eigenvalues were greater than 1 explaining 40.8, 10.4 and 10.3% of variance respectively (total variance explained 61.5%). The scree plot (Fig. 2) confirmed that 3 factors should be retained corresponding to family well-being




Fig. 1 Flow of participants through study

(Factor 1), financial well-being (Factor 2) and responsibilities and social life (Factor 3). All items except one loaded clearly on one factor. One item (PL9: "The way I see myself within the family has changed because of my illness") cross loaded on Factor 1 (0.433) and Factor 3 (-0.428). This item was retained for the family well-being subscale. The items and their loadings are shown in Table 2.

Reliability analyses were conducted for the three potential subscales using only the final items within the subscales. Cronbach's α for family well-being was

Table 1Key patientcharacteristics^a and PRRS totalscores^b

<1 year		N=135 PRRS		Univariate P			
Breast37 (27.5%)36.00 ± 14.380.060Gynacological35 (26%)40.07 ± 12.541Lung33 (24.5%)37.36 ± 11.281Melanoma30 (22%)44.13 ± 12.711Time since diagnosis41.03 ± 12.710.32940 (30%)36.61 ± 12.440.3291-2 years21 (15%)40.48 ± 15.2252 years> 2 years74 (55%)40.23 ± 12.690.003On treatment (missing N=22)0.00398 hole comparison (Bonferroni) old vsYes76 (67%)38.14 ± 13.9998 hole comparison (Bonferroni) old vsSqc group0.00398 hole comparison (Bonferroni) old vs51-6555 (41%)38.03 ± 12.689003> 5646 (34%)44.03 ± 13.079003SexFemale104 (77%)38.25 ± 13.520.121Male31 (23%)42.39 ± 10.099003Relationship statusFemale104 (77%)38.25 ± 13.520.121Male30 (22%)38.53 ± 13.288001Employment statusFemale104 (77%)38.26 ± 12.378001Grave up work due to illnes243.63 ± 12.318001Yes53 (39%)32.34 ± 10.76<0.0001No82 (61%)36.51 ± 12.3460001Grave up work due to illnes5540.01 ± 13.71Yes53 (39%)32.34 ± 10.76<0.0001No82 (61%)36.51 ± 12.346Grave up work due to illnes5 <t< th=""><th></th><th></th><th>$(Mean \pm SD)$</th><th></th></t<>			$(Mean \pm SD)$				
Gynaecological $35 (26\%)$ 40.07 ± 12.54 Lung $33 (24.5\%)$ 37.36 ± 11.28 Melanoma $30 (22\%)$ 44.13 ± 12.71 Time since diagnosis	Cancer type						
Lung $33 (24.5\%)$ 37.36 ± 11.28 Melanoma $30 (22\%)$ 44.13 ± 12.71 Time since diagnosis $<1 year$	Breast	37 (27.5%)	36.00 ± 14.38	0.060			
Manoma30 (2%) 44.13 ± 12.71 Time since diagnosis <td< td=""><td>Gynaecological</td><td>35 (26%)</td><td>40.07 ± 12.54</td><td></td></td<>	Gynaecological	35 (26%)	40.07 ± 12.54				
Time since diagnosis <1 year	Lung	33 (24.5%)	37.36 ± 11.28				
<1 year40 (30%)36.61 ± 12.440.3291-2 years21 (15%)40.48 ± 15.22>2 years74 (55%)40.23 ± 12.69On treatment (missing $N=22$)	Melanoma	30 (22%)	44.13 ± 12.71				
1-2 years21 (15%) 40.48 ± 15.22 $> 2 years74 (55%)40.23 \pm 12.69On treatment (missing N=22)Yes76 (67%)40.16 \pm 12.740.445No37 (33%)38.14 \pm 13.99Age group0.003\leq 5034 (25\%)34.54 \pm 11.73Post hoc comparison (Bonferroni) old vsyoung p = .003\geq 6646 (34%)44.03 \pm 13.07Post hoc comparison (Bonferroni) old vsyoung p = .003SexFemale104 (77%)38.25 \pm 13.520.121Male31 (23%)42.39 \pm 10.90Relationship statusPartner105 (78%)39.39 \pm 13.030.754No partner30 (22%)38.53 \pm 13.28Employment statusRetired70 (52%)42.60 \pm 12.520.001Sova up work due to illnessYes53 (39%)32.34 \pm 10.76<$	Time since diagnosis						
>2 years74 (55%) 40.23 ± 12.69 On treatment (missing $N=22$)Yes76 (67%) 40.16 ± 12.74 0.445 No37 (33%) 38.14 ± 13.99 0.003 Age group 0.003 Post hoc comparison (Bonferroni) old vs young $p=.003$ ≤ 50 $34 (25\%)$ 34.54 ± 11.73 Post hoc comparison (Bonferroni) old vs young $p=.003$ ≥ 66 $46 (34\%)$ 44.03 ± 13.07 Post hoc comparison (Bonferroni) old vs young $p=.003$ Sex $=$ $104 (77\%)$ 38.25 ± 13.52 0.121 Relationship status $=$ $104 (77\%)$ 38.25 ± 13.28 $=$ Partner $105 (78\%)$ 39.39 ± 13.03 0.754 $=$ No partner $30 (22\%)$ 38.53 ± 13.28 $=$ Employment status $=$ $=$ $=$ $=$ Retired $70 (52\%)$ 42.60 ± 12.52 0.001 Imployed/self employend $38 (28\%)$ 38.26 ± 12.37 $=$ Not in paid employment $27 (20\%)$ 31.69 ± 12.37 $=$ Gave up work due to illness $=$ $=$ $=$ Yes $53 (39\%)$ 32.34 ± 10.76 $<$ $<$ Interim 17 (13\%) 35.59 ± 10.48 $=$ $=$ Further $17 (35\%)$ 35.78 ± 12.55 $=$ $=$ Professional qualifications $33 (25\%)$ 41.52 ± 13.46 $=$ Caregiving responsibilities $=$ $=$ $=$ $=$ Yes $88 (65\%)$ 37.74 ± 12.79 0.076 $=$ No 4	<1 year	40 (30%)	36.61 ± 12.44	0.329			
On reatment (missing $N=22$) Yes 76 (67%) 40.16 ± 12.74 0.445 No 37 (33%) 38.14 ± 13.99 Age group 0.003 ≤ 50 $34 (25\%)$ 34.54 ± 11.73 Post hoc comparison (Bonferroni) old vs young $p = .003$ ≤ 66 $46 (34\%)$ 44.03 ± 13.07 Young $p = .003$ Sex Female $104 (77\%)$ 38.25 ± 13.52 0.121 Male $31 (23\%)$ 42.39 ± 10.90 Relationship status Partner $105 (78\%)$ 39.39 ± 13.03 0.754 No partner $30 (22\%)$ 38.53 ± 13.28 Bonferroni retired v not employed $p = .001$ Not in paid employed $38 (28\%)$ 38.26 ± 12.37 Bonferroni retired v not employed $p = .001$ No $82 (61\%)$ 32.34 ± 10.76 <0.0001 No $82 (261\%)$ 32.34 ± 10.76 <0.0001 No $82 (261\%)$ 35.59 ± 10.48 Yes $53 (39\%)$ 32.34 ± 10.76 <0.0001 No $82 (261\%)$ 35.59 ± 10.48 University $32 (24\%)$ $35.78 \pm 12.$	1–2 years	21 (15%)	40.48 ± 15.22				
Yes76 (67%) 40.16 ± 12.74 0.445 No37 (33%) 38.14 ± 13.99 0.003 Age group 0.003 ≤ 50 $34 (25\%)$ 34.54 ± 11.73 Post hoc comparison (Bonferroni) old vs young $p = .003$ $51-65$ $55 (41\%)$ 38.03 ± 12.68 $9ost hoc comparison (Bonferroni) old vsyoung p = .003\geq 6646 (34\%)44.03 \pm 13.077ost hoc comparison (Bonferroni) old vsyoung p = .003SexFemale104 (77\%)38.25 \pm 13.520.121Male31 (23\%)42.39 \pm 10.90754Relationship status8.25 \pm 13.520.001Partner105 (78\%)39.39 \pm 13.030.754No partner30 (22\%)38.53 \pm 13.28Employment statusRetired70 (52\%)42.60 \pm 12.520.001Employed/self employed38 (28\%)38.26 \pm 12.37Bonferroni retired v not employed p = .001No in paid employment27 (20\%)31.69 \pm 12.33Couo01Gave up work due to illnessYes53 (39\%)32.34 \pm 10.76<0.0001No82 (61\%)35.59 \pm 10.48Vinversity32 (24\%)35.78 \pm 12.55Professional qualifications33 (25\%)41.52 \pm 13.46VinversityYes88 (65\%)37.74 \pm 12.790.076No40 (30\%)37.78 \pm 14.030.413$	>2 years	74 (55%)	40.23 ± 12.69				
No 37 (33%) 38.14 \pm 13.99 Age group 0.003 \$50 34 (25%) 34.54 \pm 11.73 Post hoc comparison (Bonferroni) old vs youg $p=.003$ \$1-65 55 (41%) 38.03 \pm 12.68 Post hoc comparison (Bonferroni) old vs youg $p=.003$ \$266 46 (34%) 44.03 \pm 13.07 Post hoc comparison (Bonferroni) old vs youg $p=.003$ Sex Post hoc comparison (Bonferroni) old vs youg $p=.003$ Sex Post hoc comparison (Bonferroni) old vs youg $p=.003$ Sex Post hoc comparison (Bonferroni) old vs youg $p=.003$ Relationship status Post hoc comparison (Bonferroni old vs youg $p=.003$ Relationship status Post hoc comparison (Bonferroni Post hoc comparison (Bonferoni Pos	On treatment (missing $N=22$)						
Age group0.003≤5034 (25%) 34.54 ± 11.73 38.03 ± 12.68 Post hoc comparison (Bonferroni) old vs young $p = .003$ ≥6646 (34%) 44.03 ± 13.07 SexFemale104 (77%) 38.25 ± 13.52 0.121Male31 (23%) 42.39 ± 10.90 Relationship statusPartner105 (78%) 39.39 ± 13.03 0.754No partner30 (22%) 38.25 ± 13.28 0.001Employment statusEmployded/self employed $38 (28\%)$ 38.26 ± 12.37 Retired70 (52%) 42.60 ± 12.52 0.001Not in paid employment27 (20%) 31.69 ± 12.33 Bonferroni retired v not employed $p = .001$ No82 (61%) 43.63 ± 12.51 Education (missing $N = 2$)Secondary51 (38%) 40.91 ± 13.71 0.236 Further17 (13%) 35.59 ± 10.48 10.76University32 (24%) 35.78 ± 12.55 10.76Professional qualifications33 (25%) 41.52 ± 13.46 Caregiving responsibilitiesYes88 (65%) 37.74 ± 12.79 Yes88 (65%) 37.74 ± 12.79 0.076 No47 (35%) 41.92 ± 13.21 Caregiving responsibilities (for children/grandchildren only)Yes40 (30%) 37.78 ± 14.03 0.413	Yes	76 (67%)	40.16 ± 12.74	0.445			
≤50 34 (25%) 34.54 ± 11.73 Post hoc comparison (Bonferroni) old vs 51-65 55 (41%) 38.03 ± 12.68 $young p = .003$ ≥66 46 (34%) 44.03 ± 13.07 Sex Female 104 (77%) 38.25 ± 13.52 0.121 Male 31 (23%) 42.39 ± 10.90 Relationship status Partner 105 (78%) 39.39 ± 13.03 0.754 No partner 30 (22%) 38.53 ± 13.28 Bonferroni retired v not employed $p = .001$ Ketired 70 (52%) 42.60 ± 12.52 0.001 Employde/self employed 38 (28%) 38.26 ± 12.37 Bonferroni retired v not employed $p = .001$ Not in paid employment 27 (20%) 31.69 ± 12.33 Gave up work due to illness Yes 53 (39%) 32.34 ± 10.76 < 0.0001 No 82 (61%) 43.63 ± 12.51 Education (missing $N = 2$) Secondary 51 (38%) 40.91 ± 13.71 0.236 Further 17 (13%) 35.59 ± 10.48 State 12.55 Professional qualifications 33 (25%) 41.52 ± 13.46 Caregiving responsibilities	No	37 (33%)	38.14 ± 13.99				
25051(20)51(20)91(20)young $p = .003$ 51-6555 (41%)38.03 ± 12.68young $p = .003$ ≥6646 (34%)44.03 ± 13.07SexFemale104 (77%)38.25 ± 13.520.121Male31 (23%)42.39 ± 10.900Relationship statusPartner105 (78%)39.39 ± 13.030.754No partner30 (22%)38.53 ± 13.280.001Employment statusRetired70 (52%)42.60 ± 12.520.001Employed/self employed38 (28%)38.26 ± 12.37Bonferroni retired v not employed $p = .001$ Not in paid employment27 (20%)31.69 ± 12.33Gave up work due to illnessYes53 (39%)32.34 ± 10.76<0.0001	Age group			0.003			
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	≤50	34 (25%)	34.54 ± 11.73	· · · ·			
SexImage: second s	51–65	55 (41%)	38.03 ± 12.68	young $p = .003$			
Female $104 (77\%)$ 38.25 ± 13.52 0.121 Male $31 (23\%)$ 42.39 ± 10.90 Relationship statusPartner $105 (78\%)$ 39.39 ± 13.03 0.754 No partner $30 (22\%)$ 38.53 ± 13.28 0.001 Employment statusRetired $70 (52\%)$ 42.60 ± 12.52 0.001 Employed/self employed $38 (28\%)$ 38.26 ± 12.37 Bonferroni retired v not employed $p=.001$ Not in paid employment $27 (20\%)$ 31.69 ± 12.33 0.0001 Gave up work due to illness $$	≥66	46 (34%)	44.03 ± 13.07				
Male $31 (23\%)$ 42.39 ± 10.90 Relationship statusPartner $105 (78\%)$ 39.39 ± 13.03 0.754 No partner $30 (22\%)$ 38.53 ± 13.28 Employment statusRetired $70 (52\%)$ 42.60 ± 12.52 0.001 Employed/self employed $38 (28\%)$ 38.26 ± 12.37 Bonferroni retired v not employed $p=.001$ Not in paid employment $27 (20\%)$ 31.69 ± 12.33 Gave up work due to illnessYes $53 (39\%)$ 32.34 ± 10.76 <0.0001 No $82 (61\%)$ 43.63 ± 12.51 Education (missing $N=2$)Education (missing $N=2$) $Scondary$ $51 (38\%)$ 40.91 ± 13.71 0.236 Further $17 (13\%)$ 35.59 ± 10.48 $S.78 \pm 12.55$ Professional qualifications $33 (25\%)$ 41.52 ± 13.46 $S.78 \pm 12.55$ Caregiving responsibilities 37.74 ± 12.79 0.076 No $47 (35\%)$ 41.92 ± 13.21 4.413 Yes $88 (65\%)$ 37.74 ± 12.79 0.413 Yes $40 (30\%)$ 37.78 ± 14.03 5.413	Sex						
Relationship status 9artner 105 (78%) 39.39 ± 13.03 0.754 No partner 30 (22%) 38.53 ± 13.28 9 Employment status 70 (52%) 42.60 ± 12.52 0.001 Employed/self employed 38 (28%) 38.26 ± 12.37 Bonferroni retired v not employed $p=.001$ Not in paid employment 27 (20%) 31.69 ± 12.33 Gave up work due to illness Yes 53 (39%) 32.34 ± 10.76 <0.0001 No 82 (61%) 43.63 ± 12.51 Education (missing $N=2$) Secondary 51 (38%) 40.91 ± 13.71 0.236 Further 17 (13%) 35.59 ± 10.48 10.1026 University 32 (24%) 35.78 ± 12.55 10.48 Professional qualifications 33 (25%) 41.52 ± 13.46 10.236 Caregiving responsibilities 33 (25%) 41.52 ± 13.46 10.236 Ves 88 (65%) 37.74 ± 12.79 0.076 No 47 (35%) 41.92 ± 13.21 10.413 Yes 88 (65%) 37.78 ± 14.03 1.413	Female	104 (77%)	38.25 ± 13.52	0.121			
Partner105 (78%) 39.39 ± 13.03 0.754 No partner $30 (22\%)$ 38.53 ± 13.28 Employment statusRetired70 (52%) 42.60 ± 12.52 0.001 Employed/self employed $38 (28\%)$ 38.26 ± 12.37 Bonferroni retired v not employed $p = .001$ Not in paid employment27 (20%) 31.69 ± 12.33 Gave up work due to illnessYes53 (39%) 32.34 ± 10.76 <0.0001 No82 (61%) 43.63 ± 12.51 $<$ Education (missing $N = 2$)Secondary $51 (38\%)$ 40.91 ± 13.71 0.236 Further17 (13%) 35.59 ± 10.48 $<$ University $32 (24\%)$ 35.78 ± 12.55 $<$ Professional qualifications $33 (25\%)$ 41.52 ± 13.46 Caregiving responsibilities $<$ 0.076 $<$ No47 (35\%) 41.92 ± 13.21 $<$ Yes88 (65\%) 37.74 ± 12.79 0.076 No47 (35\%) 41.92 ± 13.21 $<$ Yes88 (05\%) 37.78 ± 14.03 $<$	Male	31 (23%)	42.39 ± 10.90				
No partner30 (22%) 38.53 ± 13.28 Employment status70 (52%) 42.60 ± 12.52 0.001 Retired70 (52%) 42.60 ± 12.52 0.001 Employed/self employed38 (28%) 38.26 ± 12.37 Bonferroni retired v not employed $p = .001$ Not in paid employment27 (20%) 31.69 ± 12.33 Gave up work due to illnessYes53 (39%) 32.34 ± 10.76 <0.0001 No82 (61%) 43.63 ± 12.51 Education (missing $N=2$)Education (missing $N=2$)Secondary $51 (38\%)$ 40.91 ± 13.71 0.236 Further17 (13%) 35.59 ± 10.48 University $32 (24\%)$ 35.78 ± 12.55 Professional qualifications33 (25%) 41.52 ± 13.46 Caregiving responsibilitiesYes88 (65%) 37.74 ± 12.79 0.076 No47 (35\%) 41.92 ± 13.21 UniversityYes80 (65%) 37.78 ± 14.03	Relationship status						
Employment status70 (52%) 42.60 ± 12.52 0.001Retired70 (52%) 42.60 ± 12.52 0.001Employed/self employed38 (28%) 38.26 ± 12.37 Bonferroni retired v not employed $p = .001$ Not in paid employment27 (20%) 31.69 ± 12.33 Gonferroni retired v not employed $p = .001$ Gave up work due to illnessYes $53 (39\%)$ 32.34 ± 10.76 <0.0001 No82 (61%) 43.63 ± 12.51 Education (missing $N=2$)Education (missing $N=2$)Secondary $51 (38\%)$ 40.91 ± 13.71 0.236 Further17 (13%) 35.59 ± 10.48 University $32 (24\%)$ 35.78 ± 12.55 Professional qualifications $33 (25\%)$ 41.52 ± 13.46 Caregiving responsibilitiesYes88 (65\%) 37.74 ± 12.79 0.076 No47 (35\%) 41.92 ± 13.21 0.413 Caregiving responsibilities (for children/grandchildren only)Yes $40 (30\%)$ 37.78 ± 14.03	Partner	105 (78%)	39.39±13.03	0.754			
Retired $70 (52\%)$ 42.60 ± 12.52 0.001 Employed/self employed $38 (28\%)$ 38.26 ± 12.37 Bonferroni retired v not employed $p = .001$ Not in paid employment $27 (20\%)$ 31.69 ± 12.33 Bonferroni retired v not employed $p = .001$ Gave up work due to illness $77 (20\%)$ 31.69 ± 12.33 0.001 Gave up work due to illness $45.3 (39\%)$ 32.34 ± 10.76 <0.0001 No $82 (61\%)$ 43.63 ± 12.51 <0.0001 No $82 (61\%)$ 43.63 ± 12.51 <0.236 Education (missing $N=2$) $51 (38\%)$ 40.91 ± 13.71 0.236 Further $17 (13\%)$ 35.59 ± 10.48 <0.236 University $32 (24\%)$ 35.78 ± 12.55 <0.001 Professional qualifications $33 (25\%)$ 41.52 ± 13.46 <0.076 No $47 (35\%)$ 41.92 ± 13.21 0.413 Ves $88 (65\%)$ 37.74 ± 12.79 0.413 Yes $40 (30\%)$ 37.78 ± 14.03 0.413	No partner	30 (22%)	38.53 ± 13.28				
Employed/self employed $38 (28\%)$ 38.26 ± 12.37 Bonferroni retired v not employed $p = .001$ Not in paid employment $27 (20\%)$ 31.69 ± 12.33 Bonferroni retired v not employed $p = .001$ Gave up work due to illness $27 (20\%)$ 31.69 ± 12.33 0.0001 Yes $53 (39\%)$ 32.34 ± 10.76 < 0.0001 No $82 (61\%)$ 43.63 ± 12.51 0.236 Education (missing $N=2$) $Secondary$ $51 (38\%)$ 40.91 ± 13.71 0.236 Further $17 (13\%)$ 35.59 ± 10.48 0.152 ± 13.46 University $32 (24\%)$ 35.78 ± 12.55 0.076 Professional qualifications $33 (25\%)$ 41.52 ± 13.46 Caregiving responsibilities $47 (35\%)$ 41.92 ± 13.21 0.413 Caregiving responsibilities (for children only) Ves $0.40 (30\%)$ 37.78 ± 14.03	Employment status						
Not in paid employment $27 (20\%)$ 31.69 ± 12.33 Gave up work due to illnessYes $53 (39\%)$ 32.34 ± 10.76 <0.0001 No $82 (61\%)$ 43.63 ± 12.51 Education (missing $N=2$)Secondary $51 (38\%)$ 40.91 ± 13.71 0.236 Further $17 (13\%)$ 35.59 ± 10.48 University $32 (24\%)$ 35.78 ± 12.55 Professional qualifications $33 (25\%)$ 41.52 ± 13.46 Caregiving responsibilitiesYes $88 (65\%)$ 37.74 ± 12.79 0.076 No $47 (35\%)$ 41.92 ± 13.21 0.413 Caregiving responsibilities (for children only)Yes $40 (30\%)$ 37.78 ± 14.03	Retired	70 (52%)	42.60 ± 12.52	0.001			
Gave up work due to illnessYes $53 (39\%)$ 32.34 ± 10.76 <0.0001 No $82 (61\%)$ 43.63 ± 12.51 Education (missing $N=2$)Secondary $51 (38\%)$ 40.91 ± 13.71 0.236 Further $17 (13\%)$ 35.59 ± 10.48 University $32 (24\%)$ 35.78 ± 12.55 Professional qualifications $33 (25\%)$ 41.52 ± 13.46 Caregiving responsibilities 37.74 ± 12.79 0.076 No $47 (35\%)$ 41.92 ± 13.21 Yes $88 (65\%)$ 37.78 ± 14.03	Employed/self employed	38 (28%)	38.26 ± 12.37	Bonferroni retired v not employed $p = .001$			
Yes $53 (39\%)$ 32.34 ± 10.76 < 0.0001 No $82 (61\%)$ 43.63 ± 12.51 Education (missing $N=2$)Secondary $51 (38\%)$ 40.91 ± 13.71 0.236 Further $17 (13\%)$ 35.59 ± 10.48 University $32 (24\%)$ 35.78 ± 12.55 Professional qualifications $33 (25\%)$ 41.52 ± 13.46 Caregiving responsibilitiesYes $88 (65\%)$ 37.74 ± 12.79 0.076 No $47 (35\%)$ 41.92 ± 13.21 0.413 Yes $40 (30\%)$ 37.78 ± 14.03	Not in paid employment	27 (20%)	31.69 ± 12.33				
No $82 (61\%)$ 43.63 ± 12.51 Education (missing N=2) 40.91 ± 13.71 0.236 Secondary $51 (38\%)$ 40.91 ± 13.71 0.236 Further $17 (13\%)$ 35.59 ± 10.48 University $32 (24\%)$ 35.78 ± 12.55 Professional qualifications $33 (25\%)$ 41.52 ± 13.46 Caregiving responsibilities $47 (35\%)$ 37.74 ± 12.79 0.076 No $47 (35\%)$ 41.92 ± 13.21 0.413 Caregiving responsibilities (for children only) $0.37.78 \pm 14.03$ 0.413	Gave up work due to illness						
Education (missing $N=2$)Secondary51 (38%) 40.91 ± 13.71 0.236 Further17 (13%) 35.59 ± 10.48 University32 (24%) 35.78 ± 12.55 Professional qualifications33 (25%) 41.52 ± 13.46 Caregiving responsibilitiesYes88 (65%) 37.74 ± 12.79 0.076 No47 (35%) 41.92 ± 13.21 0.413Caregiving responsibilities (for children only)9 $0.37.78 \pm 14.03$	Yes	53 (39%)	32.34 ± 10.76	< 0.0001			
Secondary $51 (38\%)$ 40.91 ± 13.71 0.236 Further $17 (13\%)$ 35.59 ± 10.48 University $32 (24\%)$ 35.78 ± 12.55 Professional qualifications $33 (25\%)$ 41.52 ± 13.46 Caregiving responsibilities 37.74 ± 12.79 0.076 No $47 (35\%)$ 41.92 ± 13.21 Caregiving responsibilities (for children/grandchildren only) 0.413 Yes $40 (30\%)$ 37.78 ± 14.03	No	82 (61%)	43.63 ± 12.51				
Further17 (13%) 35.59 ± 10.48 University $32 (24\%)$ 35.78 ± 12.55 Professional qualifications $33 (25\%)$ 41.52 ± 13.46 Caregiving responsibilities 37.74 ± 12.79 0.076 No $47 (35\%)$ 41.92 ± 13.21 Caregiving responsibilities (for children/grandchildren only) 0.413 Yes $40 (30\%)$ 37.78 ± 14.03	Education (missing $N=2$)						
University $32 (24\%)$ 35.78 ± 12.55 Professional qualifications $33 (25\%)$ 41.52 ± 13.46 Caregiving responsibilities 37.74 ± 12.79 0.076 No $47 (35\%)$ 41.92 ± 13.21 Caregiving responsibilities (for children/grandchildren only) 0.413 Yes $40 (30\%)$ 37.78 ± 14.03	Secondary	51 (38%)	40.91 ± 13.71	0.236			
Professional qualifications 33 (25%) 41.52 ± 13.46 Caregiving responsibilities $33 (25\%)$ 41.52 ± 13.46 Yes 88 (65%) 37.74 ± 12.79 0.076 No 47 (35%) 41.92 ± 13.21 0.413 Caregiving responsibilities (for children/grandchildren only) 0.413 0.413	Further	17 (13%)	35.59 ± 10.48				
Caregiving responsibilities 37.74 ± 12.79 0.076 No $47 (35\%)$ 41.92 ± 13.21 Caregiving responsibilities (for children/grandchildren only) 0.413 Yes $40 (30\%)$ 37.78 ± 14.03	University	32 (24%)	35.78 ± 12.55				
Yes88 (65%) 37.74 ± 12.79 0.076 No47 (35%) 41.92 ± 13.21 Caregiving responsibilities (for children/grandchildren only) 0.413 Yes40 (30%) 37.78 ± 14.03	Professional qualifications	33 (25%)	41.52 ± 13.46				
No $47(35\%)$ 41.92 ± 13.21 Caregiving responsibilities (for children/grandchildren only)0.413Yes $40(30\%)$ 37.78 ± 14.03	Caregiving responsibilities						
Caregiving responsibilities (for children/grandchildren only)0.413Yes40 (30%)37.78 ± 14.03	Yes	88 (65%)	37.74 ± 12.79	0.076			
children/grandchildren only) Yes $40 (30\%) 37.78 \pm 14.03$	No	47 (35%)	41.92 ± 13.21				
				0.413			
N_0 95 (70%) 39 79 + 12 63	Yes	40 (30%)	37.78 ± 14.03				
$33(10/0) 33.17 \pm 12.03$	No	95 (70%)	39.79 ± 12.63				

 $^{\mathrm{a}}\mathrm{Ethnicity}$ is not included as 96% of the population identified as White British

^bTotal scores on final modified scale, post item reduction

0.87 (N = 132, corrected item-total correlation range 0.60–0.76); for responsibilities and social life, 0.84 (N = 133, CITC range 0.50–0.71) and for financial

Fig. 2 Scree plot




Table 2 Item factor loading

	Item	Factor 1	Factor 2	Factor 3
PL1	My illness interferes with performing my responsibilities at home (e.g. cooking, cleaning, gardening, DIY)			-0.790
PL3	I am less able to fulfil my caregiving responsibilities (e.g. looking after children, grandchildren, another adult, pets)			-0.892
PL4	I have less patience for my caregiving responsibilities (e.g. looking after children, grandchildren, another adult, pets)			-0.674
PL5	I feel sad that my illness forces me to miss out on doing things with my children and/or other family members			-0.681
PL6	I worry about the impact of my illness on my partner (or the person who is my main support)	0.803		
PL7	I worry about the impact of my illness on my children and/or other family members	0.801		
PL8	I worry about the impact of my illness on people that I normally provide support to (e.g. friends, neighbours, parents and/or grandchildren)	0.774		
PL9	The way I see myself within the family has changed because of my illness	0.433		-0.428
PL11	I worry how my family will cope in the future	0.872		
PL13	I socialise less because of my illness			-0.653
FT3	I worry about the financial problems I will have in the future as a result of my illness or treatment		0.561	
FT11	I feel in control of my financial situation		0.500	
PF3	My family gives up things because of the financial impact of my illness		0.618	
PF4	My family have to help me financially		0.798	
PF5	The additional costs of my illness are more than I thought they would be (e.g. travel and parking, heating, healthy eating, supplements, non-prescription medication, paying for help at home)		0.686	
PF6	I have difficulty meeting the additional costs of my illness		0.874	

well-being 0.79 (N = 135, CITC range 0.47–0.71). Interitem and inter-subscale correlations are shown in Table 3.

Internal consistency

Cronbach's α for the modified PRRS total score at baseline (*N*=131, listwise deletion based on all variables) was 0.90 (CITC range 0.43–0.72); at T2 (*N*=124) α =0.90 (CITC range 0.37–0.73); and at T3 (*N*=112) α =0.91 (CITC range 0.43–0.76). In no case was α increased by the deletion of an item.

Criterion and convergent validity

As predicted, total scores on the PRRS correlated strongly and negatively with scores on the SDI (N = 135, r = -0.77). We did not find the floor effects reported elsewhere for SD-16 scores [27]; however, the distribution was skewed in that direction. As a precautionary measure, we categorised participants as distressed or not distressed based on a cutoff threshold of > 10/44 as recommended by the instrument authors [21]. 58 (43%) participants were categorised as distressed. We then used this dichotomous variable in a pointbiserial correlation with total PRRS score. As expected, the correlation was strong; N = 135, r = -0.65. Correlations between the PRRS subscales and most appropriately matched subscales of the SDI were also moderate to strong: PRRS responsibilities and social life with SDI everyday living r = -0.71; PRRS financial well-being with SDI money matters r = -0.65; PRRS family well-being with SDI self

Table 3	Inter-item	correlations	and	correlation	between	subscale scores
Table J	muci-num	contenations	anu	contenation	UCLWCCII	subscale scores

and others r = -0.52. PRRS scores showed the predicted correlations with FACT-G (N = 135, r = 0.65) and WHO-QOL-BREF (N = 135, r = 0.65).

Test-retest reliability

The median number of days between baseline and T2 was 8 (mode 7, range 3–21). Total PRRS scores showed good test–retest reliability (two-way random, absolute agreement) (N=128, ICC=0.86, 95% CI 0.80–0.90) as did the three subscales responsibilities and social life (N=128, ICC=0.83, 95% CI 0.77–0.88), family well-being (N=128, ICC=0.77, 95% CI 0.68–0.83) and financial well-being (N=128, ICC=0.86, 95% CI 0.81–0.90). Weighted kappa scores for individual items were all in the acceptable range ($\kappa \ge 0.4$) and ranged from 0.44 to 0.63.

Sensitivity to change

The median number of days between baseline and T3 was 66 (mode 60, range 56–117). Table 4 shows change over time in the anchor variable GF7 (I am content with the quality of my life right now). 47.5% of participants reported no change, 26.25% of participants reported an improvement by one or more points and the same proportion reported decline by one or more points. Participants whose GF7 score improved showed significant improvement in their PRRS scores (p = .008), those reporting decline on GF7 showed significant decline on PRRS (p = .001). As predicted, PRRS scores did not change significantly for those participants reporting no change on GF7 (p = .891). The groups do not differ at

Subscale		Range relati 8.38 (5.57) 0.475		Inter-item cor-	Correlations between subscale scores				
				relation range	Family well-being		Responsibilities and social life	Financial well- being	
Family well-being (5 items, possib	0.479–0.670 N=132								
Financial well-being (6 items, possible score 0–24)		12.85 (5.12 0–20		0.324-0.665 N=133	N=135		1.00		
		17.97 (5.01 5–24)	0.211–0.610 N=135			0.493 N=135	1.00	
Table 4 PRRS change by change on anchor variable GF7	I am content with the qu	ality of my	N	Baseline P		Mean PRRS	r	Effect size ^a	
(I am content with the quality of my life right now)	life right now			mean (SD)		change (SD)			
my me right how)	Improved by one point o		31	39.21 (13.8	,	-3.88 (7.58	,	-0.28	
	Unchanged		56	39.52 (13.0	67)	-0.12 (6.72	2) 0.891	-0.01	
	Worsened by one point of	r moro	31	40.58 (12.)	10)	4.55 (6.79	9) 0.001	0.38	

^aMean change score/SD mean baseline score

baseline. PRRS change scores correlated significantly with change on the anchor variable (r=0.42) and with change on other measures: FACTG (r=0.48) SDI (r=-0.42) and WHOQOL-BREF (r=0.32).

PRRS jobs and career subscale

The jobs and career subscale was completed by just 33, 30 and 28 patients (31, 29, 25 provided complete data) at the different time points, respectively; too few for reliability analyses. Table 5 shows baseline summary statistics for the items that make up the scale. The item PE3 had a particularly weak pattern of correlation with other subscale items (N=33, r=0.03-0.32) and two items (PE3 and PE7) demonstrated ceiling effects; however due to the small number of participants completing the scale, we will not eliminate items at this stage.

Discussion

We report the development and validation of the PRRS, a patient self-report measure of the impacts of cancer and treatment on family, financial and occupational roles and responsibilities. The format and structure are compatible with the FACIT measurement system. The scale was developed with advanced cancer patients and found to have very good reliability and validity in field testing. Although developed and validated on people with advanced cancer, the scale was developed with the intention to be sufficiently generic and appropriate for use in all stages of cancer. Indeed, due to the nature of the topics covered in the PRRS (i.e. family, finances, employment), there is potential for use in other chronic conditions. The scale is currently undergoing field testing with patients with early stage cancer. Extensive qualitative work to establish content validity would be required before any field testing in other chronic conditions would be possible. This should be a topic for further research.

The PRRS appears to be acceptable to participants both in terms of time to complete and the very low missing data rate in total and per item. Missing data on the final scale appear to be at random and are therefore assumed not to be indicative of a problem with any particular item. No participants recorded minimum scores on the final scale and only 2/135 recorded maximum scores.

Factor analysis of the Core-16 items revealed a threefactor solution corresponding to three potential subscales: family well-being, financial well-being, and responsibilities and social life. All demonstrated good internal consistency as does the PRRS-16 total score. Future studies will seek to confirm the factor structure and factorial invariance. The PRRS-16 total score showed the predicted strength of correlations with measures for criterion and convergent validity and good test-retest reliability. The scale demonstrated sensitivity to change; those patients showing improvement on the anchor variable showed significant improvement in PRRS scores, and those declining on the anchor variable showed significant decline in PRRS scores. Though significant, the effect size in both groups was small to medium. The magnitude of the effect is likely related to our choice of anchor variable and that we set just one point shift as the threshold for change on the anchor. As reported elsewhere [28] the absolute PRRS change was greater in those reporting worsening on the anchor variable than those reporting improvement.

We believe that there is utility in a total score for the PRRS, particularly when the scale is being used as part of a clinical discussion for straightforward monitoring of global change over time and as a way to quickly flag patients who might be experiencing difficulties and for whom further discussion and investigation might be appropriate. That said, the subscale scores may provide useful indication of how

Table 5 Baseline descriptive statistics for items on the jobs and career subscale (N=33)

	Item	Missing data (%)	Mean \pm SD	Median/mode	lz-score skewl	% floor (score 0)	% ceiling (score 4)
PE2	I have reduced my working hours because of my illness ^a	0	2.06 ± 1.66	3.0/0	0.57	33.3	24.2
PE3	My working hours are flexible to accommodate my treatment and appointments	0	3.00 ± 1.44	4.0/4	2.97	12.1	57.6
PE4	I feel I am able to do my job as well as I would like	0	2.30 ± 1.45	3.0/3	1.39	21.2	21.2
PE5	I worry that my illness will impact my employment in the future (including return to work) ^a	0	1.64 ± 1.34	2.0/2	0.76	27.3	12.1
FT9	I am concerned about keeping my job and income ^a	0	1.79 ± 1.52	2.0/0	0.38	30.3	18.2
PE6	I feel that my illness has limited my career ^a opportu- nities	0	2.48 ± 1.77	3.0/4	1.43	30.3	48.5
PE7	I feel supported by my employer	6.06	3.39 ± 1.12	4.0/4	4.21	3.2	71.0

^aItems are reverse scored

different domains are impacted for different individuals, allowing for more tailored approach to supportive interventions. Interpreting the profile of subscale scores rather than a total score is likely to be more appropriate and useful in some circumstances [29, 30].

We did not make a priori hypotheses on group differences on either the PRRS total score or scores on separate domains. We found that age was a significant factor with significant post hoc contrasts between those over 65 years and those 50 years or younger. Those patients already retired had significantly higher scores than those not in paid employment (but not those currently working) and those who had been forced to give up work due to illness had significantly lower scores than those who had not (either because they were not working or had continued to work). Scores for those who had caregiving responsibilities were lower than for those who did not, but not significantly so. A more nuanced exploration of the role of age and responsibilities to self and others is beyond the remit of this paper but will be presented elsewhere.

We indicated in the introduction that excellent, wellvalidated measures such as the SDI [7, 8] and COST [5, 6] exist. We believe that the PRRS provides a useful scale with more comprehensive coverage. For example, the SDI is not intended to provide a full assessment of jobs and careers; likewise the COST was not developed to measure family function and responsibilities. The PRRS has been developed following FACIT convention with the intention that it will sit within the FACIT measurement system and can be used alongside the FACT-G and or/other FACIT measures to provide comprehensive assessment.

Limitations

The main limitation relates to sample size for the validation study, particularly the jobs and careers subscale. While we have sufficient participant numbers for the factor and reliability analysis of the core questionnaire items, we were unable to evaluate the jobs and careers subscale fully. A further validation study is currently underway to address this shortfall.

We chose to develop the measure in patients with stage III/IV cancer; the stimulus for the programme of research being the increase in numbers of people living with cancer as a chronic condition. We also developed the measures in a limited number of tumour groups which although chosen to ensure a spread of key patient characteristics such as age, inevitably resulted in some sampling bias. For example two of the four cancer sites (breast and gynaecology) recruited only females resulting in a disproportionate number of female participants (77%). We did not have access to performance status scores for patients; this would be a useful group comparison in future research.

In any development study, choices are made regarding participant characteristics which limit the immediate generalizability of the findings. However, the PROACT measures were developed with an intention to be sufficiently generic for application to all disease stages and tumour types. Further validation will ensure that the measures are sufficiently generic for use in clinical trials and realworld studies.

Conclusion

The PRRS was developed to capture broad 'real life' impacts of cancer and its treatment and has potential use for both research and practice. For example, it may be used in the assessment of quality of life during the evaluation of new treatments in clinical trials and real world studies. It is potentially useful therefore to regulatory authorities scrutinising data from clinical trials and to policy makers when determining 'costs' of different treatments. The scale could also aid clinical conversations around treatment and intervention, with a growing body of evidence demonstrating the value of using PROMs in this way [31–35]. This initial evaluation shows that the PRRS is psychometrically robust with potential to inform the evaluation of novel therapies and to help drive the development of ameliorative interventions for the enhancement of extended survivorship.

Acknowledgements We would like to thank the patients, caregivers and clinic teams that participated in all of the PROACT research studies and our panel of patient and caregiver advisors. The PROACT studies were funded by Bristol-Myers Squibb.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- John, L., & Cowey, C. L. (2015). The rapid emergence of novel therapeutics in advanced malignant melanoma. *Dermatologic Therapy*, 5(3), 151–169.
- Fallowfield, L., et al. (2017). Quality of survival: A new concept framework to assess the quality of prolonged life in cancer. *Expert Review of Quality of Life in Cancer Care*, 2(4), 225–232.
- Fallowfield, L. J., et al. (2017). Therapeutic aims of drugs offering only progression-free survival are misunderstood by patients, and oncologists may be overly optimistic about likely benefits. *Support Care Cancer*, 25(1), 237–244.
- Shilling, V., et al., (2017) The pervasive nature of uncertainty: A qualitative study of patients with advanced cancer and their informal caregivers. *Journal of Cancer Survivorship*, 11(5), 590–603.

- 5. de Souza, J. A., et al. (2014). The development of a financial toxicity patient-reported outcome in cancer the COST measure. *Cancer*, *120*(20), 3245–3253.
- de Souza, J. A., et al. (2017). Measuring financial toxicity as a clinically relevant patient-reported outcome: The validation of the COmprehensive Score for financial Toxicity (COST). *Cancer*, *123*(3), 476–484.
- Wright, P., et al., (2008) Measurement and interpretation of social distress using the Social Difficulties Inventory (SDI). *European Journal of Cancer*, 44(11), 1529–1535.
- Wright, P., et al. (2011). The Social Difficulties Inventory (SDI): Development of subscales and scoring guidance for staff. *Psychooncology*, 20(1), 36–43.
- 9. Avis, N. E., et al. (2005). Assessing quality of life in adult cancer survivors (QLACS). *Quality of Life Research*, *14*(4), 1007–1023.
- Hahn, E. A., et al. (2014). New English and Spanish social health measures will facilitate evaluating health determinants. *Health Psychology*, 33(5), 490–499.
- 11. Gamper, E. M., et al. (2016). Development of an item bank for the EORTC role functioning computer adaptive test (EORTC RF-CAT). *Health and Quality of Life Outcomes, 14*, 72.
- Parrish, M. M., & Adams, S. (2003). An exploratory qualitative analysis of the emotional impact of breast cancer and caregiving among older women. *Care Management Journals*, 4(4), 191–197.
- 13. Hirschman, K. B., & Bourjolly, J. N. (2005). How do tangible supports impact the breast cancer experience? *Social Work in Health Care*, *41*(1), 17–32.
- 14. Buchbinder, M., Longhofer, J., & McCue, K. (2009). Family routines and rituals when a parent has cancer. *Families, Systems and Health*, 27(3), 213–227.
- 15. Bailey, E. H., et al. (2010). Impact of multiple caregiving roles on elevated depressed mood in early-stage breast cancer patients and same-age controls. *Breast Cancer Research and Treatment*, *121*(3), 709–718.
- Mackenzie, C. R. (2014). 'It is hard for mums to put themselves first': How mothers diagnosed with breast cancer manage the sociological boundaries between paid work, family and caring for the self. *Social Science & Medicine*, 117, 96–106.
- Jenkins, V., et al. (2013). Patients' and oncologists' views on the treatment and care of advanced ovarian cancer in the U.K.: Results from the ADVOCATE study. *British Journal of Cancer*, 108(11), 2264–2271.
- Catt, S., et al. (2017). Patient-reported outcome measures of the impact of cancer on patients' everyday lives: A systematic review. *Journal of Cancer Survivorship*, 11(2), 211–232.
- Shilling, V., et al. (2016) Patient-reported outcome measures for cancer caregivers: A systematic review. *Quality of Life Research*, 25(8), 1859–1876.
- Collins, D. (2003). Pretesting survey instruments: An overview of cognitive methods. *Quality of Life Research*, 12(3), 229–238.
- Wright, P., et al. (2007). Screening for social difficulties in cancer patients: Clinical utility of the Social Difficulties Inventory. *British Journal of Cancer*, 97(8), 1063–1070.

- Cella, D. F., et al. (1993). The Functional Assessment of Cancer Therapy scale: Development and validation of the general measure. *Journal of Clinical Oncology*, *11*(3), 570–579.
- Skevington, S. M., et al. (2004). The World Health Organization's WHOQOL-BREF quality of life assessment: Psychometric properties and results of the international field trial. A report from the WHOQOL group. *Quality of Life Research*, 13(2), 299–310.
- Patrick, D. L., et al. (2011). Content validity–establishing and reporting the evidence in newly developed patient-reported outcomes (PRO) instruments for medical product evaluation: ISPOR PRO good research practices task force report: Part 1– eliciting concepts for a new PRO instrument. *Value Health*, 14(8), 967–977.
- Patrick, D. L., et al. (2011). Content validity–establishing and reporting the evidence in newly developed patient-reported outcomes (PRO) instruments for medical product evaluation: ISPOR PRO Good Research Practices Task Force report: Part 2–assessing respondent understanding. *Value Health*, 14(8), 978–988.
- Mokkink, L. B., et al. (2010). The COSMIN checklist for assessing the methodological quality of studies on measurement properties of health status measurement instruments: An international Delphi study. *Quality of Life Research*, 19(4), 539–549.
- Ashley, L., et al. (2017) Health-related quality of life in cancer survivorship: Predictive power of the Social Difficulties Inventory. *Psychooncology*, 26(11), 1994–1997.
- Cella, D., Hahn, E. A., & Dineen, K. (2002). Meaningful change in cancer-specific quality of life scores: Differences between improvement and worsening. *Quality of Life Research*, 11(3), 207–221.
- Costa, D. S., et al. (2016). Psychometric properties of the fear of cancer recurrence inventory: An item response theory approach. *Psychooncology*, 25(7), 832–838.
- Costa, D. S., Smith, A. B., & Fardell, J. E. (2016). The sum of all fears: Conceptual challenges with measuring fear of cancer recurrence. *Support Care Cancer*, 24(1), 1–3.
- Greenhalgh, J. (2009). The applications of PROs in clinical practice: What are they, do they work, and why? *Quality of Life Research*, 18(1), 115–123.
- 32. Velikova, G., et al. (2008). The clinical value of quality of life assessment in oncology practice: A qualitative study of patient and physician views. *Psychooncology*, *17*(7), 690–698.
- Potter, C., et al. (2016). PROs for personalized care: Patient perspectives on potential use of the long-term conditions questionnaire (LTCQ) in clinical practice. *Quality of Life Research*, 25(Suppl 1), 188.
- Velikova, G., et al. (2004). Measuring quality of life in routine oncology practice improves communication and patient wellbeing: A randomized controlled trial. *Journal of Clinical Oncol*ogy, 22(4), 714–724.
- Yang, L. Y., et al. (2018). Patient-reported outcome use in oncology: A systematic review of the impact on patient-clinician communication. *Support Care Cancer*, 26(1), 41–60.