

Sussex Research

Feasibility randomised controlled trial of a one-day CBT workshop ("DISCOVER") for 15-18 year olds with anxiety and/or depression in clinic settings

Christina E Loucas, Irene Sclare, Daniel Stahl, Daniel Michelson

Publication date

09-06-2023

Licence

This work is made available under the Copyright not evaluated licence and should only be used in accordance with that licence. For more information on the specific terms, consult the repository record for this item.

Document Version

Accepted version

Citation for this work (American Psychological Association 7th edition)

Loucas, C. E., Sclare, I., Stahl, D., & Michelson, D. (2019). *Feasibility randomised controlled trial of a one-day CBT workshop ("DISCOVER") for 15-18 year olds with anxiety and/or depression in clinic settings* (Version 1). University of Sussex. https://hdl.handle.net/10779/uos.23465948.v1

Published in

Behavioural and Cognitive Psychotherapy

Link to external publisher version

https://doi.org/10.1017/S1352465819000286

Copyright and reuse:

This work was downloaded from Sussex Research Open (SRO). This document is made available in line with publisher policy and may differ from the published version. Please cite the published version where possible. Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners unless otherwise stated. For more information on this work, SRO or to report an issue, you can contact the repository administrators at sro@sussex.ac.uk. Discover more of the University's research at https://sussex.figshare.com/

Feasibility randomised controlled trial of a one-day CBT workshop ("DISCOVER") for 15-18 year olds with anxiety and/or depression in clinic settings

Christina E Loucas, Irene Sclare, Daniel Stahl, and Daniel Michelson

ABSTRACT

Background: "DISCOVER" one-day cognitive behavioural therapy (CBT) workshops have been developed to provide accessible, developmentally-sensitive psychological support for older adolescents experiencing emotional difficulties. Previous school-based evaluations of the DISCOVER model have shown positive outcomes.

Aims: The current study aimed to test the model for clinically-referred adolescents, in real-world settings.

Method: A randomised controlled trial (RCT) assessed feasibility, acceptability and preliminary outcomes of the DISCOVER intervention, in comparison with usual care, for 15-18-year-olds with emotional difficulties. Participants were recruited from outpatient clinic waiting lists in UK child and adolescent mental health services (CAMHS). Research feasibility indicators included rates of recruitment, randomisation, intervention participation (group workshops and individualised follow-up telephone calls), and data collection (at baseline and 8-week follow-up). Intervention acceptability was assessed using a structured service satisfaction questionnaire and semi-structured qualitative interviews with intervention participants. Preliminary clinical outcomes were explored using adolescent-reported validated measures of depression, anxiety and well-being.

Results: N=24 participants were randomised to intervention and usual care groups. Workshop attendance was good and high levels of treatment satisfaction were reported, although feasibility challenges emerged in recruitment and randomisation. Trends were found towards potential improvements in anxiety and well-being for the intervention group, but the effect estimate for depression was imprecise; interpretability was also limited due to the small sample size.

Conclusions: DISCOVER appears to be a feasible and acceptable intervention model for clinically-referred 15-18-year-olds with emotional difficulties. A full-scale RCT is warranted to evaluate effectiveness; protocol modifications may be necessary to ensure feasible recruitment and randomisation procedures.

Keywords

Adolescent; anxiety; depression; CBT; RCT

INTRODUCTION

Adolescence is a critical period for mental health. Approximately half of all mental disorders develop before the age of 15 years and 75% emerge by 18 years (Kim-Cohen et al., 2003). Emotional disorders are especially common in adolescence, when increased social challenges interact with immature systems for emotion regulation, posing heightened risks for both anxiety and depression (Ahmed, Bittencourt-Hewitt, & Sebastian, 2015; Steinberg, 2005). These syndromes, which are often comorbid, cause marked distress and impairment for approximately 4% of adolescents in the community, as well as accounting for most referrals to specialist child and adolescent mental health services (CAMHS) in the UK (Green, McGinnity, Meltzer, Ford, & Goodman, 2005; Wolpert et al., 2017).

Extensive evidence supports the use of psychological interventions to reduce the burden of youth mental disorders (Weisz et al., 2017), yet economic pressures limit the availability of evidence-based therapies and specialist mental health care more generally (Abdinasir & Pona, 2015). A recent review by the Children's Commissioner for England (2016) identified that only 1 in 250 young people were referred to CAMHS in 2015. Of these referrals, 28% were rejected outright, primarily because symptom presentations did not reach high thresholds for entry. For those young people who were accepted, large disparities were found in waiting times across geographic regions, ranging from 14 to 200 days. Notably, these figures do not incorporate "hidden waiting times" for the intervening period from initial assessment to treatment (Frith, 2016). Such delays can have devastating impacts on young people's quality of life and prospects (House of Commons Health Committee, 2014). For older adolescents, access is even more problematic due to poor transitions from youth to adult services, resulting in young people losing access to support during the period when they are most vulnerable (Memarzia, St Clair, Owens, Goodyer, & Dunn, 2015; Pona, Royston, Bracey & Gibbs, 2015).

Even when young people are seen in CAMHS, there is often limited provision of cognitive behavioural therapy (CBT) (Stallard, Udwin, Goddard & Hibbert, 2007; Edbrooke-Childs, Calderon, Wolpert, & Fonagy, 2015), which is widely recommended in evidence-based practice guidelines for youth anxiety and depression (Higa-McMillan, Francis, Rith-Najarian, & Chorpita, 2016; Hopkins, Crosland, Elliott, & Bewley, 2015; Weersing et al., 2017). The unwillingness (or inability) of many practitioners to implement evidence-based CBT protocols has been reported in multiple surveys (Hagermoser Sanetti, Collier-Meek, & Fallon, 2016). This is often related to a perceived lack of fit between empirically-tested (usually disorder-specific) structured treatments and the vagaries of routine practice (Southam-Gerow, Rodriguez, Chorpita, & Daleiden, 2012). "Transdiagnostic" protocols have been heralded as a way of addressing this implementation gap, offering a more parsimonious and flexible approach to dealing with real-world challenges of comorbidity and case complexity (Bearman & Weisz, 2015). Transdiagnostic interventions for anxiety and depression have been designed to target common mechanisms of emotion regulation implicated in both syndromes (Newby, McKinnon, Kuyken, Gilbody, & Dalgleish, 2015). Comorbidities can be targeted simultaneously in a single treatment protocol, therefore improving treatment efficiency, and in theory, improving access to psychological therapy (Chorpita, Taylor, Francis, Moffitt & Austin, 2004). Although direct head-to-head comparisons are lacking, between-study comparisons show that transdiagnostic CBT for adult emotional disorders is at least as effective as disorder-specific variants (Newby et al., 2015). Equivalent evidence is now emerging from transdiagnostic trials with young people (Chu, Temkin & Toffey, 2016). However, these studies have generally involved downward adaptations of emotion-focused transdiagnostic models developed with adults, which may give insufficient attention to adolescents' preferred outcomes, delivery formats, and key social contexts (Sclare & Michelson, 2016). Future developments require the active participation of young people in the conception, design and formative evaluation of new intervention models.

"DISCOVER" CBT workshops

The DISCOVER programme was born out of the need for more accessible and age-appropriate psychological support for distressed older adolescents. The delivery format and specific content of DISCOVER were co-created with a Teenage Advisory Group (TAG) to ensure that the social, emotional and relational needs of older adolescents were comprehensively addressed. The programme employs a one-day, group workshop format with individualised telephone follow-up, which has evolved from an established "well-being workshop" template (Brown, Cochrane, & Hancox, 2000). This corresponds to young people's preferences for more practical, interactive and less time-intensive modes of delivery (Persson et al., 2017; Plaistow et al., 2013).

DISCOVER follows a structured manual which covers numerous problem- and emotion-focused coping skills. These elements have been commonly applied in other evidence-based CBT interventions (Chorpita & Daleiden, 2009), and were subjected to further verification by the TAG, in order to improve fit with adolescents' perceptions of ecological relevance (Ng, Eckshtain & Weisz, 2016). The content is organised in a standardised sequence, which differs from certain other transdiagnostic programmes that deliver discrete modules according to individual participant needs (e.g. MATCH, Chorpita et al, 2013). Instead, workshop facilitators encourage participants to consider the suitability of different coping strategies from a curriculum of transdiagnostic self-regulatory skills. Further personalisation is enabled through the provision of follow-up telephone calls, which focus on progress towards individual participant goals.

The first iteration of DISCOVER was implemented in a variety of inner-London community venues as a stress management intervention for self-referred 16-18-year-olds, and tested in an uncontrolled pre-post cohort study (Sclare, Michelson, Malpass, Coster, & Brown, 2015). It was subsequently evaluated in a feasibility cluster randomised controlled trial (RCT) in secondary schools (Brown et al., 2019). The promising findings from these studies have raised the prospect of applying DISCOVER in other settings. Notably, the focus on common stressors and associated emotional symptoms is

consistent with transdiagnostic approaches previously used in clinical settings (Craske, 2012). We also identified a novel opportunity to evaluate DISCOVER for young people who had been placed on waiting lists after being referred to CAMHS with emotional difficulties. This aligned with key local and national service priorities to increase mental health service access and reduce waiting times (Department of Health & NHS England, 2015).

<u>Aims</u>

The present study aimed to investigate the feasibility, acceptability and preliminary outcomes of the DISCOVER intervention when applied to a clinical population of 15-18-year-olds with anxiety and/or depression, recruited from CAMHS waiting lists. The targeted age range was slightly wider relative to earlier evaluations of DISCOVER, based on an intention to broaden access within specialised services. A second aim was to examine the feasibility of undertaking a randomised controlled trial (RCT) of DISCOVER in CAMHS.

Objectives

The specific objectives were as follows:

- 1. To assess feasibility of intervention delivery, considering:
 - a. Attendance rate at DISCOVER workshops
 - b. Participation rate in telephone follow-up calls
- 2. To assess acceptability of intervention and associated trial procedures, considering:
 - a. Quantitative measures of service satisfaction
 - b. Qualitative feedback
- 3. To assess feasibility of research procedures, considering:
 - a. Number of eligible cases identified through CAMHS waiting lists
 - b. Consent rate
 - c. Randomisation rate
 - d. Data collection rates at baseline and 8-week follow-up

- 4. To obtain clinical outcome data to inform the design of a full-scale trial, considering:
 - Intervention effect estimates and confidence intervals, as indications of likely ranges
 for outcomes
 - b. Outcome variance estimates necessary for sample size calculations

METHODS

Design

We mounted a feasibility RCT with two parallel arms: an intervention arm (DISCOVER) and a control arm (usual care). Due to the study's feasibility objectives, an allocation ratio was not pre-specified, but determined by minimum group size requirements for DISCOVER workshops. Outcomes were assessed at baseline and 8-week follow-up.

Ethics

Ethical approval was obtained for the main study and a protocol modification (see below) by the London-Harrow NHS Research Ethics Committee (reference: 16/LO/0231). The trial was registered on clinicaltrials.gov (identifier: NCT02752945).

Participants

Eligibility criteria

Participant eligibility criteria were: (i) aged 15-18 years; (ii) fluent in English; (iii) currently on a CAMHS waiting list for specialist assessment/treatment, following a referral for anxiety and/or depression; (iv) willing and able to attend a DISCOVER workshop; and (v) position on the waiting list indicated that the young person would be unlikely to receive a CAMHS appointment within 8 weeks of completing a workshop. To determine criterion (iii), it was required that referral letters should indicate a need for assessment and/or treatment based on primary symptoms of anxiety and/or depression (although a confirmed diagnosis was not stipulated).

Young people were excluded if they were: (i) presenting with an acute risk of harm to themselves or others; (ii) presenting with severe learning difficulties; and/or (iii) unable to provide consent to participate. For exclusion criterion (iii), this included parental consent for participants aged 15 years.

Settings

The study was carried out in two outpatient CAMHS clinics in adjacent inner-London boroughs.

Compared with the general UK population, both boroughs are characterised by high levels of social disadvantage and a high proportion of black and minority ethnic residents (Stewart et al., 2009). The clinics provide a specialist multidisciplinary service for adolescents with a variety of mental health needs.

Sample size

A formal sample size calculation was not appropriate for our feasibility design. A recruitment target of N=30 was set, based on recommendations for obtaining outcome variance estimates for trial sample size calculations (Browne, 1995).

Interventions

DISCOVER

Structure

Previous iterations of DISCOVER (see Michelson, Sclare, Stahl, Morant, Bonin, & Brown, 2016) stipulated a minimum of four participants in each workshop and a maximum of 15. For the current study, the lower limit was increased to six participants. This modification was informed by the workshop facilitators' initial experience of delivering a small clinic-based group for young people. The larger group size was considered more conducive to open discussion and interaction in a clinical setting. Groups were co-facilitated by two doctoral-level clinical psychologists in accordance with a detailed manual. Workshops lasted for one day (6.5 hours) and were delivered in a CAMHS clinic on a weekday.

Content

The programme's content was rooted in cognitive-behavioural theory of emotional difficulties (Beck, 2011). CBT techniques (see Box 1) were introduced and practised through group discussion, role-play, individual written tasks and handouts. Video vignettes of teenage actors depicting common adolescent challenges were used to normalise experiences, stimulate discussion and enhance learning. Members received a workbook containing home-practice exercises and a summary of the workshop content.

Box 1 – DISCOVER workshop topics

- 1. Introductions and icebreakers
- 2. 'About stress' (psychoeducation)
- 3. 'The stress cycle' (basic CBT model thoughts, feelings, physical sensations and behaviour)
- 'Thoughts, different perspectives and thinking styles' (negative thinking patterns or biases)
- 'How to change your thinking and feel better' (distraction, thought challenging, mindfulness)
- 6. 'Behavioural changes' (graded exposure, problem-solving, time management)
- 7. 'Mind and body connections' (sleep hygiene, relaxation)
- 8. 'Tackling my problems' (goal setting and maintaining motivation)

Sclare et al. (2015)

Participants were invited to set personal goals at the end of the workshop. These were discussed approximately one week later in a 20-30-minute "telephone goal review" with one of the group

facilitators, aimed at monitoring and supporting progress. Participants were offered up to three additional reviews within the 8-week post-workshop period.

Usual care

Participants in both arms received usual care from CAMHS while on the waiting list. This comprised a clinic letter sent to the home of each participant, detailing local support services and emergency contacts in case of risk concerns. Participants were free to access any other treatment or professional support available to them outside CAMHS, including medication.

Measures

Feasibility

Please see study objectives for an outline of feasibility indicators.

Acceptability

At the end of the workshop, participants completed the 8-item Client Satisfaction Questionnaire (CSQ-8; Larsen, Attkisson, Hargreaves, & Nguyen, 1979). Items are summed to produce an overall treatment satisfaction score. We used an established categorisation system (Smith et al, 2014) to denote different aggregate satisfaction scores: poor (score of 8–13); fair (14–19); good (20–25); and excellent (26–32). Three open-ended questions were appended to the CSQ-8, exploring perceived helpfulness, suggestions for improvements and any other comments.

Participants in the intervention group were also invited to complete semi-structured exit interviews with a researcher (lead author) immediately after the follow-up outcome assessment. These 20-30-minute interviews followed a semi-structured topic guide, with questions and prompts that explored (i) views of the recruitment and assessment process; (ii) intervention content and structure; and (iii) perceived impact of the programme on well-being and service use. Interviews were audio-recorded and transcribed verbatim.

Clinical outcomes

Outcomes were assessed using three validated self-report measures at baseline (before randomisation) and 8-week follow-up. There were no changes to outcome measures after the trial commenced. The selected measures have been recommended by the UK's Child Outcomes Research Consortium (CORC; www.corc.uk.net) for routine use in CAMHS.

Primary outcome

The primary outcome was depression severity, measured by the 33-item Mood and Feelings

Questionnaire (MFQ) (Costello & Angold, 1988). This measure was particularly sensitive to change in an earlier evaluation of DISCOVER (Sclare et al., 2015). The clinical cut-off is ≥27, with higher scores indicating greater severity.

Secondary outcomes

Anxiety severity was assessed using the total anxiety score of the 47-item Revised Child Anxiety and Depression Scale (RCADS) (Chorpita, Yim, Moffitt, Umemoto & Francis, 2000). Raw scores were converted into standardised T-scores, with a clinical cut-off of ≥70. Higher scores represent greater symptom severity.

The 14-item Warwick-Edinburgh Mental Well-Being Scale (WEMWBS) (Clarke et al., 2011) was used to measure mental well-being. Higher scores represent better mental well-being.

Procedures

Recruitment

Clinicians in collaborating services screened waitlisted cases against study eligibility criteria, by scrutinising information from routinely available referral materials, with the aid of a structured proforma. The same clinicians then contacted potential participants by telephone and letter, enclosing a participant information sheet. Interested young people (or their parent/carer for younger adolescents) then opted-in to be contacted by a researcher. If agreeable, a meeting was

arranged at the clinic to obtain consent (including parallel consent from the parents/carers of 15year-olds) and to complete baseline assessments.

Randomisation and allocation concealment

Following baseline assessment, individual participant names were placed into individual opaque envelopes and sealed. Envelopes were given to an administrator, who generated the randomisation sequence by shuffling the envelopes and dividing into two piles according to the specified allocation ratio. Allocation ratios were determined by minimum workshop group size requirements. The two piles were given to the workshop facilitator or an assistant psychologist, who unsealed the envelopes and informed participants of their allocation. Workshop facilitators had no prior knowledge of participants.

Blinding

All follow-up assessments were completed by a researcher (first author) who was blind to group allocation.

Data analysis

Feasibility

Feasibility indicators were described using proportions and 95% confidence intervals.

Acceptability

CSQ-8 data were described using means and 95% confidence intervals. Qualitative data from open-ended questionnaire items and interview transcriptions were subjected to thematic analysis (Braun & Clarke, 2006). First, data from both sources were amalgamated and reviewed manually by the first author, with general annotations made for potential codes. Second, prominent features of the data were identified and initial codes were created. Third, codes were structured into emergent themes and associated sub-themes. Fourth, themes were inspected by a supervising co-author (DM) to certify that data extracts supporting each theme were meaningfully linked and different themes

could be clearly distinguished. Fifth, any discrepancies were deliberated and final refinements were made to themes and their definitions.

Data saturation was not formally monitored. However, in selecting an appropriate sample size, guidance from Guest and co-workers (2006) was followed, recommending that 6-12 interviews are sufficient to gain an understanding of common perceptions and experiences in a relatively homogenous sample.

Clinical outcomes

Given our primary focus on feasibility, the trial was not powered to detect significant differences. However, preliminary effects between groups were explored via one-way analysis of covariance (ANCOVA) for the clinical outcome measures, with baseline scores entered as covariates (Van Breukelen, 2006). Only data from those who completed both baseline and follow-up assessments were used (complete case analysis). Effect sizes were calculated as Cohen's d, where 0.20 was regarded as small, 0.50 medium and 0.80 large. Due to the small sample size and low power, emphasis was placed on confidence intervals (CIs; 95%) of the effect estimates over significance testing, allowing for examination of imprecision around effect sizes.


To provide estimates of the SD necessary for sample size calculations in a full-scale trial, 80% and 95% bootstrap CI estimates for the SD of baseline MFQ score (primary outcome) were calculated. The upper 80% CI is recommended for robust estimates of SDs (Browne, 1995).

RESULTS

Recruitment and participant flow

Three rounds of recruitment were completed from two clinics (round 1 in the first clinic and rounds 2 and 3 in the second clinic) between April and November 2016 (see Figure 1). Randomisation was unsuccessful in the first clinic (round 1) due to insufficient waitlisted cases. This resulted in a protocol amendment to reduce follow-up duration from 12 to eight weeks, thereby expanding the sampling frame available from existing CAMHS waiting lists.

Figure 1 – Participant flow



Overall, 97 waitlisted patients (round: 1, n=19; 2, n=37; 3, n=41) were eligible to take part. Of these, 28 were enrolled (round: 1, n=4; 2, n=12; 3, n=12). The sample size was too small (n=4) in the first round to permit randomisation; however, a decision was made by the Trial Management Group to deliver the workshop for these adolescents, based on clinical need. Participants in this workshop were removed from outcome analyses, but were included in estimates of feasibility.

In the second round of recruitment, n=12 participants were randomised in a 2:1 ratio, with eight participants allocated to the intervention arm and four to the control arm.

In the third round of recruitment, n=12 participants were initially randomised on a 2:1 ratio as above. However, two participants allocated to the intervention arm subsequently expressed uncertainty regarding workshop attendance. As a minimum group size of six was considered paramount to ensure a viable and acceptable group, a further decision was taken by the Trial Management Group to break randomisation and allocate an extra participant to the intervention group. This was achieved by randomly selecting a participant from the control group and reallocating to the intervention. Consequently, the allocation ratio was 3:1 (nine participants in the intervention group and three in the control group) in the third round of recruitment.

Outcomes

Feasibility of research procedures

Recruitment

Ninety-seven 15-18-year-olds on clinic waiting lists were identified as having prominent features of anxiety and/or depression, as per eligibility requirements. Thirty-one potential participants (32.0%; 95% CI = 22.9% to 42.2%) could not be contacted via the details provided in the referral, leaving 66 who were contactable (68.0%; 95% CI = 57.8% to 77.1%).

Consent

Of the remaining 66 contactable adolescents, 28 consented to take part (42.4%; 95% CI = 30.3% to 55.2%). Thirty-three adolescents declined; the two main reasons were other timetabled commitments, most commonly school (n=15); and apprehension about participating in a group (n=11). The remaining seven declining adolescents did not offer a reason. Five additional adolescents declined clinic contact entirely (i.e. they no longer needed support or were accessing another service) and were therefore discharged from the waiting list.

Baseline assessments

All 28 of the consenting participants completed baseline assessments (see Table 1 for participant characteristics).

Randomisation

Four consenting participants could not be randomised from the first round of recruitment due to insufficient sample sizes. Twenty-four participants were randomised in the next two rounds of recruitment (85.7%; 95% CI = 67.3% to 96%): 17 were randomised to the intervention arm and seven to the control arm. Randomisation was broken for one participant in the third recruitment cohort, such that a participant was randomly selected from the control group and re-allocated to the intervention (see justification above).

Follow-up assessments

Twenty out of 24 randomised participants (83.3%; 95% CI = 62.6% to 95.3%) completed follow-up assessments. Attempts were made to contact all randomised participants, but four could not be reached.

Feasibility of intervention delivery

Workshop attendance

Fifteen out of 17 participants in the intervention arm attended at least part of a DISCOVER workshop (88.2%; 95% CI 63.6% to 98.5%). One young person did not attend because of anxiety about participating in a group, and another young person failed to attend because of parental objections to involvement in mental health services. Another participant started, but did not complete the full workshop due to anxiety experienced in the group.

Telephone follow-up

Telephone goal reviews were completed with 11 out of 15 workshop attenders (73.3%; 95% CI 44.9% to 92.2%). Nine adolescents participated in one call, one received three calls and one received four calls; four could not be reached.

<u>Acceptability</u>

Satisfaction ratings

All 14 participants who completed the full workshop also completed the CSQ-8. The mean treatment satisfaction score was 26.86 (SD=3.88; 95% CI 24.82 to 28.89), with 13 out of 14 participants (93%; 95% CI 66% to 100%) indicating overall service satisfaction in the "good" or "excellent" range.

Qualitative feedback

Eleven participants took part in feedback interviews and 14 participants responded to open-ended questions from the CSQ-8. Participants were satisfied with and posed no suggestions about modifying procedures for initial approach, consent, questionnaire administration, randomisation and follow-up process. Data on treatment acceptability were categorised into three overarching themes reflecting different areas of perceived benefit (see Table 2 for participant quotes underpinning each theme). Suggestions for improvements have been described separately.

Being acknowledged

Participants appreciated the opportunity to access more immediate support and attention from professionals instead of remaining unseen and "forgotten" on a waiting list. The availability of telephone goal reviews also contributed to a sense of being kept in mind and validated as an individual whose needs are worthy of attention.

Valuing the group experience

Group interaction was highly valued for reasons such as finding comfort in shared difficulties, and shared learning of coping strategies. One participant expressed limited gain from the workshop content, but appreciated the group experience. There was a general appreciation of the group structure and dynamics being collaborative and facilitating engagement.

Developing improved ways of coping

Young people valued learning a range of practical coping strategies that they could explore before deciding which was most helpful for them. Participants particularly endorsed methods for reappraising problem orientation and habitual responses to stressors. Changing perspectives and responses tended to be linked to improvements in relationships and associated reductions in emotional distress.

Although there was consensus that DISCOVER had helped participants become better resourced for coping with stress, there was a felt need for further support to explore individual problems and consolidate learning. There was also concern about being taken off the waiting list, for fear of problems worsening and losing access to specialist care.

Suggestions for improvement

The most commonly suggested improvement was a shortened workshop programme, with several participants preferring a shorter duration to minimise fatigue. Other participants were concerned about losing content, leading to the suggestion of splitting the workshop across separate days.

Mixed views were expressed around the value of the video vignettes: some participants found the video characters relatable and helpful for modelling coping strategies, whereas others considered the scenarios depicted to be somewhat unrealistic. There were also reports of discomfort around the use of telephone calls for the goal reviews, with alternative methods of written communication (e.g. email or text messages) preferred.

Clinical outcomes

Intervention effect estimates and confidence intervals

As shown in Table 3, there were no conclusive differences in clinical outcomes between the intervention and control groups. However, trends were observed towards potential intervention effects on anxiety and mental-wellbeing. The effect estimate for depression was imprecise. Cls were wide and the presence of effects on primary and secondary outcomes cannot be ruled out at this stage.

Outcome variance estimates

Table 4 presents the estimates of SDs for future sample size calculations. The upper 80% CIs are recommended for any future power analyses of a full-scale trial.

DISCUSSION

The present study aimed to test the preliminary implementation of DISCOVER CBT workshops for waitlisted depressed and/or anxious adolescents in CAMHS using a feasibility RCT design. Although we observed specific challenges in participant recruitment and randomisation, rates of intervention participation and follow-up were encouraging. Moreover, the intervention was generally well-received by participants based on satisfaction scores and qualitative feedback. In terms of preliminary clinical outcomes, potential trends towards improvements favouring the intervention

group, were identified on some measures. Potential for impact was corroborated through participant exit interviews.

With respect to trial recruitment challenges, approximately one third of eligible patients could not be reached using contact information obtained from clinic registers. In real-world clinical settings, referral sources often provide incomplete or inaccurate information (Foot, Naylor & Imison, 2010; Gandhi et al., 2000). This effectively inflates the total sampling frame needed to reach an intended sample size.

Among those adolescents who were successfully contacted, the research consent rate (42%) was considerably lower than the benchmark of >80% reported in other psychological treatment trials with comparable populations (e.g. Chapman et al., 2016; Goodyer et al., 2017). Non-participation primarily related to concerns about the group format and timing of workshops. Pre-group preparation (e.g. clarifying treatment expectations and outlining group rules) has been recommended for alleviating anxieties before embarking on any group-based psychological intervention (Bernard et al., 2008). This would be a useful avenue to explore for boosting uptake in further clinical evaluations of DISCOVER. Relevant modifications could include the use of online videos as an engaging and accessible method for socialising potential participants to the workshop model (Campinha-Bacote & Dexter, 2012), especially considering that the DISCOVER curriculum already makes extensive use of video materials.

With regards to timetabling, there was an inevitable trade-off between the efficiency of a one-day group intervention and scheduling restrictions, such that the group workshops could not be planned around the schedules of individual participants. Moreover, the potential for delivering community-based DISCOVER workshops on non-school days has previously received limited support (Sclare et al, 2015). However, young people in other studies have expressed a desire for greater flexibility from mental health services (Abdinasir, 2017); hence there may be value in exploring the delivery of clinic-based DISCOVER workshops on non-school days.

The above-mentioned recruitment barriers meant that randomisation was not feasible in the first round of recruitment, and was broken in the third. One option for expanding the sampling frame in a future trial would involve extending eligibility criteria to include young people with impairing but subthreshold symptoms, who might otherwise be denied access to specialist mental health care (Balázs et al., 2013; Children's Commissioner for England, 2016). This would offer a pragmatic way to boost sample size, while building service capacity to treat young people at an earlier and less severe stage in their mental health presentation. From a service provider perspective, there is also a direction towards scaling-up innovative early interventions for youth mental health, in line with policy initiatives and other drivers of service redesign (Department of Health & NHS England, 2015; McGorry, Bates & Birchwood, 2013).

Participation across subsequent phases of the study flow was encouraging. The rate of "loss to follow-up" (17%) was within the range of 15-20% missing data commonly reported in psychological and educational research (Enders 2003, as cited in Dong & Peng, 2013), while the intervention dropout rate (11.8%) compared favourably with the benchmark of 18.5% identified in a recent meta-analysis of pre-treatment drop-out from CBT in child and adolescent populations (Fernandez, Salem, Swift, & Ramtahal, 2015). Participants engaged well with the goal review follow-up calls, with 73.3% of workshop attenders completing at least one call. This is similar to the rate of 78.8% reported in a previous evaluation of DISCOVER (Brown et al., 2019). In the qualitative feedback, participants described positive aspects of the calls, such as feeling acknowledged, but also advocated for alternative written communication methods in preference to telephone conversations. This mirrors previous findings that telephone calls may be perceived as less acceptable by adolescents relative to other modes of mental health care delivery (Bradford & Rickwood, 2012).

Both quantitative and qualitative feedback suggested high levels of acceptability for the workshops. All but one participant in the intervention arm rated overall service satisfaction as good or excellent, and the mean satisfaction score (M=26.86; SD=3.88) was similar to CSQ-8 scores in other trials involving more conventional CBT formats, such as 12-session individual CBT for youth with anxiety (M=26.0; SD=4.5; Khanna & Kendall, 2010) and depression (M=26.75; SD=4.19; Shirk, DePrince, Crisostomo, & Labus, 2014). Within the workshop format, interactions with fellow group members and facilitators were highly valued, consistent with other CAMHS research citing collaboration and reduction of power differentials as crucial for meeting the developmental needs of older adolescents (Harper, Dickson & Bramwell, 2014).

Considering the DISCOVER service model more generally, participants appreciated being able to access more immediate support while on a waiting list in CAMHS. Comments were made by some participants about a need for mental health care after ending their involvement with DISCOVER. This would be consistent with a stepped-care model, where an initial "low-intensity" (less costly and less time-intensive) psychological intervention is followed by another treatment of incremental intensity (Clark, 2011).

We also assessed clinical outcomes at baseline and 8-week follow-up. Findings were inconclusive for all outcomes, however trends were observed towards potential intervention effects on anxiety and well-being, while the effect on depression was imprecise. We reiterate that the study was not designed to provide reliable estimates of effectiveness and caution should be applied in interpreting outcomes derived from small sample sizes (Ioannidis, 2005). Nevertheless, the observed pattern of results raises important questions about the transdiagnostic effects of the intervention. Around 70% of the total sample had been referred to CAMHS for depression (with around half of these participants presenting with comorbid anxiety), while the overall rate of anxiety was slightly lower at around 60%. Hence, the observed lack of impact on depression is not obviously explained by low baseline prevalence. Pending replication of this finding, it is possible that treatment elements of

more specific relevance to depression may need to be enhanced in any future modifications of the programme. For example, this may call for a greater emphasis on behavioural activation, which is a well-established component of CBT for depression (Weersing, Rozenman, Gonzalez, 2009).

Measuring depression and anxiety as co-primary outcomes could assist with better understanding the relative effects in a future definitive trial. Lastly, we must note a recent meta-anaysis of 447 trials of psychotherapy for youth internalising problems, which found strongest post-treatment effects for anxiety (0.61) and weakest effects for depression (0.29) (Weisz et al., 2017). Hence, optimising treatments for depression appears to be a key issue for the field of youth psychotherapy as a whole.

Limitations

The feasibility design and relatively small sample size necessarily limit the conclusions that can be drawn from this trial. Although we observed trends towards certain improved outcomes favouring the intervention, the study was underpowered and the effect size estimates included wide confidence intervals.

Potential allocation bias was also introduced due to randomisation being compromised. Additionally, selection bias cannot be ruled out, considering that only around one-quarter of all eligible participants in the study's sampling frame were successfully contacted and consented. Formal assessment of selection bias could not be undertaken given the relatively sparse clinical and demographic information available from referral letters. Lastly, interviews could not be undertaken with the two participants who attended the workshop but did not complete follow-up assessment. Hence, bias from attrition might have been introduced, as interviews were not conducted with participants who potentially had less favourable experiences of the intervention.

A final limitation pertains to the qualitative analysis. Codes were ordered into thematic categories in consultation with a senior co-author, although the initial codes were derived without independent

checks. Having two researchers involved in all aspects of the qualitative analysis might have enhanced the reliability of findings.

Recommendations for future research

The present feasibility trial has demonstrated that a full-scale RCT of DISCOVER is warranted in specialist youth mental health clinics, building on existing studies of DISCOVER in self-referred community samples (Brown et al., 2019; Sclare et al., 2015). Our feasibility data point towards short-term improvements in participants' emotional functioning after participating in the DISCOVER workshop, along with an expressed need for additional therapeutic support to consolidate therapeutic gains. A future trial of DISCOVER would provide more definitive evidence about the potential sensitizing effects of the intervention within a stepped-care framework. This should incorporate protocol refinements to enhance the likely recruitment rate, and compare prospective rates of service use, as well as clinical outcomes for adolescents who received either DISCOVER or usual care. A parallel economic analysis would allow for the assessment of potential cost efficiencies stemming from reductions in waiting times and duration of care.

Clinical implications

Just under half of all young people who were contactable opted-in to the study, suggesting significant demand for a brief psychological intervention, that can be accessed promptly by adolescents who would otherwise be waiting for usual care. Issues around service capacity are particularly salient for older adolescents who often "fall through the gap" between adult and child services, failing to access treatment at a crucial developmental stage (Memarzia et al., 2015; Pona et al., 2015). The specific innovations of DISCOVER (e.g. age-appropriate content; interactive, video-supported group workshop format) serve as a promising platform for delivering a brief, developmentally-attuned, first-line transdiagnostic intervention for clinic-referred adolescents with emotional difficulties. Nevertheless, there is scope for further optimisation of this intervention model, such as providing participants with preparatory information to address concerns about the group workshop format; and offering a choice about the preferred mode of contact for 1:1 follow-up

sessions, rather than limiting this to telephone calls. The potential for introducing focal treatment elements that are specific to anxiety and depression also warrants consideration.

Conclusion

The current study provides preliminary evidence for the feasibility and acceptability of delivering the DISCOVER one-day CBT workshop intervention for waitlisted, clinic-referred 15-18-year-olds with emotional difficulties, while establishing key research parameters needed to design a full-scale trial. Future evidence on clinical effectiveness and service-level outcomes (including associated direct and indirect costs) will help to determine whether DISCOVER can be scaled up to provide accessible, age-appropriate and cost-effective mental health care for the high volume of adolescents presenting to specialist services with anxiety and depression.

REFERENCES

- Abdinasir, K. (2017). Stick with us: Tackling missed appointments in children's mental health services.

 London: The Children's Society. Retrieved from

 https://www.childrenssociety.org.uk/sites/default/files/stick with us tackling missed appointments in children s mental health services.pdf
- Abdinasir, K. & Pona, I. (2015). *Access denied: A teenager's pathway through the mental health system.* London: The Children's Society. Retrieved from http://www.childrenssociety.org.uk/sites/default/files/AccessDenied_final.pdf
- Ahmed, S. P., Bittencourt-Hewitt, A., & Sebastian, C. L. (2015). Neurocognitive bases of emotion regulation development in adolescence. *Developmental Cognitive Neuroscience*, *15*, 11-25. https://doi.org/10.1016/j.dcn.2015.07.006
- Balázs, J., Miklósi, M., Keresztény, A., Hoven, C. W., Carli, V., Wasserman, C., ...Wasserman, D.
 (2013). Adolescent subthreshold-depression and anxiety: psychopathology, functional impairment and increased suicide risk. *Journal of Child Psychology and Psychiatry*, *54*(6), 670-677. doi: 10.1111/jcpp.12016.
- Bearman, S. K., & Weisz, J. R. (2015). Review: Comprehensive treatments for youth comorbidity evidence-guided approaches to a complicated problem. *Child and Adolescent Mental Health,* 20(3), 131-141. doi: 10.1111/camh.12092
- Beck, J. S. (2011). *Cognitive Behavior Therapy: Basics and Beyond* (2nd ed.). New York: Guilford Press.
- Bernard, H., Burlingame, G., Flores, P., Greene, L., Joyce, A., Kobos, J. C., ...Feirman., D. (2008).

 Clinical practice guidelines for group psychotherapy. *International Journal of Group*Psychotherapy, 58(4), 455-542. doi: 10.1521/ijgp.2008.58.4.455
- Bradford, S., & Rickwood, D. (2012). Adolescent's preferred modes of delivery for mental health services. *Child and Adolescent Mental Health*, *19*(1), 39-45. doi: 10.1111/camh.12002

- Braun, V. & Clarke, V. (2006). Using thematic analysis in psychology. *Qualitative Research in Psychology*, *3*(2), 77-101. http://dx.doi.org/10.1191/1478088706qp063oa
- Brown, J. S. L., Blackshaw, E., Stahl, D., Fennelly, L., McKeague, L., Sclare, I., & Michelson, D. (2019).

 School-based early intervention for anxiety and depression in older adolescents: A feasibility randomised controlled trial of a self-referral stress management workshop programme

 ("DISCOVER"). *Journal of Adolescence*, 71, 150-161.

 https://doi.org/10.1016/j.adolescence.2018.11.009
- Brown, J. S. L., Cochrane, R., & Hancox, T. (2000). Large-scale health promotion stress workshops for the general public: A controlled evaluation. *Behavioural and Cognitive Psychotherapy, 28*(2), 139-151. https://www.cambridge.org/core/journals/behavioural-and-cognitive-psychotherapy/article/large-scale-health-promotion-stress-workshops-for-the-general-public-a-controlled-evaluation/91D41A622BACB46CEA853889A9CCA4C8
- Browne, R.H. (1995). On the use of a pilot sample for sample size determination. *Statistics in Medicine*, *14*(17), 1933-1940. doi: 10.1002/sim.4780141709
- Campinha-Bacote, D., & Dexter, D. (2012). *Pre-group preparation in college counseling centers:*Through the use of an audio-visual aid. (Electronic Thesis or Dissertation). Retrieved from https://corescholar.libraries.wright.edu/cgi/viewcontent.cgi?article=2249&context=etd all
- Chapman, R., Loades, M., O'Reilly, G., Coyle, D., Patterson, M., & Salkovskis, P. (2016). 'Pesky gNATs': investigating the feasibility of a novel computerized CBT intervention for adolescents with anxiety and/or depression in a Tier 3 CAMHS setting. *The Cognitive Behaviour Therapist, 9*, e35. https://doi.org/10.1017/S1754470X16000222
- Children's Commissioner for England. (2016). *Lightning review: access to child and adolescent*mental health services, May 2016. London: Children's Commissioner for England. Retrieved

- from https://www.childrenscommissioner.gov.uk/wp-content/uploads/2017/06/Childrens-commissioners-Mental-Health-Lightning-Review.pdf
- Chorpita, B.F., & Daleiden, E.L. (2009). Mapping evidence-based treatments for children and adolescents: application of the distillation and matching model to 615 treatments from 322 randomized trials. *Journal of Consulting and Clinical Psychology*, 77(3), 566-579. doi: 10.1037/a0014565.
- Chorpita, B. F., Taylor, A. A., Francis, S. E., Moffitt, C., & Austin, A. A. (2004). Efficacy of modular cognitive behavior therapy for childhood anxiety disorders. *Behavior Therapy*, *35*(2), 263-287. https://doi.org/10.1016/S0005-7894(04)80039-X
- Chorpita, B. F., Yim, L. M., Moffitt, C., Umemoto L. A., & Francis, S. E. (2000). Assessment of symptoms of DSM-IV anxiety and depression in children: A revised child anxiety and depression scale. *Behaviour Research and Therapy*, *38*(8), 835-855.

 https://doi.org/10.1016/S0005-7967(99)00130-8
- Chorpita, B. F., Weisz, J.R., Daleiden, E.L., Schoenwald, S.K., Palinkas, L.A., Miranda, J., ...Gibbons, R.D. (2013). Long-term outcomes for the Child STEPs randomized effectiveness trial: a comparison of modular and standard treatment designs with usual care. *Journal of Consulting and Clinical Psychology*, *81*(6), 999-1009. doi: 10.1037/a0034200.
- Chu, B. C., Temkin, A. B., & Toffey, K. (2016). *Transdiagnostic mechanisms and treatment for children* and adolescents: An emerging field. Oxford: Oxford handbooks online.
- Clarke, A., Friede, T., Putz, R., Ashdown, J., Martin, S., Blake, A., ...Stewart-Brown, S. (2011).

 Warwick-Edinburgh mental well-being scale (WEMWBS): validated for teenage school students in England and Scotland. A mixed methods assessment. *BMC Public Health, 11,* 487.

 https://doi.org/10.1186/1471-2458-11-487

- Clark, D. M. (2011). Implementing NICE guidelines for the psychological treatment of depression and anxiety disorders: The IAPT experience. *International Review of Psychiatry*, *23*(4), 318-327. doi: 10.3109/09540261.2011.606803.
- CORC. Child Outcomes Research Consortium (CORC): www.corc.uk.net.
- Costello, E. J. & Angold, A. (1988) Scales to assess child and adolescent depression: checklists, screens, and nets. *Journal of the American Academy of Child & Adolescent Psychiatry, 27*(6), 726-737. doi: 10.1097/00004583-198811000-00011
- Craske, M. G. (2012). Transdiagnostic treatment for anxiety and depression. *Depression and Anxiety,* 29(9), 749-743. doi: 10.1002/da.21992
- Department of Health and NHS England (2015). Future in mind: Promoting, protecting and improving our children and young people's mental health and wellbeing. London: Department of Health and NHS England. Retrieved from

 https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/414024/C
 hildrens Mental Health.pdf
- Dong, Y. & Peng, J. (2013). Principled missing data methods for researchers. *Springerplus*, 2(1), 222. doi: 10.1186/2193-1801-2-222
- Edbrooke-Childs, J., Calderon, A., Wolpert, M., & Fonagy, P. (2015). *Children and Young People's Improving Access to Psychological Therapies: Rapid Internal Audit, National Report.* London:

 CYP-IAPT. Retrieved from https://www.ucl.ac.uk/drupal/evidence-based-practice-unit/files/pub_and_resources_project_reports_rapid_internal_audit_2015.pdf
- Fernandez, E., Salem, D., Swift, J. K., & Ramtahal, N. (2015). Meta-analysis of dropout from cognitive behavioral therapy: Magnitude, timing, and moderators. *Journal of Consulting and Clinical Psychology*, 83(6), 1108-1022. doi: 10.1037/ccp0000044.

- Foot, C., Naylor, C., & Imison, C. (2010). *The quality of GP diagnosis and referral.* London: The King's Fund. Retrieved from:

 https://www.kingsfund.org.uk/sites/files/kf/Diagnosis%20and%20referral.pdf
- Frith, E. (2016). *CentreForum Commission on Children and Young People's Mental Health: State of the Nation*. London: CentreForum. Retrieved from http://centreforum.org/live/wp-content/uploads/2016/04/State-of-the-Nation-report-web.pdf
- Gandhi, T. J., Sittig, D.F., Franklin, M., Sussman, A. J., Fairchild, D. G. & Bates, D. W. (2000).

 Communication breakdown in the outpatient referral process. *Journal of General Internal Medicine*, *15*(9), 626-631. doi: 10.1046/j.1525-1497.2000.91119.x
- Green, H., McGinnity, A., Meltzer, H., Ford, T., & Goodman, R. (2005). *Mental health of children and young people in Great Britain, 2004. A survey carried out by the Office for National Statistics on behalf of the Department of Health and the Scottish Executive.* Basingstoke: Palgrave Macmillan. Retrieved from http://content.digital.nhs.uk/catalogue/PUB06116/ment-heal-chil-youn-peop-gb-2004-rep2.pdf
- Goodyer, I. M., Reynolds, S., Barrett, B., Byford, S., Dubicka, B., Hill.J., ...Fonagy, P. (2017). Cognitive behavioural therapy and short-term psychoanalytical psychotherapy versus a brief psychosocial intervention in adolescents with unipolar major depressive disorder (IMPACT): a multicentre, pragmatic, observer-blind, randomised controlled superiority trial. *Lancet Psychiatry*, *4*(2), 109-119. https://doi.org/10.1016/S2215-0366(16)30378-9
- Guest, G., Bunce, A., & Johnson, L. (2006). How Many Interviews Are Enough? An Experiment with Data Saturation and Variability. *Field Methods*, *18*(1), 59-82.
- Hagermoser Sanetti, L. M., Collier-Meek, M. A., & Fallon, L. M. (2016). Fidelity with flexibility:

 Treatment acceptability and individualised adaptations of evidence-supported treatments.

 In S. Maltzman (Ed.), *The Oxford handbook of treatment processes and outcomes in*

- *psychology: A multidisciplinary biopsychosocial approach* (pp. 289-308). New York: Oxford University Press.
- Harper, B., Dickson, J. M., & Bramwell, R. (2014). Experiences of young people in a 16-18 Mental Health Service. *Child and Adolescent Mental Health*, *19*(2), 90-96.

 https://doi.org/10.1111/camh.12024
- Higa-McMillan, C. K., Francis, S. E., Rith-Najarian, L., & Chorpita, B. F. (2016). Evidence base update: 50 years of research on treatment for child and adolescent anxiety. *Journal of Clinical Child and Adolescent Psychology*, 45(2), 91-113. doi: 10.1111/camh.12024
- Hopkins, K., Crosland, P., Elliott, N., & Bewley, S. (2015). Diagnosis and management of depression in children and young people: summary of updated NICE guidance. *BMJ*, *350*, h824. doi: 10.1136/bmj.h824.
- House of Commons Health Committee (2014). *Children's and adolescents' mental health and*CAMHS: Third Report of Session 2014–15. House of Commons London: The Stationery Office

 Limited. Retrieved from
 - https://www.publications.parliament.uk/pa/cm201415/cmselect/cmhealth/342/342.pdf
- Ioannidis, J.P. (2005). Why most published research findings are false. *PLoS Medicine*, 2(8), e124.
- Khanna, M. S., & Kendall, P. C. (2010). Computer-assisted cognitive behavioral therapy for child anxiety: Results of a randomized clinical trial. *Journal of Consulting and Clinical Psychology,* 78(5), 737-745. doi: 10.1037/a0019739
- Kim-Cohen, J., Caspi, A., Moffitt, T. E., Harrington, H., Milne, B. J., & Poulton, R. (2003). Prior juvenile diagnoses in adults with mental disorder: developmental follow-back of a prospective-longitudinal cohort. *Archives of General Psychiatry, 60*(7), 709-717. doi: 10.1001/archpsyc.60.7.709

- Larsen, D. L., Attkisson, C. C., Hargreaves, W. A., & Nguyen, T. D. (1979). Assessment of client/patient satisfaction: Development of a general scale. *Evaluation and Program Planning, 2*(3), 197-207. https://doi.org/10.1016/0149-7189(79)90094-6
- McGorry, P., Bates, T., & Birchwood, M. (2013). Designing youth mental health services for the 21st century: examples from Australia, Ireland and the UK. *British Journal of Psychiatry, 202*(s54), s30-s35. doi: 10.1192/bjp.bp.112.119214.
- Memarzia, J., St Clair, M. C., Owens, M., Goodyer, I. M. & Dunn, V. J. (2015). Adolescents leaving mental health or social care services: predictors of mental health and psychosocial outcomes one year later. *BMC Health Services Research*, *15*, 185. https://doi.org/10.1186/s12913-015-0853-9
- Michelson, D., Sclare, I., Stahl, D., Morant, N., Bonin, E. M., & Brown, J. S. (2016). Early intervention for depression and anxiety in 16-18-year-olds: Protocol for a feasibility cluster randomised controlled trial of open-access psychological workshops in schools (DISCOVER).

 Contemporary Clinical Trials, 48, 52-58. doi: 10.1016/j.cct.2016.02.003
- Newby, J. M., McKinnon, A., Kuyken, W., Gilbody, S., & Dalgleish, T. (2015). Systematic review and meta-analysis of transdiagnostic psychological treatments for anxiety and depressive disorders in adulthood. *Clinical Psychology Review, 40,* 91-110. doi: 10.1016/j.cpr.2015.06.002
- Ng, M.Y., Eckshtain, D. & Weisz, J.R. (2016). Assessing fit between evidence-based psychotherapies for youth depression and real-life coping in early adolescence. *Journal of Clinical Child and Adolescent Psychology*, 45(6),732-748. doi: 10.1080/15374416.2015.1041591
- Persson, S., Hagquist, C. & Michelson, D. (2017). Young voices in mental health care: Exploring children's and adolescents' service experiences and preferences. *Clinical Child Psychology and Psychiatry*, 22(1), 140-151. doi: 10.1177/1359104516656722.

- Plaistow, J., Masson, K., Koch, D., Wilson, J., Stark, R. M., Jones, P. B., & Lennox, B. R. (2014). Young people's views of UK mental health services. *Early Intervention in Psychiatry*, 8(1), 12-23. doi: 10.1111/eip.12060.
- Pona, I., Royston, S., Bracey, C., & Gibbs, A. (2015). *Seriously awkward: How vulnerable 16-17 year olds are falling through the cracks*. London: The Children's Society. Retrieved from:

 http://www.childrenssociety.org.uk/sites/default/files/seriously_awkward_youth-at-risk_full_online-final.pdf
- Sclare, I., Michelson, D., Malpass, L., Coster, F., & Brown, J. (2015). Innovations in Practice: DISCOVER

 CBT workshops for 16-18-year-olds: development of an open-access intervention for anxiety

 and depression in inner-city youth. *Child and Adolescent Mental Health, 20*(2), 102-106. doi:

 10.1111/camh.12060
- Sclare, I. & Michelson, D. (2016). Innovations in CBT for young people: Improving access and outcomes for vulnerable youth. In: R.G. Menzies, M. Kyrios, & N. Kazantzis (Eds.).

 Innovations and Future Directions in the Behavioural and Cognitive Therapies (pp. 34-38).

 Samford Valley, Queensland: Australian Academic Press.
- Shirk, S. R., DePrince, A. P., Crisostomo, P. S., & Labus, J. (2014). Cognitive behavioral therapy for depressed adolescents exposed to interpersonal trauma: An initial effectiveness trial.

 *Psychotherapy, 51(1), 167-179. doi: 10.1037/a0034845.
- Smith, D., Roche, E., O'Loughlin, K., Brennan, D., Madigan, K., Lyne, J., ...O'Donoghue, B. (2014).

 Satisfaction with services following voluntary and involuntary admission. *Journal of Mental Health*, *23*(1), 38-45. doi: 10.3109/09638237.2013.841864.
- Southam-Gerow, M. A., Rodríguez, A., Chorpita, B. F. & Daleiden, E. (2012). Dissemination and implementation of evidence based treatments for youth: Challenges and recommendations.

- Professional Psychology: Research and Practice, 43(5), 527-534. http://dx.doi.org/10.1037/a0029101
- Stallard, P., Udwin, O., Goddard, M., & Hibbert, S. (2007). The availability of cognitive behaviour therapy within specialist child and adolescent mental health services (CAMHS): A national survey. *Behavioural and Cognitive Psychotherapy*, *35*(4), 501-505.

 https://doi.org/10.1017/S1352465807003724
- Steinberg, L. (2005). Cognitive and affective development in adolescence. *Trends in Cognitive Science*, *9*(2), 69-74. doi: 10.1016/j.tics.2004.12.005
- Stewart, R., Soremekun, M., Perera, G., Broadbent, M., Callard, F., Denis, M., ...Lovestone, S. (2009).

 The South London and Maudsley NHS Foundation Trust Biomedical Research Centre (SLAM BRC) case register: development and descriptive data. *BMC Psychiatry*, *9*, 51.

 doi:10.1186/1471-244X-9-51
- Weersing, V. R., Rozenman, M., & Gonzalez, A. (2009). Core components of therapy in youth: do we know what to disseminate? *Behavior Modification*, *33*(1), 24-47. doi: 10.1177/0145445508322629
- Weersing, V. R., Jeffreys, M., Do, M-C. T., Schwartz, K. T. G & Bolano, C. (2017). Evidence base update of psychosocial treatments for child and adolescent depression. *Journal of Clinical Child & Adolescent Psychology, 46*(1), 11-43. doi: 10.1080/15374416.2016.1220310
- Weisz, J. R., Kuppens, S., Ng, M. Y., Eckshtain, D., Ugueto, A. M., Vaughn-Coaxum, R., ...Fordwood, S.
 R. (2017). What five decades of research tells us about the effects of youth psychological therapy: A multilevel meta-analysis and implications for science and practice. *American Psychologist*, 72(2), 79-117. doi: 10.1037/a0040360.
- Wolpert, M., Jacob, J., Napoleone, E., Whale, A., Calderon, A., & Edbrooke-Childs, J. (2017). *Child-and Parent-reported Outcomes and Experience from Child and Young People's Mental Health*

Services 2011–2015. London: Child Outcomes Research Consortium (CORC). Retrieved from: http://www.corc.uk.net/media/1543/05052017 corc-report.pdf

Van Breukelen, G. J. (2006). ANCOVA versus change from baseline: more power in randomized studies, more bias in nonrandomized studies [corrected]. *Journal of Clinical Epidemiology,* 59(9), 920-925. doi: 10.1016/j.jclinepi.2006.02.0