

Sussex Research

Mapping of podoconiosis cases and risk factors in Kenya: a nationwide cross-sectional study

Hadley Matendechero Sultani, Collins Okoyo, Henry M Kanyi, Sammy M Njenga, Wyckliff P Omondi, Isabella Ayagah, Morris Buliva, Isaac Ngere, John Gachohi, Jacinta Muli, Melanie Newport, Kebede Kassaye

Publication date

09-06-2023

Licence

This work is made available under the Copyright not evaluated licence and should only be used in accordance with that licence. For more information on the specific terms, consult the repository record for this item.

Document Version

Published version

Citation for this work (American Psychological Association 7th edition)

Matendechero Sultani, H., Okoyo, C., Kanyi, H. M., Njenga, S. M., Omondi, W. P., Ayagah, I., Buliva, M., Ngere, I., Gachohi, J., Muli, J., Newport, M., & Kassaye, K. (2021). *Mapping of podoconiosis cases and risk factors in Kenya: a nationwide cross-sectional study* (Version 1). University of Sussex. https://hdl.handle.net/10779/uos.23481917.v1

Published in

American Journal of Tropical Medicine and Hygiene

Link to external publisher version

https://doi.org/10.4269/ajtmh.21-0248

Copyright and reuse:

This work was downloaded from Sussex Research Open (SRO). This document is made available in line with publisher policy and may differ from the published version. Please cite the published version where possible. Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners unless otherwise stated. For more information on this work, SRO or to report an issue, you can contact the repository administrators at sro@sussex.ac.uk. Discover more of the University's research at https://sussex.figshare.com/

County/SubCounty	No. of	No. of	LF prevalence	
	villages	participants	No. positive	%(95%CI)
Overall	48	6,228	9	0.1 (0.1-0.3)
Demographics				
Sex				
Male	48	2,847 (45.7%)	5	0.2 (0.1-0.4)
Female	48	3,381 (54.3%)	4	0.1 (0-0.3)
Age group				
<10	48	383 (6.2%)	0	0
10-17	48	932 (14.9%)	1	0.1 (0-0.7)
≥18	48	4,913 (78.9%)	8	0.2 (0.1-0.3)
County/SubCounty				
Baringo			0	0
Baringo South	2	201	0	0
Tiaty	2	200	0	0
Busia			0	0
Teso North	2	221	0	0
Homabay			0	0
Homabay Town	2	199	0	0
Makueni			0	0
Kibwezi East	2	208	0	0
Kibwezi West	2	203	0	0
Marsabit			1	0.2 (0-1.5)
Moyale	2	147	1	0.7 (0.1-9.0)
North Horr	2	199	0	0
Saku	2	201	0	0
Meru	2	205	1	0.3 (0-2.6)
Igembe Central	2	205	0	0
South Imenti	2	157	1	0.6 (0-9.2)
Nakuru	2	202	0	0
Nalvasna	2	203	0	0
Subukia	2	201	0	0
Narok	2	140	1	0.7 (0.2-2.4)
Narok East	2	140	1	0.7 (0.2-2.4)
Slaya	2	104	0	0
	Z	194	0	0
	n	205	0	0
Turkene	Z	205	0	
Turkana North	2	160	1	0.3(0-2.5)
Turkana Most	2	207	1	0.0 (0.1-7.0)
	2	207	1	
Tarbai	2	206	1	0.5(0.1-5.5)
West Pokot	2	200	0	0.5 (0.1-5.5)
Kacheliha	2	207	0	0
	2	207	0	0
Taveta	2	605	0	0
Tana River	۷	005	1	0 2 (0 1-0 6)
Bura	2	610	1	0.2(0.1-0.0)
Galole	2	529	1 2	0.2 (0-1.2)
Garsen	2	619	- 1	0.2 (0-1.1)

Supplemental Table S1: Number of villages and participants surveyed and LF prevalence per subcounty in 15 counties in Kenya